## Relationship Between Sub-Period and Total Period Regression

Pradeep Apte

The problem with which we deal here may be stated as follows. Suppose we have two sample-data sets, each pertaining to successive periods or groups. Let these be denoted as  $[\mathtt{X}_1,\mathtt{Y}_1]$  with  $\mathtt{n}_1$  observations and  $[\mathtt{X}_2,\mathtt{Y}_2]$  with  $\mathtt{n}_2$  observations.

We first compute regression slope-coefficient using these two data sets separately and denote these as  $\beta_1$ , and  $\beta_2$  for the first and second samples respectively.

Now combining these two sub-period data sets we have yet another sample i.e.  $\{(X_1, X_2), (Y_2, Y_2)\}$ . With  $n_1 + n_2$  number of

observations and calculate slope coefficient again, denoted by  $\beta.$  What relationship does exist between  $\beta_1,\ \beta_2$  and  $\beta$ ?

Relevance of such a problem can be easily imagined. Research workers often handle the situations where the given data is divided into sub-groups and regression parameters for the sub-groups taken separately and these subgroups taken or pooled together are calculated. For example; growth rates, incremental capital output ration (ICOR) consumption function etc. are often estimated, using regression, for, sub-periods as well as for the total period. Generally, (perhaps due to experience with arithmatic averages), one expects that growth rate or ICOR, for the total period will be some-where between the subperiod parameter estimates. [1] But this does not always happen and a seemingly paradoxical situation emerges, where the parameter of the 'total' period lies entirely above or below the subgroup parameters. Such a result needs some straightforward explanation which standard text-books on statistics do not often provide. [2]

A similar question has been dealt with by Placket (1950).
[3] But the question was posed, somewhat differently, as follows. If we have a sample which is initially given and later