SERVANTS OF INDIA SOCIETY'S LIBRARY, POONA 4.

BULLETIN No. 109 OF 1921


THE CROPS OF THE BOMBAY PRESIDENCY

THEIR GEOGRAPHY AND STATISTICS

ВY

P. C. PATIL, L.Ag.


DEPUTY DIRECTOR OF AGRICULTURE, CENTRAL DIVISION



Dhananjayarao Gadgil Library

(| 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 | 111 |

GIPE-PUNE-045754



FOREWORD.

ANY attempt to provide, for the general public, a means of obtaining exact knowledge with regard to the area and distribution of the crops of Western India is the direct result of the series of famines which afflicted the country during the second half of the nineteenth century. The report of the famine commission which sat after the famine of 1877 made, for the first time, a plea for a thorough knowledge of the existing agricultural conditions as one of the great pre-requisites for being able to meet the situation created by a long drought. And as a result, the famine relief code, when prepared, ordered that a statistical account with maps should be prepared for each district. This is the origin of the Statistical Atlas of the Bombay Presidency of which two editions have appeared hitherto, the first in 1888 and the second in 1906, and which has formed by far the most accessible source of information hitherto as regards the distribution of crops over the districts of Western India.

But something more than this has been needed for those who are interested not only in meeting drought and fighting famine, but also in trying to improve the general agricultural conditions. We require not only the general statistical tables which are published in the Statistical Atlas, or which are brought up to date each year in the Annual Season and Crop Report, but also we need to know the distribution of the crops inside the districts themselves, their relationship to the rainfall, to the soil, or to the supply of irrigation water, and the way in which one crop is replaced by another when the climate or the soil is altered even to a comparatively small degree. This is what Mr. Patil has endeavoured to provide in the present bulletin. By a series of maps and tables, specially prepared for the purpose, he has succeeded in showing, very vividly, the distribution of the crops of the country, their relation to the various factors in evidence and to one another. By carrying his figures and his maps down to the subdivisions (talukas) of the districts, he has been able to show these relationships more clearly than has ever been done before.

The figures given and illustrated are for a single year, 1914-15. This represents a normal year perhaps better than any other recent one that might have been chosen, and is one in which the figures had not been affected seriously either by war conditions or by the rise in prices which has been the most striking economic feature of the last five years. It may be possible to issue another bulletin on similar

lines when agricultural arrangements have again become stable under the new conditions, but that is a matter for the future. At present the figures, diagrams, and maps now given do show a very definite relationship between the agricultural practices and the other factors considered, which it is of extreme importance to recognise in studying the development of the agriculture of Western India.

HAROLD H. MANN,
Acting Director of Agriculture,
Bombay Presidency.

Poona, May 24, 1921.

CONTENTS.

						:	Page
	Foreword	••					i
	Table of Cont	ents	••				iii
I.	Introduction	••	••	••			1
	Table of Aver	rage Rainfa	all of Typic	al Centres	••		4
11.	Rainfall in th	e Bombay	Presidency	7			6
III.	Rocks and So	ils of the l	Bombay Pr	esidency	••		9
IV.	The Crops of	the Bomba	y Presider	ıcy			13
	General Stat	ement No	. I of the	Crops of th	e Bombay	Presi-	14
	General Stat	ement No	. II of the	Crops of t	he Bombay	Presi-	
	dency	~	• • •	••	• •	• •	24
	Jowar (Andre			••	••	• •	25
	Bajri (Pennis	etum typho	rrdeum)	••	••	:•	26
· VII.		••	••	••	••	••,	27
	Wheat	••	••	••	••	• •	30
	Cotton	••	••	••	••	• •	31
	Sugarcane	••	••	• •	••	• • •	33 🎉
	Potato	••	••	• •	•	••	35
	Tobacco	••	••	••	••	• •	36
	Oil Seeds	••	••	••	••	• •	37
	Fruits	••	••	••	••	• •	45
	Spices and C				••		48
XVI.	Vegetables as	nd Root C	rops	••	••		49
XVII.	Yields of Cro	p s	••	••	••		51
VIII.	Irrigation	••	••	••	••		57
XIX.	Live Stock	••	••	••	••	••	61 •
	•		PART II	.			
rops o	of each Distric	t of the B	ombay Pre	sidency		••	71
		A	PPENDIX	ζ,			
lossar	y of Botanica	l, English	and Vernac	ular names		••	129
lossar	v of Technica	l Agricultu	ral terms		••		134

THE CROPS OF THE BOMBAY PRESIDENCY, THEIR GEOGRAPHY AND STATISTICS.

I.—INTRODUCTION.

The purpose of the following account of the geography and statistics of agriculture in the Bombay Presidency is to show the distribution of the crops in the area dealt with, explaining as far as possible the relation between the crops and the climatic, soil, and economic conditions which are responsible for this distribution. It is not possible within the limited space in the present bulletin to do more than touch on the factors which govern the production and distribution of crops in general. The statistical data are collected from various sources, but chiefly from the annual season and crop reports, and from the original returns from each subdivision or taluka, which are received in the office of the Director of Agriculture. In the case of sugarcane and fruits, information had to be specially collected from the officers in charge of each taluka (Mamlatdars). In the matter of rainfall statistics, the information available in the rainfall returns issued by the Director of Agriculture has been supplemented by materials collected from raingauges maintained by the Public Works Department in the Nasik, Ahmednagar, Poona, and Satara districts. The material collected from these various sources is worked out and presented in statements and in specially prepared maps and graphs in such a form as to show as vividly as possible the distribution of the crops and to enable conclusions to be drawn for any taluka or district.

The Presidency maps show the density and distribution of crops, each dot in these maps representing a definite area under the crop. The area covered by fairly large native states, for which statistics are not available, are shown by cross lines. The graphs accompanying the Presidency maps show the relative importance of each district in the production of a particular crop: the graphs at the bottom of the district maps visualise the relative importance of the various talukas in growing these crops.

The acreage statements show the areas of each important crop and how these are distributed. At a glance one can find how the land occupied by a particular plant or class of plants compares with that of any other, and in the same statement is found how a particular crop in any district is distributed in the different talukas.

Crop Notes.—A short account of important crops or groups of crops accompany the maps. The notes on these crops are not intended to be exhaustive. They, however, indicate the importance,

distribution and use of each, and also their limitations. Economic, historical, and other facts bearing on their distribution are mentioned.

In addition to maps and statements with regard to crops, the following are included, with explanatory notes:—

- (a) A map showing the distribution of rainfall.
- (b) A statement showing the distribution of live stock.
- (c) A statement showing the relative areas under irrigation in the various districts.
- (d) Data regarding the normal and average outturns of crops, including fodder crops.

Explanation of statistical statements, maps, and graphs.—The areas under different crops are taken for the year 1914-15. This was a normal year and the area under the different crops had not been disturbed either by famine or by the war which commenced soon after the crops were sown. Moreover, it was the last year in which the crop returns were received in the office of the Director of Agriculture by talukas. Since 1915-16, such returns have only been furnished by districts, though these are as a matter of fact far too large units for statistical bulletins and crop distribution maps. The area for fruits is taken for the year 1917-18 because of the recent increase in the amount of land under certain kinds of fruit. The area for sugarcane is taken for 1914-15, but side by side the area for 1917-18 is also given as, owing to the expansion of canal irrigation, the amount of land under sugarcane is steadily increasing.

• The number of cattle is taken from the quinquennial census of the year 1915-16 as the next quinquennium shows a considerable diminution due especially to the fodder famine of 1918-19.

Complete Presidency maps are only given for the important crops. iA crop may, however, be important because of the large area which it occupies, or because it is a staple food crop, or it may be important because it is a crop of high value and hence of greater economic importance than its area would indicate.

In the Presidency crop maps, each dot represents 1,000 acres except in the case of sugarcane, when each dot only represents 10 acres. In the same way in the district maps, for the first six crops of the district, a dot represents 1,000 acres, and for the remaining three crops illustrated (except in a few cases) one dot represents 100 acres. Graphs and dots are shown on a uniform scale and they show the relative importance of each taluka or district for all crops on the same basis. In the small maps drawn for fruits and oilseeds, a dot is made to represent different areas, but on each map the value of a dot is given.

It may be noticed in this place that the areas given in this bulletin are the areas of the tracts which regularly report to Government. In most districts there are some portions which for one reason or another do not furnish returns, and for these estimates are made every year on the basis of the tracts surrounding them. The matter is not important except in the Ratnagiri district, and except in that one case, the actual area reported is used in framing the maps and tables. The reason is that the estimates are, after all, estimates, and their accuracy is doubtful, while the error introduced by ignoring them is very small. But in the Ratnagiri district the case is different. About half the total non-reporting area in the Bombay Presidency is in that one district and there amounts to 533,000 acres, as most of the so-called 'khoti' villages do not supply statistics. In this case therefore the question is important, for the non-reporting area amounts to about three-fourths of the cultivated area in the district, and in addition, the crops there grown (rice, nagli, and cocoanuts) are not produced to a very large extent in the greater part of the Bombay Presidency. For this reason, for Ratnagiri, two sets of figures are given in the statistical tables, and the estimated area is taken into account in preparing the maps showing the distribution of these crops.

Acknowledgments.—I have to thank Mr. G. F. Keatinge, C.I.E., the Director of Agriculture, for many useful hints which he has given during the preparation of this bulletin, and my thanks are due to Dr. H. H. Mann, late Principal of the Agricultural College, Poona, and now acting Director of Agriculture, for other help. He went over the manuscript and attended to the proofs and press work as I was away in the United States of America while the book was in press. My thanks are due also to my friend Mr. S. G. Mutkekar, B.Ag. (Bombay), M.Sc. (Mass.), for much assistance. Mr. K. R. Tilak, the artist of the Department of Agriculture, Bombay, drew the maps for me, and Mr. N. V. Bhagwat, B.Ag., and others helped me in calculating the figures given.

As has already been stated, the principal sources of information have been the annual season and crop reports issued by the Government of Bombay and the taluka returns of crops kept in the office of the Director of Agriculture. But in addition to these, I have used the Report of the Irrigation Commission, rainfall returns from various sources, and many other Government publications in the preparation of the data presented in the following bulletin.

Average rainfail, of typical centres in each rainfall tract.

	٥		1			Mon	thly Norm	al Rainfe	ılı.		,•				*	
	Station.	District.	January.	Febru- ary.	March.	April.	May.	June.	July,	August.	Septem - ber.	October.	Novem- ber.	Decem- ber.	Annual Normal Rain- fall.	
		•							1]		l .			
		Ì					}		places.		1		,		ŀ	
3	Karwar .	North Kanara .	0.08	0.01		0.21	3.08	87 80	89.12	20.28	10.58	5.48	1.47	0.21	118.36	
2	Ratnagiri .	. Ratnagiri .	0.10	0.02	0.04	0.08	1.09	28.12	83.45	18.09	11.44	8.24	0.71	0.08	96.47	
8	Malvan .	. Do	0.08	0.08		0.19	1.68	25 · 19	29.04	18.95	9-86	2.56	0.96	0.12	83.62	
4	Alibag .	. Kolaba .	0.07	0.03	0.08	0.11	0.67	23.42	29.47	16.39	13.16	2.26	0.42	0.05	86.08	
5	Umbergaon .	Page	0.03	0.09	0.04	٠	0.18	13.68	24.61	13.12	9-42	1.07	0.29	0.08	62.56	
6	Jalaipur .	. Surat .	0.08	0.08	0.02		0.02	9.12	20.93	8.83	6.18	0.98	0.88		46.56	
					Ì	Mallad	or	Maval	tracts.				ļ			
7	Yellapur .	. North Kanara .	. 0.08	0.08	0.25	1.25	2.45	20.37	87.80	21.37	7.11	5.24	1.27	0.83	97.55	
8	Malcompeth .	. Satara .	. 0·i1	-0.02	0.18	1.40	1.62	46.26	110.28	75.09	29.11	5.83	1-48	0.81	271 62	
9	Lonavla .	. Poons .	. 0.05	0.03	0.02	0.26	0.75	27.86	68.85	46.88	19.62	8.46	0.89	0.17	168-84	
10	Peint .	. Nasik .	0.04	0.09	0.01	0.02	0-50	11.99	33.71	25.40	10.66	2.02	0.48	0.08	84.98	
11	Bansda .	. Surat Agency .	. 0.04	0.18	0.04	0.08	0.18	11-87	83.42	22.13	11.92	1.18	0.47	0.07	81:51	
]	Tr	act lying	between	Maval	and scan	ty rainfa	lltract.				İ	
12	Savanur	. Savanur State .	. 0.06	0.02	0.12	148	2.54	4.24	4.74	8.64	8.20	4.78	1.08	0.55	26.40	
18	Shirala .	. Satara .	. 0.08	0.04	0.17	1.04	1.71	5.95	12.10	6949	4.01	4.14	1.01	0.21	36.95	
14	Gadhingiaj .	. S. M. C. States,	0.09	0.05	0.43	1.86	2.16	6.74	11-57	6-65	4.05	3.99	1.19	0.89	39-17	

II.—RAINFALL IN THE BOMBAY PRESIDENCY.

Any map of the rainfall of the Bombay Presidency will indicate at once the extraordinary variations which occur in different parts of its comparatively limited area,— differences which are perhaps more striking than in any other part of India.

The Bombay Presidency receives its rain mainly between June and December, the greater part of the fall being supplied by the South-West Monsoon between June and October, supplemented in a few parts by later rain in November and December. The South-West Monsoon touches North Kanara, the southernmost district, first and works its way up towards Gujarat. As it advances, the amount of precipitation for which it is responsible decreases from south to north along the coast. Bhatkal, at the extreme south on the coast of Kanara receives 150 inches of rainfall. Ratnagiri about two hundred miles north, receives 96 inches, Alibag another hundred miles in the same direction has only 86 inches, while Umbergaon in the north of the Thana District receives only 62 inches.

The high range of the Western Ghats, inasmuch as it arrests the surcharged clouds, receives more rainfall than the coast line over which the clouds come. Thus Ratnagiri, Guhagar, Alibag and Mahim being on the coast line receive 96 inches, 92 inches, 86 inches and 62 inches of rainfall respectively, while Sangameshwar, Chiplun, Lonavla and Igatpuri being on or near the Ghats in the same latitude as the above places respectively receive 146 inches, 141 inches, 168 inches and 130 inches respectively. It will thus be seen that the western face of the Western Ghats arrests much of the rain. The line of the Western Ghats, which runs nearly parallel to the coast and the ramifications of the Ghats on the eastern side, cause considerable variation in the amount of rain received in the Deccan and Karnatak districts, as the current of the south-west monsoon has to pass across the Ghats. The Deccan districts have a slight slope to the east, away from the Western Ghats, and most of the rain which crosses the Ghats falls in the uplands. The rainfall lines on the map show how the rainfall decreases eastward and, to a less extent, westward from the summit of the Ghats. Khandesh, and especially East Khandesh receives a little additional rainfall from the east, from wind currents originating in the Bay of Bengal.

By about September the force of the south-west monsoon decreases fast, and, soon after, a series of storms arriving apparently from the north-east supplies rain to Ahmednagar, Sholapur, Bijapur and the eastern parts of Dharwar and Belgaum. Sometimes these storms spread further west, but as a whole they form perhaps the most doubtful factor in the annual rainfall.

The Bombay Presidency can be divided into the following tracts on the basis of rainfall:—

- (1) The three Konkan districts (Ratnagiri, Kolaba and Thana) Kanara, and a narrow strip of about fifteen or twenty miles on the crest of the Western Ghats from Kanara to the Nerbudda in the western part of West Khandesh. This area receives more than 80 inches of rainfall annually.
- (2) The tract called mallad in Belgaum, mura in Satara, maval in Poona, and the konkani talukas of the Nasik district. This is a narrow strip of a width of fifteen or twenty miles to the east of the areas included under (1). It receives from thirty-five to fifty inches of rain per annum, and in some places a little more.
- (3) The so-called desh tract, comprising the eastern parts of the Belgaum, Satara, Poona and Nasik districts, together with the whole of the Ahmednagar, Sholapur, Bijapur and almost the whole of the Dharwar district. These areas receive from eighteen to thirty inches of rainfall annually.
- (4) Khandesh, consisting chiefly of the Tapti valley and surrounding areas in the Bombay Presidency, has rainfall features of its own. It receives from twenty to thirty inches of rain per annum.
- (5) Gujarat, or the coast tracts, with the flat land behind them, to the north of the Konkan, receives from twenty to forty inches of rainfall per annum on the average, the amount tending to become smaller as one proceeds further north.

We will deal shortly with each of these tracts in succession.

(1) The heavy rainfall tract.—This tract receives, as has already been noted, eighty inches or more of rain annually, the stations which are actually on the crest of the Western Ghats recording very large quantities up to from 250 to 300 inches per annum. As one proceeds to the north from Bhatkal, at the extreme limit of the coast of the Bombay Presidency to the South, as far as Umbergaon in the Thana district which may be taken as the northern limit of the tract under discussion, the total annual rainfall tends continually to decrease from 150 to 62 inches. Owing to the mountainous nature of the country and the heavy rainfall during only about three or four months of the year (June to September), this tract is a very rugged one, is cut up into narrow valleys and the country is highly eroded. • Forest occupies a large proportion of the area in Kanara and along the whole length of the Western Ghats themselves. Bare rock (laterite) occupies large areas in the Ratnagiri district. With the exception of these large laterite plateaux there are very few large flat areas of land except on the sea coast and near the tidal creeks. The hill sides in much of this tract are carefully terraced for cultivation.

The soil is very variable, being formed in various parts of the tract from laterite, trap, or, in the south, granite and schists. On the whole it is shallow and is not retentive of water. As a result, although this tract receives large quantities of rain it grows no crop in the *rabi* season, except in the few level areas with deep soil. Even in the *kharif* season, if the rains hold off for a week or two the crops suffer considerably.

(2) The Mallad tract.—This tract receives from thirty-five to fifty inches of rain per annum. It lies immediately to the east of that last dealt with, and usually has an elevation above the sea of two thousand feet or more. That is to say, the general level of the plateau is about two thousand feet, while the strip is crossed by ranges of hills many of which are a thousand feet higher. The strip of country is in fact mountainous though not as much so as that last dealt with, while it is not nearly so level as the desh further to the east. The distribution of the rain is very similar to that in the heavy rainfall tract already dealt with.

The soil, though not usually very deep, is more retentive of water than that in the former tract, and much, in the valleys, grows both *kharif* and *rabi* crops. It is formed chiefly from trap rock, except in the extreme south where granites and schists appear, and in a few parts, of the Belgaum district especially, where laterite is found. Most of the good areas of land are levelled and embanked, so as to be useful for rice cultivation, or to avoid further erosion.

(3) The Desh tract.— To the east of the mallad tract lies what is usually called the desh, and in this region the land becomes much more level, while the rainfall is much smaller and more uncertain as one proceeds to the east. The rainfall, in fact, varies from a little over thirty inches on an average in the western part of the area to an average of about twenty-two inches in the eastern tract. The latter is that portion of the Bombay Presidency which is chronically liable to drought and hence to famine. In these eastern parts, moreover, the chief dependence is on the rain in August and September, and hence the people mainly rely on rabi crops—rabi jowar, wheat, oilseeds, or cotton planted late.

For the most part this tract consists of level plateaux of fair extent, though there are low rocky hills in view throughout the area. The land is often, however, badly cut up near the rivers and very little effort has been made to prevent scouring or eroding of the soil. The soil is usually deep and retentive, becoming more so as one proceeds towards the east.

(4) The Khandesh tract.—The Khandesh tract consists almost entirely of the valley of the river Tapti, and the surrounding country. At this latitude the Western Ghats are no longer the formidable barrier that they are further south, but on the other hand the Satpura range, which forms the northern boundary of Khandesh and of the Bombay Presidency, intercepts a great deal of the rain which is passing north in the south-west monsoon current. At the same time, the country is open to the east and a certain amount of rain from the Bay of Bengal current of the monsoon reaches Khandesh. As a result of these features, the rainfall of Khandesh, though only from twenty to thirty inches annually in amount, is much more certain than in the desh country further south. It is, however, almost limited to the south-west monsoon, and little late rain in September or October can be relied upon.

The soil of Khandesh, especially in the east, tends to be very deep and retentive wherever it is level and erosion is not great. As a result, in spite of the absence of late rain, rabi as well as kharif crops are grown, more particularly in the east. The general level of the country in Khandesh is about a thousand feet lower than in the desh country to the south, but it slopes up to the latter on the south and up to the Satpura range on the north.

(5) The Gujarat tract.—The level plain which stretches from the coast to the east and north from the confines of the Thana district, is known as Gujarat. It is little elevated above the sea level, and is mostly flat. The rainfall, which in the extreme south is from fifty to sixty inches annually, soon becomes not more than forty inches at • Surat and less and less from that point as one goes north. It is, more over, entirely a rainfall of the south-west monsoon, and it stops as a rule in early September. After this time no appreciable amount of rain falls until the following June.

The southern part of Gujarat consists of stiff black soils, formed from trap. But from Broach northward this soil is gradually replaced by large alluvial plains of sandy loam. The district known as the Panch Mahals does not properly belong to Gujarat, and lies among the Satpuras and is hence hilly, the hills being composed of granite and gneiss, with deep or shallow soils in the valleys according to the configuration of the country.

The map presented shows the general distribution of the total rainfall, the differences in the average annual fall being shown by differences in shading.

III,—ROCKS AND SOILS OF THE BOMBAY PRESIDENCY.

The crop characteristics of an area like the Bombay Presidency are determined not only by climatic considerations but also by the character of the soils, and these are again determined by the rocks or other sources from which they are derived. In this respect the geological features of the Bombay Presidency are simple. The greater part in fact of the area with which we are dealing in the present bulletin consists of what is probably the largest outbreak of volcanic rock in the world, the well-known Deccan trap. Except in the extreme south, and in the Panch Mahals in the north-east, with the alluvial plains of Northern Gujarat, and a few islands of sandstone and the soils derived from it in the south Deccan, the whole of the Presidency owes its characteristic features to the Deccan trap. The bare and rugged country of the Konkan is a trap country: equally so is the treeless hill country of the Deccan. From it is derived the famous black cotton soil or regur which occupies so much of the Deccan cultivated area.

From the trap, too, is chiefly derived the laterite, which is a feature of the Konkan and the *mallad* area, particularly toward the south. This is a porous argillaceous rock, generally red in colour owing to impregnation with peroxide of iron, which gives rise to light dry hungry soils in the regions named.

Granite and gneiss appear in the extreme south and form the prevailing rocks in the Kanara district and also in parts of the adjoining districts of Dharwar and Belgaum. These same rocks reappear in the Satpura range to the north and form the characteristic feature of the geology of the Panch Mahals.

The sandstone of the Belgaum, Bijapur, and a part of the Dharwar districts form an island in the midst of the surrounding trap area. But the soils derived from it give a characteristic cropping which is considerably different from that of the surrounding regions.

The soils characteristic of the various parts of the Bombay Presidency are as follows:—

- (1) The Northern Gujarat Tract.—This tract, consisting essentially of the districts of Ahmedabad and Kaira, is really a part of the great alluvial areas of northern and central India. The soil is formed from primitive and metamorphic rocks of Central India and Rajputana, and is usually level and deep. It varies from what is little more than drift sand in some parts of Ahmedabad to a splendid loam in parts of Kaira. The colour is light fawn to rich brown. On the whole it is a rich soil if only it is supplied with sufficient water, and in the better portions it is fairly retentive. Kharif crops are the mainstay of the people, but the more retentive lands grow rabi crops and especially wheat. It lends itself admirably to irrigation, and much of the best well irrigation in the Bombay Presidency is on this land.
- (2) The Southern Gujarat Tract.—This tract is also essentially alluvial, but the source of the alluvium is quite different from

that in the previous case. In the Surat and Broach districts, which form this tract, the main source of the alluvium is trap rock, and granite and gneiss only to a completely subordinate extent. The soil is usually deep and black in colour, and it has all the characteristics of black cotton soil. It is very retentive of moisture, and cracks deeply in the dry parts of the year. This last feature is not, however, found in the bhatha or riverine lands or in the gorat or former riverine lands. Except for the last two classes of land, this area is not considered as suitable for irrigation. The bhatha and gorat soils, however, grow large quantities of irrigated garden crops.

Large tracts of this black soil area, especially in the southern part, grow *kharif* crops such as *jowar*, cotton, and where the rainfall is sufficient, rice. In the northern part of the area *rabi jowar* and wheat find a place in rotation with cotton.

- (3) The Panch Mahals Tract.—This district, which lies among the Satpura Hills and is considerably different from any other district in the Bombay Presidency, is an undulating area where the predominant rock is granite or gneiss. The soil in the higher lands is shallow and light coloured, while the lowlying areas are composed of a deep, rich, dark coloured loam, retentive of water and capable of giving two crops a year. These crops are usually maize in the kharif season and wheat in the rabi season. The standard of cultivation in this district is, however, low.
- (4) The Khandesh Tract.—The soils of Khandesh are essentially derived from trap rock, and are hence of the black cotton soil type. Their depth and retentiveness, however, varies a good deal according to their distance from the Tapti river. Along the flat areas near the Tapti river the soil is deep and retentive and the characteristic crops are cotton, jowar and wheat. Further from the river the soil is not so deep, but as the rainfall is fairly reliable cotton is commonly grown in rotation with jowar or bajri. The soils, both deep and shallow, are well adapted for irrigation and grow large areas of chillies and plantains in rotation with dry crops.
- (5) The Deccan Tract.—This tract which comprises the districts of Ahmednagar, Nasik, Poona, Satara and Sholapur consists essentially of soil derived directly from trap rock. It can be divided into three strips more or less parallel with the Western Ghats:—
 - (a) The high lying strip along the Western Ghats is hilly and covered with jungle, having small hamlets scattered wherever tillable soil is to be found. This area receives

heavy rainfall and grows *kharif* crops such as rice and millets. The soil is usually shallow and porous.

- (b) The next strip to this is less hilly than the last and contains a good deal of tillable land. It generally receives sufficient and certain rain, and grows both *kharif* and *rabi* crops, such as rice, *jowar*, peas in the kharif season, and *jowar*, gram and wheat in the rabi season. Irrigated crops are also grown, and in some parts, notably in the Nasik district, local irrigation dams enable the people to grow rice, sugarcane and wheat under irrigation. The soil in this part of the country is still shallow, but much deeper than in the strip last dealt with.
- (c) The next strip is the desh and the principal characteristics of this area have been already described. The soil is deep, black and retentive on the whole except on the high lands and hills.
- (4) The Karnatak Tract.— This tract which lies to the south of that last dealt with and which includes the districts of Belgaum, Bijapur and Dharwar, consists in the northern portion of the three strips whose characteristics were detailed under the last head. To the south of this there are however differences, as we pass from the trap area into the transition tract between the trap rocks to the north and the metamorphic rocks to the south. This transition tract is rich, and, as the late rainfall of August and September is more certain than it is further to the north, rabi crops can be grown with more assurance. This is the great area for rabi cotton, and there is also found a very large development of wheat in the area in question. This area, occurring in the Dharwar district contains some of the deepest and most retentive black soils in the Bombay Presidency.
- (5) The Konkan Tract.—This tract, which lies wholly between the coast and the Western Ghats, contains the districts of Thana, Kolaba, Ratnagiri and North Kanara. It is hilly, with shallow or porous soils everywhere, and except near the coast there is little level land. The greatest amount of level land occurs in the northern districts, Thana and Kolaba. The common rock, except in the north, is latterite, beneath which trap can usually be found, though this latter is replaced by granite in the south towards Vengurla. The soils are usually divided into rice lands, garden lands, and varkas lands. The rice land may be either malla or low-lying rich land, or kuryat or land lying on the slopes of low hills which are terraced in order to hold water and prevent erosion, or kharvat, that is to say, land reclaimed from the sea. The garden lands are so called from the fact that they have

command of well or other water for irrigation, and have been levelled and otherwise prepared for garden crops, which in this case include betel palms, cardamoms and pepper in the Kanara district, and fruits, ginger and sugarcane over most of the tract.

The varkas, or upland lying on the slopes or tops of hills and plateaux in the Konkan, consist of shallow and poor soil, which is usually cultivated by a system of long fallows, with crops of millets of various kinds and of niger seed respectively in the years in which they are cultivated at all. They are, in spite of an assured rainfall, among the most barren lands in the Bombay Presidency.

IV.—THE CROPS OF THE BOMBAY PRESIDENCY.

The tables printed on pages 14—19 give a general statement as to the area which is devoted to the several crops and classes of crops in the various districts of the Bombay Presidency. They are based on the records of a single year, namely, 1914-15, with certain corrections which will be afterwards noted for progressive changes which have occurred with several crops. This year has been chosen as representing the normal distribution of crops before the war and the resulting economic disturbances in the world had occurred.

	•		14				
v ş			GENERA	l State	MENT]	No. I or	CROP
							Ceres
		}	Jos	Var.	·	Bajri	
	Names of the Districts.	Kharif.	Rabi.	Total.	P.C.		
		Area.	Area.	Area.		Area.	P. C.
1	2	В	4	5	6	7	8
, 1	Ahmedabad .	259,648		259,648	3.99	163,703	3.1
2	Kaira .	22,725	766	23,491	0.36	152,916	2.0
8	Panch Mahals .	8,266		8,266	0.13	44,976	0.8
4	Broach .	3,873	136,491	140,364	2.15	7,915	0.1
· 5 ·	Surat .	88,654	12,863	101,517	1.55	5,648	0.15
6	West Khandesh .	74,483	36,346	110,829	1.72	359,198	7.10
7	East Khandesh .	321,568		321,568	4.95	326,579	6.8
8	Nasik .	38,233	38,803	77,036	1.18	856,514	16-6-
9	Ahmednagar .	5,342	707,845	712,687	10.95	1,068,812	20.8
10	Poona .	121,823	510,639	632,462	9.74	728,935	14.2
11	Sholapur .	.}	1,128,601	1,128,601	17:35	4,08,516	7.8
12	Satara .	248,382	275,421	523,803	8.02	421,740	8.2
13	Belgaum .	813,309	224,408	537,717	8-27	195,863	3.84
14	Bijapur .	358,127	953,553	1,311,680	20-16	373,594	7 21
15	Dharwar	446,661	165,956	612,617	9.42	9,187	0.11
16	Thana .	690		690	0.01	8	
17	Kolaba .						
4 18	Ratnagiri .					20 20	::
19	Kanara .	664		564	0.008		
	Total	2,312,348	4,191,192	6,503,540	100.00	5,119,119	100.0

15

OF THE BOMBAY PRESIDENCY.

_	
and	Duises.

Rice	θ.	Whea	t	Other food-grains and pulses.								
				Barley.	Maize.	Ragi.	Gram.	Other food grains.	Total area.			
Area.	P. C.	Area.	P. C.	Area.	Area.	Area.	Area.	Атеа.				
9	10	· 11	12	18	14	15	16	17	18			
68,468	4.07	145,815	9.5	11,398	4,194	4,372	24,798	54,711	737,107			
89,222	6.87	11,398	0.7	1,662	421	61,657	10,174	132,138	483,079			
69,101	4-12	13,960	0.8	3,192	111,028	16,118	58,978	69,008	389,627			
13,279	0.79	69,678	4.8	••	12	198	2,691	59,216	293,348			
96,267	5-87	28,229	1.8	1	8	16,181	1,888	57,302	307,036			
45,738	2.78	144,658	9.5	••	3,019	10,603	82,989	128,378	835,357			
629	0.003	48,159	3.2	••	455		10,956	272,983	981,329			
65,868	8.88	269,069	17.6	1	1,553	110,849	68,700	256,976	1,706,561			
14,884	0.88	124,324	8.2	195	1,078	22,084	41,244	250,322	2,235,630			
74,282	4.42	73,686	4.8	619	1,116	36,737	44,561	143,107	1,785,50			
6,996	0.49	22,088	1.4	1,874	10,080	34	23,318	119,338	1,715,84			
41,778	2.48	45,487	2.9	8,115	8,684	.89,626	59,236	197,430	1,340,894			
102,786	6·12	99,165	6-5	543	12,322	31,888	34,886	262,626	1,277,796			
8,936	0.28	163,331	10.7	2,602	1,366	541	38,160	176,987	2,072,19			
153,717	9·15	263,865	17.8	21	28	37,094	85,647	288,290	1,399,96			
\$26, 550	19-43	213	••	••	24	50,110	2,049	44,451	424,090			
254,944	15·18	1		••		41,647	769	47,579	844,940			
76,491	4.55				1	35,868	30	43,793	156,20			
285,844 173,281	12·83 10·81			••	473	152,068 3,125	116 388	168,876 6,417	556,92 184,24			
678,207	100.00	1,522,616	100.00	25,228	155,857	518,732	486,412	<u> </u>	18,620,75			

			 					
		Fodder		Sugarci	ane.		Fruits	and vege
	Names of the Districts.	crops.	Sugarcane.		Other.	Total.	Fruits.	P. C.
		Атев.	Area.	P. C.	Агеа,	Area.	Area.	
1	2	19	20	21	22	23	24	25
1	Ahmedabad .	11,701	636 753	1·2 0·82	5	641	822	2.26
2	Kaira	45,417	123 87	0·2 0·09		123	213	0.58
3	Panch Mahals .	2,551	80 66	0·1 0·07		80	33	0.09
4	Broach	3 823	17 50	0·0 0·05	70	87	366	1.00
5	Surat	1,399	2,723 3,833	5·00 4·15	833	3,056	944	2.60
6	West Khandesh	198	753 1,317	1·3 1·43		753	433	1.19
7	East Khandesh	42	354 1,218	0·6 1·32	••	854	2,762	7.61
8	Nasik	5,721	5,431 10,041	10·1 10·88	1	5,432	2,367	6.52
9	Ahmednagar	3,280	4,079 10,324	7·55 11·19	••	4,079	2,718	7-49
10	Poons	1,805	13,196 17,207	24·3 18·64	76	18,272	4,825	13.29
11	Sholapur	1,931	1,566 3,065	2·9 3·32	181	1,747	1,499	4.18
, 12	Satara	1,861	8,533 16,742	15·7 18·14		8,533	2,027	5.28
13	Belgaum	27,806	10,364 16,060	19·00 17·40		10,364	1,796	4.95
14	Bijapur	1,181	635 1,701	1·2 1·84	20	655	1,516	4.18
15	Dharwar	15	1,924 5,284	3·6 5·73	115	2,039	2,795	7.70
16	Thana	110	399 510	0·7 0·55	••	399	3,632	10.01
17	Kolaba	1	32 20	0.00	61	93	422	1.16
18	Ratnagiri		539 929	0.90 1.70 1.18	••	539 929	703 1,393	1·93 3·77
19	Kanara	1	1,093 2,885 2,934	1·18 5·3 3·18	••	2,885	6,412	17.67
	Total .	108,793	54,269 54,659 92,305	100.00	862	55,131 55,521	36,295 36,975	100.00

des. Includ crops.	ing root	Oil-seeds.	(Details	C	ondiments s	uch as chille	s, etc.
getable.	Total.	To	tal.				Total.
Arca.	Area.	Area,	P. C.	Ares.	P. C.	Area.	Area.
26	27	28	, 29	80	81	32	\$3
40,689	41,511	44,577	3.33	•		2,493	2,453
27,579	27,792	27,930	2.08		**	4,248	4,246
6,777	6,810	49,583	8.70	••	••	1,428	1,428
6,030	6,896	13,038	0.88		.••	276	276
47,104	48,048	25,684	1 92		···	1,545	1,545
17,547	17,980	69,808	5 22		•••	4,256	4,256
21,965	24,727	24,668	1.84		,••	9,627	9,627
28,255	80,622	167,049	12.49			11,391	11,391
80,269	82,987	179,018	13.88		, 	5,716	5,716
84,091	68,916	114,132	8.53			11,907	11,907
89,825	40,824	165,672	12-89		.••	7,970	7,970
71,752	78,779	138,686	10-87		.••	19,607	19,607
5,283	7,079	63,921	4.78			17,131	17,131
26,399	27,915	142,696	10-67		••	3,751	8,751
24,518	27,813	68,690	6.18	·		50,496	50,496
5,779	9,411	11,011	0.82	l	••	2,004	2,004
9,238	9,655	4,750	0:35	••	•••	196	196
4,183	4,886 9,683	14,303 83,596	1·07 2·48	::	••	809 1,762	809 1,762
8,290 2,923	9,335	11,957	0.89	1,395	4,160	795	6,350
529,701 533,808	565,986 570,788	1,337,168 1,856,454	100.00	1,395	4,160	155,584	161,139 162,092

	ar a ta ta sa				ibre.		1	Drugs and	
	Names of the Distric	ŧ.	Со	Cotton. San, ambadi and other.			Tobacco,		
·				<u> </u>		<u> </u>	<u> </u>		
			Area.	P. C.	Area.	Area.	Area.	P. C.	
1	2		84	35	86	97	38	39	
1	Ahmedabad	••	284,843	6.8	2,849	287,192	2,964	3.45	
2	Kaira		98,886	0.8	4,587	42,928	26,117	30-42	
3	Panch Mahals	:.	84,417	0.8	9,962	44,879	1,854	1.57	
4	Broach		287,832	5.6	148	287,980	2,303	2.68	
5	Surat		122,976	2.9	3,944	126,920	651	0.76	
8	West Khandesh	٠	478,199	11-4	7,251	485,450	864	1.00	
7	East Khandesh		939,129	22.4	9,825	948,954	3,728	4.33	
8.	Nasik		110,881	2.6	7,690	118,571	2,093	2-44	
ġ	Ahmednagar	••	272,742	8-4	26,338	299,080	4,165	4.85	
10	Poons		25,720	0.6	10,026	35,746	1,127	1.30	
11	Sholapur	••	117,840	2.8	80,431	148,271	2,645	3. 08	
,18	Satara	٠	17,878	0.4	7,622	25,500	8,166	9-50	
13	Belgaum	••	257,997	6.5	11,421	269,418	27,892	32-49	
14	Bijapur	••	693,558	16.6	11,804	705,862	1,406	1.64	
15	Dharwar	••	568,866	13 60	6,568	578,484	368	0.48	
16	Thana	••	2	0.00	1,783	1,735	1	6.00	
17	Kolaba	••		0.00	201	201	·••	0.00	
18	Batnagiri .		8	0.00	2,595 7,014	2,598 7,014		. 0∙00	
19	Kanara	••		0.00	169	169	1	0.00	
	Total	•~	4,198,719	100.00	155,164 159,583	4,313,883 4,358,302	85,840	On .00	

			1	1	í	ł	
Marcotics.							
	Other.		Dyes.	Other non- food grops.	Gross area eropped.	Deduct area cropped more than	Remained net cropp- ed area.
Betelnut.	Other.	Total.					
Area,	Агра,	Area,	Area.	Ares.	Area.	Area.	Aron
40	41	42	48	44	45	46	47
••	1	2,964	292	498	1,128,916	24,993	1,103,923
••]]	26,117	1,806	287	659,220	24,798	684,427
		1,354		5	495,817	75,611	420,206
		2,308		66	557,817	1,101	556,216
	1	652		160	514,500	61,819	453,181
	40	904	51	89	1,414,791	23,879	1,390,912
	217	8,940	8	64	1,998,708	17,681	1,976,077
••	260	2,353	••	297	2,047,997	61,627	1,986,370
••	481	4,848		814	2,814,750	59,696	2,755,054
••	636	1,768	••	216	1,983,262	86,756	1,896,506
••	206	2,851	••	245	2,085,856	85,116	2,050,240
••	591	8,757		1,173	1,618,790	116,326	1,502,464
	852	28,244	••	44	1,701,803	28,419	1,673,384
••	79	1,485	1	18	2,955,211	7,760	2,947,451
	1,759	2,127		8	2,124,088	21,378	2,102,710
4	896	401	••	98	449,254	6,978	442,281
851	18	864		. 9	860,209	11,368	848,841
1,257 8,821	40 182	1,817 4,008	::	6	180,661	11,868	168,793
18,019	408	18,428	1	45	283,414	22,588	210,826
19,651	5,474 9,437	110,965 118,651	1,659	3,587	25,819,064	699,202	24,619,862

In examining the general characteristics of this distribution of crops, the first point to be noticed is the fact that no less than 76 per cent, of the cropped area is occupied by food crops. The relatively large area thus occupied is on the whole a matter for satisfaction. Among these food crops, cereals and pulses take up 73.5 per cent. of the total acreage, while vegetables and fruits only require 2.2 per cent. and sugarcane only 0.2 per cent.

The non-food crops consist of oil-seeds, fibres, condiments like chillies, narcotics like tobacco, and fodder crops. Oil-seeds occupy 5·3 per cent. of the cropped area, while fibres, of which almost the whole is cotton, take no less than 17·2 per cent. When set out as a percentage of the total cropped area the amount of space devoted to chillies and similar crops seems very small, for it is only 0·64 per cent. of the cropped area, but this really represents not less than 160,000 acres, while the heading "drugs and narcotics", eighty per cent. of which is tobacco, and practically all the rest betel nut, accounts for over 110,000 acres or 0·44 per cent. of the area under crops. The area under fodder crops is perhaps the most disappointing of all the headings we have to consider, for they, so vital in a dry country for keeping livestock in good condition, only occupy 109,000 acres or 0·43 per cent. of the area.

Of the cereals jowar, bajri, rice and wheat are the most important judged by the area devoted to them, and in this order. They occupy 26, 20, 7 and 6 per cent. respectively of the total area under crop. A better idea of their relative importance is perhaps given by taking the percentage of the total area under food crops. The figures then come to 34 per cent. for jowar, 27 per cent. for bajri, 9 per cent. for rice and 8 per cent. for wheat. Thus these four crops take no less than 78 per cent of the area devoted to the growing of food. Two other food grains are important in this connection. The first of these is one of the smaller millets, known as nagli, nachani or ragi in various parts of the Bombay Presidency, and this occupies 3 per cent. of the total food area, chiefly in the uplands of the Konkan and the mallad tract. The second of the remaining food grains is the pulse, gram, which takes 2 5 per cent. of the food area.

Of the oil-seeds safflower, sesamum, groundnut, nigerseed and linseed are important in this order. This is, I suppose, the only large area in the world where safflower stands first among oil-seed crops. Among the fibres I have already said that almost the whole area is under cotton. The only other one which has the least importance is sann-hemp, or, as it is called on the market, Bombay hemp.

The following table gives the population, the cultivated area and the cultivated area per head of population in each of the districts of the Bombay Presidency:—

Distri	c t.		Population.	Net cropped area.	Acreage of cultivated area per capita.
Ahmedabad			827,809	Acres.	1.00
Kaira Kaira	••	• • •	691,744	1,103,923	1.33
Panch Mahals	••	••		634,427	0.92
ranca manais Broach	• •	•••	322,695 306,717	420,206	1.30
Burat	••	• • •	654,109	556,216	1.81
West Khandesh	••	• • •	580,723	453,181.	0.69
	••	; •		1,390,912	2.39
East Khandesh	••	• • •	1,034,886	1,976,077	1.91
Nasik	••	•••	905,030	1,986,370	2.19
Ahmednagar	••		945,305	2,755,054	2.91
Poons	••		1,071,512	1,896,506	1.77
Sholapur	••	••[768,330	2,050,240	2.67
Satara	••		1,081,278	1,502,464	1.39
Belgaum	• •	• • •	943,820	1,673,384	1.77
Bijapur	••		862,973	2,947,451	3.42
Dharwar			1,026,005	2,102,710	2.05
Thana			882,309	442,281	0.20
Kolaba	••	[594,166	348,841	0.59
Ratnagiri	••		1,203,638	168,793	0.14
-		- 1		*571,826	0.48
Kanara	••		430,548	210,826	0.48
Sombay City	• •		979,445		<u> </u>
	Total		16,113,042	24,619,862	1.53

^{*} Including estimates for non-reporting area.

This table will be found exceedingly interesting especially as regards the figure for the amount of cultivated land per head of population. It will be seen that taking the Bombay Presidency as a whole the cultivated area per head of population is 1.5 acres. Ratnagiri has the least, or slightly under half an acre per head. Small areas per head are, however, usual in rice growing areas, for rice is, on the whole, an intensively grown crop. Bijapur, on the other hand, has the greatest, or nearly $3\frac{1}{2}$ acres, this being a district of extensive but poor cultivation, with very little irrigation.

It is interesting to compare these figures with those prevailing in some of the other countries of the world. Thus we have as follows:—

Area of Cultivation per head


United States of America . 3.5
France . 1.5
United Kingdom . 0.4
Japan . 0.3
India as a whole . 1.0


Taken altogether, the figures explain why the Konkan districts are not self-supporting, especially Ratnagiri, in spite of the fact that they grow rice. They have always to import food from the outside, and a large part of their population has to go outside, chiefly to Bombay, for employment. The Khandesh districts are apparently well supplied with cultivated area in relation to the population, until it is recognised that nearly half the area is occupied by non-food crops, and that hence the food crop area per head is really small, and that therefore, except in good years, these districts are not self-supporting in the matter of food.

If we take the Bombay Presidency as a whole, without taking any account of the special conditions of each district, we find that taking cereals and pulses alone, which form almost the whole of the food of the population, it will be seen that the province produces enough food to supply about 713 pounds a year per head, or about two pounds of grain per day for each member of the community. This estimate is based on what are supposed to be normal yields, which are only attained in good years. The areas and amounts of produce corresponding with them which are used for this calculation are shown in the following table:—

No.	Name of crop.		Acreage.	Outturn in lbs.	Remarks.
1	Jowar	Ī	6,503,540	4,357,371,800	Figures of average
2	Bajri		5,119,119	2,047,647,600	yields of grain per
3	Wheat		1,522,616	875,504,200	
2 3 4	Rice	(0		2,064,194,610	
		(2		196,004,190	
5	Maize		155,857	83,539,352	
6	Nagali	= 0		549,855,920	ending 1917.
•	L. mB-ma	(2		123,172,000	Figures are estimates
7	Gram		486,412	199,428,920	for the unreporting
8	Tur		513,573	265,003,668	
9	Other cereals	a		276,825,244	District are marked
. •	Other coronin	``\}		25,088,032	(2).
10	Other pulses	lõ		451.398.960	۳,
	Other panses	· Č		12,036,418	
	<u> </u>				
		(1) 18,585,535	11,130,769,274	
		(2		356,500,640	

No.	Population.	Quantity of grain and pulses, lbs.	Quantity available per year per capita in lbs.	Quantity available per day per capita in lbs.
1	16,113,042	(1) 11,130,769,274 (2) 356,500,640	712 9	1.95
		11,487,269,914		





V.—JOWAR (Andropogon Sorghum).

Jowar is the most important cereal, and at the same time the largest and most widely grown crop in the Bombay Presidency. As a food crop it is better than rice as the grain contains more proteids, while it also has more starch than wheat. The grain is chiefly eaten as a bread stuff in unleavened cakes. Particularly in the eastern districts it almost entirely replaces rice and wheat, and is a quite indispensable food.

Its only rival in importance as a food crop is bajri, and these two claim about forty per cent. of the total cropped area. Most of the produce is, however, consumed locally, the export not amounting to more than one per cent. of the total produce. In bad years of drought the Bombay Presidency imports jowar from Sind and Madras, while in good years there is an export to Kathiawar and Cutch.

Jowar is equally important as a fodder crop, as jowar straw is the main dependence of the cattle in the larger part of the Bombay Presidency and a failure of the kharif jowar crop means inevitably an appalling death of cattle, and the other results of famine.

Statement No. I shows that Bijapur claims the largest area of any district and contains more than twenty per cent. of the gross-cropped area in the Bombay Presidency. Sholapur stands second with 17 per cent., while Ahmednagar, Poona, Satara and Dharwar each have between eight and a little over ten per cent. The greater part of the remaining area lies in Belgaum, East Khandesh, Ahmedabad and Broach.

Statement No. II shows that jowar claims fifty-four per cent, of the total cropped area of the Sholapur district, and over forty-four percent. of Bijapur, while in the Satara, Poona and Belgaum districts more than thirty per cent. of the cropped area, and in the Dharwar, Ahmednagar, Broach and Ahmedabad districts more than twenty percent. of the cropped area are under this crop.

Agriculturally speaking, three main divisions of the *jowar* crop may be made: (1) the early rain or *kharif* varieties, which are either red or white; (2) the late rain or *rabi* varieties, which are all white; and (3) the irrigated hot-season crop grown for fodder alone, whose area is very small.

Of the total area of 6,503,540 acres of jowar 4,191,192 acres or nearly two-thirds is grown in the rabi season, and the remainder or about one-third in the kharif season. Sholapur, Bijapur and Ahmednagar together claim about three-quarter of the rabi area; Poona, Satara and Dharwar also return a large area of rabi jowar. Dharwar, Bijapur and East Khandesh claim half of the total kharif area, while Belgaum, Ahmedabad, Satara and Poona grow very large quantities of kharif jowar.

Jowar is generally a dry or jirayat crop when grown for grain; wher grown for purely fodder purposes either in the rabi or in the hot season it receives irrigation. It rabi jowar be irrigated the outturn is naturally very much greater.

There are over a hundred varieties of jowar in the Bombay Presidency. Among the best known varieties are Sathi, Perio, Chapti, Sholapuri, Ais-puri, Garya, Gudhaghi, Bedri, Dagdi, Maldandi, Dukri, Gidgap, Argad, Utavli, Nilwa and Sundhia.

The distribution of the crop is regulated by the amount and distribution of rainfall, but the depth and character of the soil plays an equally important part. In the Konkan and Mallad tracts, where the rainfall is very high, jowar has no place at all. In the parts of the Deccan where the soils are shallow it gives place to bajri, while in the Panch Mahals and in part of the Belgaum district maize takes the place of jowar. It is essentially a crop of deep and heavier soils, while the best results are obtained in centres with an average rainfall of from thirty to forty-five inches. Kharif jowar is sown in the month of July, either alone or in mixture with other crops. Rabi jowar is sown in September and October, and in many centres it is mixed with other crops, notably with safflower. On an average the crop occupies the land for five months.


The average yield of jowar per acre* comes to about 670 lbs. for the dry jirayat crop. The average yield of fodder per acre, in addition to the grain, comes to about 1,700 lbs. and the normal proportion of grain to fodder is about 1 to 2.5.


VI.—BAJRI (Pennisetum typhoideum).

Bajri stands second in importance as a food crop, occupying about 5,100,000 acres or about 20 per cent. of the gross cropped area in the Bombay Presidency. It is the staple crop in a large tract, and is the chief tood of large classes of the people, but it is grown only where it gives better results than jowar. It is a light soil millet, while jowar is chiefly found in heavier soils, though on much of the land the crops are alternative, according as the weather conditions are more favourable for one or the other.

It is entirely a rains crop, and the area fluctuates considerably from year to year. When the monsoon opens with liberal rainfall, jowar occupies a larger area, while when the early rains are deficient, the land put under bajri tends to increase. In the Deccan the tract where bajri growing is most intense lies in the western part of the Desh; the crop does well on somewhat less rainfall than is required by kharif jowar.

Quinquennial report on the average yield per sore of principal crops for the period ending in 1916-17.





It is grown almost entirely as a food crop for the people themselves, and hence there is no export whatever of this crop to foreign countries. Unleavened bread made from bajri is preferred to jowar bread in many parts of the country in the cold season.

Statement No. I shows that Ahmednagar claims 20.8 per cent. of the total area under bajri, Nasik 16.6 per cent. and Poona 14.25 per cent. These districts are followed by Sholapur, Satara, West Khandesh, East Khandesh and Bijapur, each of which grows between 7 to 8 per cent. of the total area. Ahmedabad claims 3.9 and Kaira 3.0 per cent.

There are two varieties of bajri grown in Gujarat, chiefly (as the above figures show) on the light goradu soil of Kaira and Ahmedabad. These varieties—deshi and bhavnagri—have special characters, and are on the whole superior in yield to the variety grown in the Deccan. In the Deccan there is only one type, though bajri grown in the Akola taluka of the Ahmedaagar district is reputed to be of better quality and to give a higher yield.

Statement No. II shows that bajri claims about 42 per cent. of the total cropped area in Nasik, about 37 per cent. in Poona and also in Ahmednagar. Next in importance come Satara, West Khandesh and Kaira, where it covers from 20 to 26 per cent. of the total area.

Bajri as a nutritious food stands very high, containing about ten per cent. of proteids and seventy per cent. of starch. It, hence, compares very favourably with jowar as a food.

It is always a mixed crop in Gujarat. In certain parts of the Deccan, tur (Cajanus indicus) is grown in rows with bajri. In Khandesh, however, it is grown alone. If heavy rain falls when the crop is inflower it suffers much, and hence the crop is generally sown late to avoid heavy showers during this stage of its growth. The normal average outturn in the Bombay Presidency comes to 400 lbs.* of grain and 1,000 lbs. of fodder per acre. Bajri straw, however, makes poor fodder, much poorer than jowar straw. The proportion of grain to straw generally is about 1 to 2.5.

VII.—RICE (Oryza sativa).

Rice stands third among the food crops of the Bombay Presidency, and as its cultivation very largely occurs in the areas where jowar and bajri are not and cannot be grown, it is for these tracts of great importance. As a food crop rice is not equal either to jowar or bajri as the grain is starchy and somewhat deficient in proteids. But, nevertheless, it is indispensable in the districts where it is grown. As a fodder crop it is far inferior to jowar, both in the quantity and quality of the straw which it yields, and as a result the cattle in districts devoted to rice growing are usually very inferior.

Quinquennial report on the average yield per sore of principal crops for the period ending 1916-17.


The rice produce of the Bombay Presidency is not sufficient to meet the needs of the people. In most districts the peasants and labouring classes go without rice and depend on millets only. Even in the Konkan districts where it is a staple crop and the main food, the local produce often falls short of their requirements, and then rice is imported from Madras, the United Provinces, the Central Provinces and Burma. In good years, however, large quantities of rice are exported from Bombay to Europe, though a large part of it comes from other provinces and only passes through Bombay.

Statement No. I shows that the area under rice is about 1,678,000 acres, and this represents 9 per cent. of the land under food crops or 6 6 per cent. of the gross cropped area in the Bombay Presidency. The districts which chiefly contribute to this total are Thana with 20 per cent., Kolaba with 15 per cent. and Ratnagiri with 13 per cent. of the whole area under rice. Kanara (10 per cent)., Dharwar (9 per cent.), and Belgaum (6 per cent.) follow, in this order of importance, as rice growing centres. A fairly large area is also found in the districts of Gujarat, where the processes of cultivation are peculiar, and where it is found fairly well scattered over the districts. Rice is not grown at all in the eastern Deccan.

Statement No. II shows that in three districts—Kanara, Thana, and Kolaba—rice occupies more than 70 per cent. of the whole cultivated area. In Ratnagiri, where there is more varkas land (which is not suitable for rice cultivation) than anywhere else, the proportion falls to a little under 40 per cent. No other district shows more than 18 per cent. found in Surat, or between 13 and 14 per cent. found in Kaira and in the Panch Mahals.

The distribution of the rice area is governed either by the rai nfall or by the existence of facilities for artificial irrigation. Most varieties of rice are semi-aquatic and need water at the most for from four to five months. The eastern limit of the ordinary rice crop in the Deccan and Karnatak districts follows more or less clearly the line of mean monsoon rainfall of about 50 inches. In the Nasik district, where there are a number of dams along the rivers, rice extends beyond this line along the rivers. In the Gujarat districts rice is grown on a fairly large scale, and in this tract there are a number of small tanks which partly protect the crop. The growing period of most of the varieties of rice corresponds to that of the south-west monsoon, namely from June to October.

In the Konkan districts, in Gujarat and in the heavy rainfall strip of the Deccan called *Maval* or *Mallad*, along and close to the Western Ghats, rice is generally transplanted. On the other hand, in the salt rice ands of the coast and in the uplands in parts of Ratnagiri and Kanara, broadcast sowing is practised. This is also the case



NAMES OF DISTRICTS

A C R E S IN T HOUS AND S

17 NAMA
2 K BLA BA
3 MA NA RA
4 DUAR WAR
3 SEL E AUM
6 SURAT
7 K A IRA
8 PROFINA
18 P

in Gos and in the Karwar and Ankola talukas of the Kanara district. In the eastern part of the *Mallad* tract, in Belgaum and Satara, wehre the rainfall is only from forty-five to sixty inches, rice is sown with a drill Likewise in parts of Broach and the Panch Mahals rice is often drilled between rows of cotton. Drilled rice is also not infrequently grown on the Deccan canals on sugarcane land. Hot weather rice, known as vaingan, in the Konkan and Kanara, and *kharif* rice under the irrigation from rivers in the Nasik and Khandesh districts are sown by broadcasting sprouted seed.

In the Konkan districts, including Kanara, as also in the Mallad tracts, early varieties of rice ripening in one hundred to one hundred and twenty days are grown on high-lying areas which do not retain sufficient water, while late varieties maturing in 120 to 160 days are grown in low-lying fields.

In the Konkan districts, except Kanara, and in the Mallad tracts, seedlings for transplanted rice are grown in a specially treated seed-bed. The treatment of the land is called "rab." It consists in heating the surface soil of the seed-bed by burning materials like dried cattle dung, leafy loppings of trees, grass, etc., on the surface. This system of preparing the seed-bed of rice is almost peculiar to the Bombay Presidency and has been proved to be very effective. In Gujarat and Kanara the rice seedlings are grown in a seed-bed manured with ordinary farm manure,

Rice freely responds to manuring. But in the richer trap soils of the North Konkan and the *Mallad* tract where transplanting is practised, the crop hardly receives any manure. On the other hand, the crop on the poorer laterite soils of the South Konkan, on areas where drill sowing is resorted to, on the land under river irrigation in the Nasik district, and on that under tanks in Gujarat, is regularly manured.

For successful rice cultivation a good and constant supply of water is essential. This can be best secured by having level plots with proper embankments. In the Konkan and on the Ghats, where the general configuration of the land is uneven, this levelling is brought about by a very skilful and elaborate process of terracing according to the contour of the land. This prevents scouring, and enables the small plots to hold evenly fair depths of water all over the field.

In the Konkan, on the Western Ghats, and also in Gujarat, rice is sown year after year in the same land. Occasionally, well drained rice lands go under sugarcane. In the Nasik district the lands under river irrigation however, receive, in regular rotation, rice, sugarcane and wheat or bajri.

There are hundreds of varieties of rice. Some are early, ripening in from 90 to 120 days, and some are late, ripening in from 120 to

180 days. Early varieties give generally a coarser grain, and occupy uplands which are not capable of retaining very much water. Late varieties give generally finer grain, and usually occupy the low-lying fields. Of the early varieties Varangal, Dodka, Panwel and Waksal are a few of the well known types. Among late varieties Kamod, Ambemohor, Raybhog, Kolamba and Garri Patni may be mentioned. Ambemohor, Kamod, Jiresal, Pankhali are a few of the scented varieties.

The average outturn of rice per acre comes to about 1,200 lbs. of grain and 1,600 lbs. of straw. Rice straw is used as fodder, though it is not considered to be very sustaining, and also for thatching. The proportion of grain to straw generally comes to about 1 to 1.25.

VIII.—WHEAT (Triticum sativum).


Wheat occupies the fourth place among cereals, and the fifth among all crops in point of acreage in the Bombay Presidency, having 1,522,616 acres or about 6 per cent. of the gross cropped area to its credit.

Wheat is one of the crops in the Bombay Presidency which are almost entirely grown rot for consumption but for sale. Compared with the population, the total produce is small, but a large part of the people eat bajri and jowar almost exclusively and hence in good years wheat is exported to England. In bad years wheat is imported into the Bombay Presidency from other provinces such as Rajputana, Central India, the Punjab, the Central Provinces, the United Provinces, etc. When exported, Indian wheats fetch poor prices relatively to those for American or Argentine produce. This is only partly due to any intrinsic fault in the wheat, but is mainly caused by the way in which it is marketed and exported. British merchants complain of the dirt and mixture of inferior grain in Indian wheats.

Statement No. I shows that Nasik stands at the head of the districts in the Bombay Presidency as wheat producers containing 17.6 per cent. of the total wheat area, followed closely by Dharwar which has 17.3 per cent. After this come Ahmedabad (10.7 per cent.), West Khandesh (9.5 per cent.), Ahmednagar (8.2 per cent.) and Belgaum (6.5 per cent.) in that order.

Statement No. II shows that wheat occupies 13.3 per cent. of the cropped area of the Nasik district, about 13.0 per cent. in Ahmedabad, 12.5 per cent. in Broach, 12.3 per cent. in Dharwar and 10.2 in West Khandesh.

Most of the crop is a *jirayat* or dry crop. Only 5 per cent. is irrigated. Nasik, Ahmednagar, Satara, Ahmedabad and West Khandesh claim fairly large areas under irrigation.



NAMESOF DISTRICTS

A C R E S IN T H O U S A N D S

INASIN

E DHARWAR

3 BIJAPUR

3 BIJAPUR

4 ANNEDMEAR

7 9 FLS AUN

8 PRONA

9 RORCH

M 6 STARA

12 SWAT

13 NOLAPUR

15 NOL

Kata or Shetgahu, Wagia, Daudkhani, Bakshi, Mundi, Potya and Khapli are the principal varieties grown. Khapli and Bakshi (Kala. Kusal) are irrigated wheats; the others are generally taken as dry crops though in some tracts they receive a little irrigation.

The distribution of the wheat crop as well as its outturn per acre is governed by the moisture in the soil during the growing period of the crop, as well as on the length of the period available to the crop for making growth. When the weather gets hot the crop ripens quickly. The crop is, of course, always and entirely a rabi or cold weather crop.

The normal average outturn of wheat for the Bombay Presidency comes to about 575 lbs. of grain and about 1,000 lbs. of fodder per acre. The proportion of grain to straw generally is about 1 to 1.6.

IX.—COTTON (Gossypium).

Cotton has been cultivated in India from very early times, and cotton cloth has been, from before the dawn of history, the clothing of people in this country. There was a time when Indian cloth was highly valued in many parts of Europe. In recent years there has been a very great increase in the cotton area, due to the demand for the crop for export to European and in very recent times to Japanese markets. Bombay stands first among the provinces of India as a grower of cotton, and, in fact, out of the total number of about 750 presses and 250 mills in India, the Bombay Presidency claims about 220 presses and 180 mills.

Cotton is one of the most important crops of the Bombay Presidency. It occupies the third place in acreage, and claims about 17 per cent. of the gross cropped area. The proportion of the area under cotton in Bombay to the total area under that crop throughout India is about 17.7 per cent. according to the Indian Cotton Committee's report.

Statement No. I shows that East Khandesh possesses the largest share of the crop among all the districts, namely 22 per cent. of the total area under cotton in the Presidency. Bijapur, Dharwar and West Khandesh follow with about 17, 14 and 11 per cent., respectively. Ahmedabad and Ahmedagar, Belgaum and Broach contain from 6 to 7 per cent. each, while Surat, Sholapur and Nasik have about 3 per cent. each.

If we look at the proportion of the area under cotton to the total of the cropped area of each district we again find East Khandesh standing highest, where the cotton area form no less than 47 per cent. of the gross cropped area of the district. Broach has 43 per cent. of its area under cotton and West Khandesh, Dharwar, Ahmedabad, Surat and

Bijapur return between 25 and 33 per cent. of their gross cropped area as under this crop (see Statement Π).


The Bombay Presidency exported before the war large quantities of raw cotton to Japan, Italy, Germany, Belgium, France and the United Kingdom, but the European trade has now largely fallen off while that with Japan has greatly increased. Cotton yarn has been in the past exported to China (Hong-Kong), Turkey in Asia and the Strait Settlements, while piece-goods have been exported to many centres in Asia and Africa. On the other hand, large quantities of cotton manufactures,—cotton twist and yarn, and piece-goods,—have been imported from England and other European countries.


Roughly speaking there are four commercial types of cottons in Bombay—Dholeras, Broach, Oomras (or Khandesh) and Kumpta-Dharwar. Dholeras include all the Cottons (Wagad, Lalio, Goghari and Mathio) grown in the north of Gujarat. Broach includes Navsari, Surat, Broach and Kanvi grown in the south of Gujarat. "Oomras" include only one type which is common in the West Khandesh, East Khandesh, Nasik, Ahmednagar, Poona, Sholapur and part of the Bijapur districts: Kumpta-Dharwar or Karnatak cottons include the types known in the trade as Kumpta and Dharwar American.

Of the total area under cotton, about 60 per cent. grows short stapled (1/2" to 5/8") and 40 per cent. long stapled (5/8" to 1") varieties. The following table gives the commercial types, varieties, length of fibre (staple) and the ginning percentages of the cottons grown in the Presidency:—

				Staple.	Ginning Percentages.
•				inohea	per cent.
Dholeras		Wagad	••	 6-8 to 7-8	33
		7 -1.	••	 5-8 to 6-8	33
		Contoni	••	4-8 to 5-8	40
		26-43:-	••	4-8 to 5-8	32 .
Broach		Navsari		7-8 to 1	31
		Quant	•	7–8	32
	- 1	D1	••	5-8 to 6-8	32
		Fanni	••	 4-8 to 6-8	35
Kumpta-Dharwar		Kumpta		7–8	26
	1	Dharwar-Ameri	can	 6-8 to 7-8	30,
Oomras		Khandesh	••	 4-8 to 5-8	38

The average normal yield of cotton per acre for British India comes to 102 lbs. of lint. It may be remembered that in Sind the outturn per acre is 170 lbs. In Egypt it is 140 lbs., but both in Sind and





in Egypt the crop is irrigated. In the United States of America, the yield of lint per acre is said to be 200 lbs. The average yield of Gujarat and Karnatak cottons comes to 128 lbs. and that of Khandesh to about 120 to 150 lbs. per acre.

In Bombay, the cotton is a jirayat or dry crop, though in Ahmedabad and Kaira it occasionally gets irrigation. Except in the Karnatak it is a Kharif crop, being sown as soon as the land is well soaked with rain. At times it is sown in the dry soil in anticipation of early rain. In Khandesh, picking commences in October and is over before December. In Gujarat, cotton is picked in January and February. Kumpta Cotton is sown in about the second week of August, and Dharwar American two to three weeks later. Picking of cotton in this area commences by the second fortnight of February, and lasts till the end of April.

Cotton does well in deep black soil having 20 to 50 inches of rainfall. In the Broach and Surat cotton tracts, where the soil is deep and the rainfall between 30 to 40 inches, long staple cotton (Broach variety) does well. As the Karnatak receives the benefits of both the monsoons, it also grows a long stapled variety (Kumpta). In the Deccan districts, on the other hand, which receive between 20 to 30 inches of rain per annum, short staple cotton is usually grown, being generally one of the types of "Khandesh".

X.-SUGARCANE.

India is supposed to be the original home of Sugarcane and she still claims the largest acreage under this crop of any country in the world. Till the seventeenth century India supplied sugar to Europe. Now she imports as much as, 7,000,000 tons every year, most of it coming, before the war, from Mauritius, Java, Austria-Hungary, the Straits Settlements, Hong-Kong and China. Among the different provinces of India, Bombay occupies only the sixth place as regards area under this crop, for the United Provinces stand first, and the Punjab second, while Bihar and Orissa, Bengal and Madras, occupy respectively the third, fourth and fifth places. But while the area in Bombay is small as compared with other provinces, the yield per acre is larger than anywhere else, and in outturn of gul or crude sugar, Bombay stands third, the amount of product only being exceeded in the United Provinces and in the Punjab.

In the Bombay Presidency sugarcane only occupied 92,000 acres in 1917-18, and this itself represents a considerable increase in the area three years before. It should, however, be remembered that owing to the intensive nature of the cultivation, it produces gul worth between five and eight crores of rupees a year. It should further be remembered

that most of the *gul* produced in Bombay is consumed locally, though a very large quantity of sugar is imported annually from outside India. Hence the importance of this crop is very great.

Considerable quantities of gul are supplied by the Karnatak and the Deccan districts to the Central Provinces, to Berar, to the Nizam's Territories, and to Rajputana, while the Karnatak also sends gul to Mysore. The Konkan and Gujarat draw their supplies of gul from the Karnatak and the Deccan.

The average normal yield of crude sugar (gul) per acre for British India comes to 2,719 lbs. while for the Bombay Presidency it is no less than 6,950 lbs. The average yield for the thin varieties of cane is about 3,500 to 4,500 pounds per acre, for thick varieties under well irrigation, it is about 5,000 to 7,000 pounds per acre, and for thick canes under canal irrigation it reaches 7,000 to 11,000 pounds per acre. The amount of gul for the ordinary thick cane grown (pundya) is about 12.25 per cent. of the weight of cane. The outturn of gul in Bombay compares favourably with that of any other country in the world. The cost of cultivation, however, is rather high as compared to that of other countries. The green tops of the cane are fed to cattle. The proportion of gul to tops comes to about 1 to 1.3.

So far there are no sugar factories in Bombay actually working. Recently two large companies have been floated, and with the extension of irrigation canals, Bombay would appear to present a good field for the development of sugar manufacture.

Statement No. I shows that Poona, Satara and Belgaum are the principal districts producing sugarcane, each claiming about 18 per cent. of the total area under cane in the Bombay Presidency. Ahmednagar and Nasik follow, each growing about 11 per cent., each of the total area. Dharwar claims about 6 per cent., Surat about 4 per cent., while Sholapur and Kanara have about 3 per cent. each.

Statement No. II shows that Kanara has the largest proportion of the cropped area (more than 1.2 per cent.) of the district under sugarcane. Though Poona claims the largest acreage under this crop, the proportion of the crop to the total area comes to only 0.6 per cent.


There are many varieties of cane grown of which the principal are as follows:—


Thick and juicy canes —

Pundya (pale yellow) grown on the Deccan canals and in the Deccan districts.

Kabirya (striped) in Sholapur, in the Walva taluka of the Satara district, and in parts of Kanara.

Malbari (pale yellow) in the Surat District.





Medium thick canes less juicy than the above group-

Kavangiri or Kala Malbarl (medium thick red cane) in the Konkan and in the Nasik District.

Bangdya: in parts of Nasik, though it is now disappearing.

Thin and hard canes—

Khadya, in the western parts of Nasik and in West Khandesh. Wansi, in parts of the Mallad tract.

The area occupied by the medium and thin varieties is only relatively small.

Sugarcane is always an irrigated crop in Bombay. It requires heavy manuring and regular watering to get the best results, and it does better on level well-drained soils. Soils two to four feet deep with good drainage are specially suitable, and in such soils it requires watering once in ten days. On shallow soils it requires watering every sixth to eighth day. In the Karnatak districts all the cane is planted in December, while on the Deccan canals most of the planting is done in February and March. In the Surat district, cane is chiefly planted in November and December. On the Deccan canals the cost of cultivation of this crop comes to between 500 and 700 rupees per acre, though many people spend still more. On newly opened canals two ratoon crops are taken for about fifteen years: in the older sugarcane tracts the second ration does not pay, and only one ration is taken. In the oldest canal tracts rations have been almost entirely abandoned, and only plant cane is taken. Even on the new canals after the ration is taken, the land goes for a year or two under some dry crop.

The system of weights in different centres varies. In the Karnatak, gul is sold by the atki of about 247 lbs.; at Kopargaon in the Ahmednagar district by the palla which weighs 250 lbs.; while at Baramati in the Poona district gul is sold by the atki of 200 lbs. These are a few of the many different weights used for gul.

XI.-POTATO.

The potato crop only occupies a small area, but it is an intensive crop and is hence perhaps worthy of special attention. It is an introduced crop in this country, and first appears in the Bombay Presidency about 1840. It is said to have been first tried near Junnar in the Poona district, while later on it was introduced by Chinese prisoners at Mahableshwar about 1857.

Potato occupies about 8,200 acres in the Bombay Presidency. The Poona district, growing 6,345 acres, claims 77 per cent. of the total area, while Nasik, though it only grows 458 acres stands second. If we take the taluka as the unit, the Khed taluka of the Poona district

occupies 62 per cent. of the total area in the Bombay Presidency. The Sirur taluka of the Poona district claims 7 per cent., the district of Dharwar about 4 per cent., and the Junnar taluka, the Nasik and the Purandhar talukas about 3 per cent. each.

Khed, Mahableshwar, Belgaum, Dharwar and Ahmedabad (in the last case in the bed of the Sabarmati river) are well-known centres though the area at the latter four places is not very great.


The potato crop is grown both in the *kharif* and *rabi* season in the Khed tract and near Mahableshwar. At Belgaum it is purely a *rabi* crop while at Dharwar it is a purely *kharif* crop. Near Ahmedabad potato is grown in the bed of the Sabarmati in the *rabi* season. *Kharif* potatoes have generally to depend on the rainfall. The crop takes about 3½ months to mature. In the method of growing usually adopted, about 1,000 to 1,100 lbs. of potatoes are required per acre for seed, and the outturn in average years is from about 5,000 to about 8,000 lbs. per acre, though over 16,000 lbs. per acre has been reached. In good years it is a paying crop, but if the crop fails, the loss is very great on account of the high cost of seed.


Large quantities of Italian seed potatoes come to the Bombay Presidency every year, as freshly imported seed has in the past nearly always given better returns than seed potatoes locally grown. It is a crop which has a considerable future in the *Mallad* tract and on the Deccan canals. In the Khed taluka, rabi potatoes are grown in fields which have grown bajri in the previous kharif season, while near Belgaum, the crop is taken in the rice fields after the rice crop is harvested.

XII.—TOBACCO.

Of the plants grown for their narcotic power, tobacco is probably the most important crop all over the world. It occupies 85,840 acres in the Bombay Presidency, of which the Kaira and Belgaum districts together claim about two-thirds. The Satara district stands third with 8,000 acres, followed by Ahmednagar and East Khandesh with about 4,000 acres each. Ahmedabad, Broach, Belgaum and Sholapur have about 2,000 acres each. The Konkan districts do not grow any tobacco. The cultivation of tobacco receives specially careful attention in the Nadiad, Petlad and Anand talukas of Kaira, the Walva and Karad talukas of Satara, and the Chikodi and Hukeri talukas of the Belgaum district.

Tobacco requires very careful treatment of the seed-bed and of the field. The field is well prepared and manured. Seedlings are ready for planting out when about forty days old. Seedlings are planted about 18 inches to 24 inches apart each way, in lines, in the month of August. After about six weeks they are topped keeping





ten to fifteen leaves, the lowest three of which are subsequently removed. After topping of the plants suckering continues, and at the same time the land has to be heed and weeded. Tobacco stands in the field for about five months. It is both an irrigated and a jirayat or dry crop. In Gujarat, tobacco is generally irrigated; in the Deccan it is generally taken as a jirayat crop.

The method of harvesting and curing varies according to the kind of tobacco to be made. Harvesting, drying in the field, pitting or heaping, tying into bundles and stacking require considerable skill and attention.

The outturn of the *jirayat* crop varies between 600 to 1,000 pounds per acre, and of the irrigated crop between 1,000 and 1,600 pounds per acre.

XIII.—OIL SEEDS.

Oil seeds taken together claim 5.3 per cent. of the total cropped area and occupy 1,337,163 acres. It may be noted here that though large quantities of cotton seed are exported for its oil, cotton has not been included in the statistics as an oil seed. The attached table shows the distribution of the important oil seeds by districts.

The figures in the attached table show that safflower, sesamum, groundnut, niger, linseed and castor occupy the major portion of the area which is occupied with oil seeds, and taken together, they form a very important group of crops in the Bombay Presidency. Many of these supply edible oils, a few supply medical oils, while others supply lubricants and other oils required in different kinds of industries. Most oil cakes are useful as cattle food, while all of them can be used as concentrated manures.

The following table gives the percentage of oil content as well as the percentage of oil that is generally expressed from the common oil seeds, as grown in Western India:—

Name of oil seed.					Percentage of oil expressed.		
		44	to	49	38	to	42
••		45	to	50	40	to	45
• •	Í	46	to	50	40	to	42
		. 37	to	44	32	to	38
	- 1	30	to	32	20	to	25
	1	35	to	40	28	to	.32
••		18	to	21	12	to	14
	••						

The following table gives the uses to which the oils and the bye-products of the important oil seeds are put:—

Name of oil	Use of		Remarks.
seed.	Oil.	Cake.	Ivolitatas.
Safflower	Used in cookery and is said to be useful for oil paints.	and manure.	Flowers of some varieties of safflo- were supply dyeing material.
Sesamum	Used in cookery	1 30 m	
	5 N. 1 . 1 . 1		
Groundnut	Used in cookery and is found useful in soap making		Kernels in the pods are eaten raw or roasted. The straw makes
			good cattle food.
Niger seed	Used in cookery		Seeds are used in chutney.
Linseed	Used in cookery,	99	
**. *	paints and; var- nishes.		
Castor	For medicinal use, for lubrication and in hard soaps.	Manure.	1
Cocoanut	Used in cookery and	Cattle food.	
E MAN MAN AND AND AND AND AND AND AND AND AND A	in making vegeta- ble margarine and in candle making.	. :5, '	cookery and is eaten. Husk makes fibre.

Most oils are also used for lamp oils though they burn slowly.

Most of the oil expressed in the Presidency is consumed locally, every tract having a liking for the oil of its staple oil seed; only part is exported. Part of the edible cake is used as cattle food, part is used as manure, and part (generally factory cake) is exported. Castor cake is solely used as manure. Oil cakes are extensively used for manuring sugarcane, and the value of different cakes is determined

by the percentage of nitrogen they contain. The following table gives the average percentage of nitrogen and phosphoric acid contained in the cakes. These cakes which supply nitrogen at the cheapest rate are generally used as manure, especially for sugarcane.

Name	of cake.		Percentage of Nitrogen.	Percentage of Phosphoric acid.	Remarks.
Groundnut	cake		8	1.5	
Safflower	27	••	6	2	Is found specially useful for sugarcane.
Sesamum			5	2	
Niger	"	!	4	2	
Castor	**	••	4	2	Extensively used for cane and serves to check white-ants.
Cocoanut	"	•••	3 .	1 1	CHOOK WHITE-SHES.

It may here be remarked that the most economical plan would be to feed oil cakes to cattle and use their droppings as manure, provided the whole of the latter could be preserved, as most of the nitrogen and phosphoric acid contained in the cakes is obtained in the dung and urine voided by the cattle. In practice, however, it is difficult to conserve even half the quantity of dung and more so to conserve any appreciable quantity of urine in India, where cattle are not stall fed.

The following table gives the average analysis of oil cakes generally used as cattle food. The value of cake as cattle feed depends mostly upon its albuminoid and oil contents. Some cakes like coconnut cake and groundnut cake get rancid when stale and it is necessary to feed these fresh:—

Name of cake.	Percentage of Albuminoids.	Percentage of oil.	Remarks.
Groundnut cake	48	10-13	Gets rancid if stored for a long time.
Safflower:	32	7–10	
Sesamum	36	10-12	Is easily digestible.
Niger "	27	10-12	

Oil seeds represent a fairly large proportion of the exports of Bombay. The average quantities of oil seeds, cake and oil exported annually, previous to the war, were as under. (The table is worked out from the figures for 1907-08 to 1912-13):—

Name of com	modity.	Quantity.	Value.	
				Rs.
Oil seeds		63,247 Tons	• •	95,76,838
Cake		10,552 Tons		8,02,688
Oil	. ••	94,961 Gallons	••	2,47,832
		Tota	ıl	1,06,27,358

It may be noted that cotton seed, poppy seed, etc., are included in the calculation.

Safflower.—The total area of this crop in the Presidency is 426,617 acres, occupying the largest acreage of all the oil seeds. Most of the area is found in the rabi portion of the Desh tract, Ahmednagar, Sholapur, Bijapur and Poona districts, claiming together about three-fourth of the total area. Next come Dharwar and Belgaum. The Konkan and Gujarat do not grow any safflower except a little in Ahmedabad.


It is usually grown as a subordinate crop with rabi jowar, wheat, and gram, though sometimes it is sown alone on the head lands. Often four to five rows of safflower are sown round the main rabi crops named above to prevent cattle getting to it, the spiny nature of safflower serving as a temporary hedge. The area reported is the area supposed to be occupied by safflower alone.


The seed contains 32 per cent. oil and yields 25 per cent. in the country ghani which deals with large quantities all over the Bombay Presidency. Large quantities of safflower seed are also exported. The cake is used as cattle food and is also found to be a very useful concentrated manure for sugarcane on the Deccan Canals. Well prepared safflower cake contains about 38 per cent. albuminoids and at least 6 per cent. nitrogen.

In certain villages in Kaira district the farmers used to gather flowers of this crop which in old times supplied an indigenous dye.

The stalks and other parts of the safflower plant are of no value.

Sesamum (Til).—The total area under this crop in the Bombay Presidency is 255,429 acres, and it occupies the second place among oil seeds. Ahmednagar claims no less than 22 per cent. of the area





under sesamum, followed by West Khandesh and the Panch Mahals with 13 to 14 per cent. each. Nasik has 10 per cent., and East Khandesh, Bijapur, Ahmedabad and Sholapur grow about 5 to 6 per cent. each of the area in the province.

There are three varieties of sesamum commonly grown, namely, white sesamum, red sesamum and black sesamum. The white variety is earlier and also richer in oil, containing about 48 per cent. of oil; the red and black varieties contain about 46 per cent. of oil. The seeds yield about 40 to 45 per cent. of oil on the weight of seed in the country ghani. The cake makes very good food for cattle containing over 30 per cent. of albuminoids. The stalks of til are not eaten by cattle.

Til flourishes on lighter soils. It does not stand heavy rainfall when young. In Gujarat it is sown mixed with bajri. In some districts in the Deccan it is sown alone though in others it is a subordinate crop. The area reported is the area supposed to be occupied by til alone, allowing for mixtures.

Large quantities of sesamum are exported to Belgium, France, Germany and Italy.

When sesamum is grown alone it yields from 300 to 400 lbs. of seed per acre.

Groundnut.—Groundnut is one of the more important oil seed crops in India. It is largely cultivated both in the Madras and Bombay Presidencies. In the Bombay Presidency it occupies 214,912 acres and stands third among the oil seed crops. Of the total area under groundnut Satara grows 42.4 per cent., Sholapur 22 6 per cent. and Belgaum, Nasik and Poona about 8 to 9 per cent. each. The remainder is distributed between East and West Khandesh, parts of the Karnatak, and Gujarat.

Although large quantities of groundnut are crushed locally and fairly large quantities are used as food, the crop forms a considerable portion of the exports in oil seed. Groundnut generally goes to France. Groundnut is imported into Bombay from Madras and the Nizam's territories and exported by rail to the United Provinces and the Punjab.

Groundnut cake is a rich cattle food containing about 45 per cent. albuminoids. It is also a very useful manure for sugarcane on the Nira and Godavari Canals, and usually contains from seven to eight per cent. of nitrogen.

The crop is grown as jirayat or dry as well as irrigated. It does well on light and medium soils. In the tracts receiving more than 40 inches of rainfall early groundnut, the seed of which is generally imported from abroad, does not require irrigation. Groundnuts are

chiefly cultivated in the western talukas of Satara, Poona and Nasik where most of the crop grown now consists of the early foreign varieties. Lately the crop—Spanish peanut variety—is extending in Khandesh and in some parts of Gujarat such as North Ahmedabad. In Khandesh the Spanish peanut is likely to take a place in rotation with cotton as the receipts of this crop compare very favourably with those of cotton.

Big Japanese, Pondichery, Spanish peanut and Small Japan are the principal foreign varieties grown in the Bombay Presidency. These were introduced by the Agricultural Department. Of these Spanish peanut and Small Japan are the early varieties and are generally grown without irrigation. The other varieties are later, though not so late as the country variety and require one or more waterings where the rainfall is scanty. Even these varieties are harvested about a month earlier than the country variety, and in the Satara district even Big Japanese and Pondichery are grown as a dry crop in the Western talukas.

Groundnut is a *kharif* crop and is sown with the first fall of rain. Early varieties are harvested in September and October and late ones in November and December.

The average outturn per acre varies from 800 to 1,600 lbs. of unhusked seed containing from 65 to 75 per cent. of kernels. The percentage of oil in the kernels comes to from 45 to 50. From an acre of groundnut about 800 lbs. of very good fodder is obtained.


Niger seed.—The total area under this crop is 202,798 acres, and of this nearly half is claimed by the Nasik District. Poona follows with about 15 per cent. Ratnagiri will come next; Satara claims about 7 per cent. Other districts grow less than 5 per cent. Gujarat does not grow niger seed, except a small area in the Panch Mahals.


Niger seed is a *kharif* crop. It does well on shallow soils, and in the Konkan and the *mallad* tracts it is rotated with coarse millets. In some parts of the country a light sprinkling of niger seed is found in the *bajri* crop.

The seed contains from 41 to 43 per cent. of oil and yields in the country ghani from 30 to 32 per cent. of oil, which is used for household purposes and especially in cooking. The oil cake makes a good concentrated food for cattle containing about 28 per cent. of albuminoids. The cake contains about 4 per cent. of nitrogen, and is found to be good manure for the sugarcane crop in the Nasik District.

The outturn per acre usually obtained is about 300 lbs.

Linseed.—The total area under this crop is 265,429 acres, and it thus takes the fifth place among the oil seeds in the Bombay Presidency. It is sown in the rabi season in deep, moisture holding, lands. It is generally grown alone and is the sole crop of the year. About 42 per





cent. of the total area is claimed by Bijapur. Nasik, Dharwar and Sholapur together make up about 40 per cent. while the remaining 18 per cent. of the area is found in other districts, except in the Konkan and in Gujarat where no linseed is grown.

The crop is grown for its seed which supplies exclusively oil and cake. In Europe and America the same plant is grown very largely as a fibre crop but this is never the case here. Most of our linseed is exported to the continent of Europe, and before the war this export amounted to about 130,000 tons per annum.

The seed contains from 37 to 44 per cent. of oil and yields about 32 to 38 per cent. in the *ghani*. The cake is among the most useful cattle foods but it is here also used as a manure.


A good crop with full even plant on deep black soil will yield about 500 lbs. seed per acre.

Castor seed.—The total area under castor is 89,235 acres, about three quarters of which is claimed by the Surat, Ahmedabad, Kaira and West Khandesh districts. The Panch Mahals has about 7 per cent. of the total area and Dharwar about 4 per cent.

There are a number of varieties of castor. The seeds of these contain from 45 to 52 per cent. oil and yield from 40 to 42 per cent. of oil in the country ghani. The oil is used as a lubricant and lamp oil in villages, and is largely in demand generally, as a lubricant, in soap making and in medicine. The oil cake is used as manure alone, and its quality is very variable according to the amount of husk left in the cake in pressing. When well decorticated it gives white castor cake containing about 6 to 7 per cent. of nitrogen but the quality more common in Bombay contains much husk and only about 4 per cent. of nitrogen? Castor cake cannot be fed to cattle as it causes first purging and then death. The stalks and the husk of the capsules are of no use except for fuel. Considerable quantities of castor cake are supplied by, oil mills for the sugarcane crop on the Deccan canals. There is a large export of castor seed from Bombay to the United Kingdom, the continent of Europe, and the United States of America.

The average outturn per acre may be taken at about 450 to 550 lbs. of seed.

Cocoanut.—The cocoanut palm is a tropical crop, being found in the countries and islands bordering on the Indian and Pacific oceans, and it is from the islands and sea coasts in the more tropical parts of these oceans that the world's supply is chiefly derived. Only the southern part of the Bombay Presidency is sufficiently tropical to grow cocoanut as a large commercial crop, and hence the total area under cocoanut is only about 28,000 acres, 50 per cent. of which is found in Ratnagri, 40 per cent. in Kanara and 3 per cent. each in the Thana and Kolaba districts.



XIV.—FRUITS.

The common fruits grown and used in the Bombay Presidency are plantains or bananas, mangoes, guavas, oranges and other citrus fruits, pomegranates, grapes and figs. The mango probably occupies a larger acreage than is ever returned by the talukas as in most places it is planted round the villages and on unoccupied land. With this qualification, the area under the different fruit crops is shown in the attached table, district by district. The following notes on the distribution of each of these crops may be useful.

Distribution of fruits in the Bombay Prisidency.

								•	
No.	Name of Dis- tricts.	Grapes.	Plan- tains.	Pome- granates.	Guavas.	Figs,	Oranges and Le- mons.	Mangoes,	Total.
1	Ahmedabad		95	252	293		114	68	822
2	Kaira		9	9	30	1	84	80	213
8	Panch mahais		8	21	4		6	¦	34
4	Broach		65	4	235	ļ	13	49	266
5	Surat	1	411	2	114	9	50	858	945
6	West Khandesh		157	1	97	••	. 88	141	484
7	East Khandesh	2	2,059	8	415	1	220	62	2,762
8	Nasik	836	255	19	672	1	67	903	2,181 .
9	Ahmednagar	- 56	807	254	601	5	1,399	98	2,720
10	Poons	57	1,029	742	620	426	1,785	166	4,825
11	Sholapur		1,032	58	158	٠	84	167	1,499
12	Satera	5	1.555	77	247	••	75	68	2,027
- 18	Belgaum		683	7	128		20	971	1,807
14	Bijapur	l !	1,068	:	66		88	349	1,516
15	Dharwar	 	476	8	442		52	1,824	2,797
16	Thana	l !	1,651				10	1,971	3,632
17	Kolaba		89				8	880	422
18	Ratnagiri		167 331		1 2			535 1,069	703 1,393
19	Kanara		5,200		8	••	4	1,205	6,412
	Total	457	16,261 331	1,452	4,124	443	4,171	9,395 1,060	86,308 1,393

Plantains or Bananas.—The total area under plantains in the Bombay Presidency is 16,218 acres. The largest centres of production are located in the districts of Kanara, East Khandesh, Thana, Satara, Bijapur, Sholapur and Poona each having an approximate area of 1,000 acres. In Gujarat and the Konkan (except in Thana) the crop occupies very little area. The fruit is very popular among all

nationalities and considerable quantities of plantains are exported from Bombay to Karachi and the Punjab.

The chief centres of cultivation are as follows:-

- (1) Near Bassein in the Thana District.
- (2) Sirsi and Siddapur in the Kanara District.
- (3) Pachora, Jamner, Bhusaval, Raver, Jalgaon, (Asoda), and Chalisgaon in East Khandesh.
 - (4) Khatav, Walva, Wai and Karad in the Satara District.
 - (5) Bagewadi and Bijapur in the Bijapur District.
 - (6) Sholapur, Madha, and Barsi in the Sholapur District.
 - (7) Junnar and Khed in the Poona District.

Some of these tracts are handicapped for want of transport facilities and though they are more suitable than others, transport absorbs so much of the profit that plantain cultivation cannot extend as would otherwise be the case.

In most parts the crop occupies the land for about two years (i.e. for one harvest) though in some parts it is kept from four to ten years on the same land.


Mangoes:—This is perhaps the fruit par excellence of India, and the mangoes of Bombay are unrivalled in any other part of the country. Except in the sandy parts of Gujarat it is grown in all districts fairly generally, but the climate of the Konkan seems the ideal for the best varieties. These best varieties which are always in demand at a good price are the Alphonso or Hapus and the Payari. In the Konkan the mango begins to bear in the fifth year, but elsewhere it takes longer, but once established a mango plantation lasts for generations.


Fairly large quantities of mangoes from Madras and Goa find their way to Bombay, while large quantities leave Bombay for the Punjab, Sind, Baluchistan, and even for England and France.


Guavas:—The total area under guavas is 4,111 acres. The crop is well spread in the districts of the Deccan and Karnatak. It practically finds no place in the Konkan, and of the Gujarat districts, only Ahmedabad returns a fair area. Nasik, Poona, and Ahmedagar grow between 600 to 700 acres each. Dharwar and East Khandesh over 400 acres. The fruit is very popular with the Indian section of the population, and the Bombay Presidency does not export guavas. On the other hand, it imports this fruit from Agra and Delhi.


The crop is concentrated in the following talukas or their parts:-

- (1) Dholka and North Daskroi of Ahmedabad.
- (2) Broach and Ankleswar of Broach.
- (3) Chorashi of Surat.









this lies

area else

large cer

This

Tain

by beteli entirely s

detailed !

(4) Jalgaon of East Khandesh. (5) Nasik and Malegaon of Nasik (6) Pathardi and Kopergaon

Ahmednagar.

(7) Haveli of Poona. (8) Gadag and Dharwar of Dhar

Oranges and other citrus fruits.—T and other citrus fruits is 5,146 acres, 35 per cent. and Ahmednagar 27 per cent desh, Nasik, and Ahmedabad grow betwo. each. Oranges are not grown in the Konkan and extent in Gujarat. On the other hand there is a fair at loes in the Thana district. But on the whole, the Bombay Presidency is an importing centre for citrus fruits, receiving oranges in particular

Citrus fruits are concentrated in the following talukas or their parts:-

small export trade to the Punjab, Kathiawar and Karachi.

from Nagpur, East Africa, Italy and Persia. There is however a

- (1) North Daskroi of Ahmedabad.
- (2) Chalisgaon of East Khandesh.
- (3) Rahuri, Shrigonda, Ahmednagar and Parner of Ahmednagar. (Rahuri, alone claims about 500 acres.)
 - (4) Purandhar, Haveli and Sirur in the Poona District.

Pomegranates.—The total area under pomegranates is about 1,451 acres. More than half the area is claimed by the Poona district, where most is grown in the Haveli, Bhimthadi and Purandhar talukas. The Dholka taluka of the Ahmedabad district claims 15 per cent. of the total area in the Bombay Presidency. The Ahmednagar district grows about 15 per cent. most of it being claimed by the Pathardi, Rahuri, Ahmednagar and Sangamner talukas. Except in a few places like the Dholka taluka and a few villages in the Poona district the fruit is not of a superior quality. Fruits of better quality are imported from Muskat, Harvai (Baluchistan) and Peshawar.


Grapes.—The total area under grapes is only 457 acres, threefourth of the area being concentrated round about Nasik and its subburbs. Of the remaining area about one-half is found at and near Ahmednagar and nearly half in the Junnar taluka of the Poona district. Grapes from Nasik are sent to Bombay, Poona, Calcutta, Lahore, Agra and other cities; while some fruit comes into the Bombay Presidency from Quetta, Malta and even from Spain. The cultivation of grapes is extending very fast at and near Nasik, which is found particularly suitable for this crop, and also in Khandesh, especially near Jalgaon (Nasirabad).


is.—The total area under figs is about 443 acres. Most of in the Purandhar Taluka of the Poona District. In fact, the awhere is negligible. But from the Purandhar Taluka all the—tres are supplied by passenger train.


XV.—SPICES AND CONDIMENTS.

group is represented in the Bombay Presidency principally ut, cardamoms and black pepper. These crops are almost grown in the Kanara, Ratnagiri and Kolaba Districts as below:—

1		B .	Betelnut	Cardamon	Black Pepper	
			Acres.	Acres. Kolaba District.	Acres.	
	··	ŀ		Kolaba District.		
	Alibeg	••	370		••	
2	Panwel		1	1	••	
_		Total	871			
				Ratnagiri District.		
1	Vengurla		1,308		••	
2	Maivan		398			
8	Deogad		105			
4	Rajapur		254		••	
5	Ratnagiri		263		••	
6	Sangameshwa	r	15			
-7	Chiplun		117		••	
8	Khed		3	.	••	
9	Dapoli		1,358			
_	· 	Total	3,821		••	
				Kanara District.		
1	Karwar		33		••	
2	Ankola	•	880	. 7	26	
8	Kumpta		654		58	
4	Honavar	•	864	\	39	
5	Siddapur	••	383	736	2,123	
6	Siral		5,795	550	1,620	
7	Yellapur	•	2,419	100	297	
8	Haliyal	•-	169	2	2	
		Total	18,019	1,395	4,160	







Betelnut.—The total area under Betelnut is about 22,211 acres, most of it (practically 82 per cent.) being claimed by Kanara. Ratnagiri grows 3821 and Kolaba 371 acres.

Cardamons.—The total area under cardamons is 1,395 acres. All of it is found in the Kanara district, where the Siddapur and Sirsi talukas claim 78 per cent. of the area, while the Yellapur taluka has about 100 acres under this crop. Almost all this crop is consumed in the country.

Black Pepper.—The total area under pepper is 4,160 acres. Half of the area is grown in the Siddapur taluka of the Kanara district, while Sirsi grows 1,620 and Yellapur 297 acres. Other talukas claim less than 60 acres each. Other talukas of the district practically do not grow pepper.

XVI,-VEGETABLES AND ROOT CROPS.

The market, soil, climate, supply of water and habits of the people in each locality are very important factors in determining whether vegetables shall be cultivated and also what vegetables shall be grown. The following are the principal root and vegetable crops grown in the Bombay Presidency:—

Roots and Bulbs.	Leafy vegetable.	English vegetables.	Other vegetables.
Potatoes	Methi (3). Math (4)	Calbage.	Brinjals. (8)
Sweet-potatoes	Rajgira. (5)	Cauliflower.	Chillies.
Suran (1) or Elephant foot.		Knol-kol or Kohl-rabi.	Bhendi. (9)
Goradu (2) or Yam	ı	Beet. ·	Chavli. (10)
Turmeric .	Chakwat. (6) Alu. (7)	Tomatoes.	Cucurbits.
Ginger.		i	
Oniona.			
Garlic.			
Carrots.			· ·
		1	1

^{1.} Amorphophallus campanulatus. 2. Dioscorea. 3. Trigonella foenumgraecum. 4. Amarantus gangeticus. 5. Amarantus paniculatus 6. Chenopodium album. 7. Colocasia antiquorum. 8. Solanum melongena. 9. Hibiscus esculentus. 10. Vigna catiang.

D A 5 5-7

Potatoes.—This crop has been previously dealt with. (See page 35.)

Sweet-potatoes.—Sweet-potatoes are grown in small patches all over the Bombay Presidency, though not very extensively. The total area is in fact only about 12,000 acres. The crop, however, does well on almost any kind of soil. There are two main varieties, white and red, the latter being more common. The crop is grown chiefly in the cold season though in some places it is cultivated also in the rains and even in the hot weather.

Suran or Elephant foot.—This crop is important in certain parts of Gujarat, especially in the valleys of the rivers Tapti, Purna and Ambika in the Surat district. It does not appear to thrive well in the Deccan. In Gujarat the cultivation is very careful and methodical. It is in fact a four year crop though the crops are harvested and replanted every year. The crop requires porous, well drained and rich soil. The produce is stored by Surat merchants in cellars and sold locally or sent to Bombay. More than three quarters of the suran from this tract goes to Bombay, though there is a fair local demand.

Turmeric, Ginger, Goradu Chillies, etc.—In certain parts of Gujarat, especially in the Ambika and Purna valleys, in the Jalalpur taluka of the Surat district, these crops are sown as a mixture. The system and method of planting is very methodical, a plant of each crop having a definite place in each bed. The cultivation is very intensive and regular rotation is followed. The crop is planted about the middle of May. In addition to rain water it requires irrigation from fifteen to seventeen times, the first five times in May and June and the remainder between October and January. The crop is harvested in February. Many variations of this method of cultivation are found in this part of Gujarat, certain of the constituents of the mixture being omitted, or even (like turmeric) being often grown alone. The cultivation is always, however, equally careful, and among the most intensive in India.

Turmeric is a very important crop in many parts of Satara and adjaceet districts in the Deccan, especially in the Krishna valley, being grown as a single crop. The same is the case both in the Sholapur and Belgaum districts, though not to the same extent.

Ginger is grown as a pure crop in parts of the Kaira and Thana districts, and in parts of Baroda State. Ginger, turmeric, and yams require, like *suran*, rich free working garden soils. Most of these crops, as grown in Gujarat, are marketed in Bombay.

Onions.—This crop is very important in the Deccan notably in the north of the Poona district, and in several talukas of the Nasik district,

notably, Niphad, Sinnar, Dindori and Nasik. It is also grown in certain parts of the Satara district (Walva taluka), and in West Khandesh near Dhulia. In Gujarat it is purely a crop of garden lands. In the Konkan and in Kanara its cultivation is now tending to spread.

Onions like free working, well drained soil of fair depth. They are usually taken as a cold weather crop, but have been found to do well even in the rainy season and in the hot weather.

Garlic is an important crop in the Sinnar and Niphad talukas of the Nasik district where it is very carefully cultivated. Elsewhere it is found in small areas all over the Bombay Presidency, the most important centres apart from Nasik, being Dharwar, Belgaum, Poona and Satara. It is a semi-cold weather crop grown under irrigation.

Other Vegetable Crops.—Little need be said about other vegetable crops. The most important of these are Brinjals, Bhendi, Beans of various kinds, and different varieties of cucurbits. Brinjals are grown almost everywhere, in garden land, but the chief cultivation is in Gujarat, particularly in the goradu or besar lands of Kaira district and the Baroda State. Beans, as vegetables, are also universally grown while cucurbits are found round a large proportion of rural dwellings, and are always in keen demand in all cities and towns.

English Vegetables, such as cabbage, cauliflower, beet root, and tomatoes have a very good market in all the cities of Western India, and specially in Bombay. These vegetables are usually grown in the cold weather in free working, well manured soils. Poona, Belgaum, and Nasik and other surrounding places are well known centres which supply these to Bombay and other markets.

XVII.—YIELDS OF CROPS.

The statement of the normal outturn of principal produce per acre given on page 53 is taken from the officials records of the Bombay Government. These yields are said to have been fixed without elaborate and careful tests many years ago and the same have been handed down with slight modifications made at revisions conducted every five years. The normal outturn is supposed to be that which a cultivator has a right to expect with good cultivation in a normal year.

Another statement (page 56) of yields is worked out on the basis of the estimated actual outturns of crops for a number of years. These estimates are formed annually for every district, on the basis of the condition of the crops during their period of growth. It will be noticed

that these are in almost every case considerably below the normal outturn as defined above.

A third statement (page 58) shows the average yield of such important crops as give fodder in addition to grain. The procedure followed in order to obtain these figures is as follows:—From the material available on the Government farms (from the year 1904 to 1920) the proportion of fodder to grain for each crop has been fixed separately for each division of the Bombay Presidency. This having been done, the amount of fodder is calculated on the bases of the average yields, given on page 54:—

Statement showing the normal outturn per acre of stated crops of each district.

(Taken from the season and crop report quinquennium ending on 1917.)

, ***	Ì	Name of crops.										
Name of District		Jowar.	Bajri.	Rice.	Wheat.	Gram.	Tur.	Udid mug and math.	Maize			
,		Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.			
Ahmedabad		1,080	840	1,440	560	500	560	510	1,08			
Kaira		1,050	970	1,320	600	500	690	560	1,05			
Panch Mahals		1,050	825	1,200	700	600	570	510	1,19			
Broach		1,160	700	900	600	500	600	625	1,16			
Surat	•	760	600	1,560	560	500	680	640	1,16			
West Khandesh		650	500	1,080	600	500	960	600	72			
East Khandesh		650	500	1,080	600	500	960	600	72			
Nasik	••	520	860	1,080	460	850	960	520	52			
Ahmednagar		540	340	1,040	460	880	960	500	54			
Poona	••	500	840	1,120	850	820	800	510	50			
Sholapur	• • •	540	320	900	400	860	960	520	50			
Satara	••	720	360	1,120	480	880	1,200	550	72			
Belgaum	••	800	400	1,140	560	400	960	510	80			
Bljapur	••	540	820	800	400	840	960	500	54			
Dharwar		1,000	450	1,140	500	500	1,200	540	90			
Thana	••	700	840	1,200	400	400	870	400	50			
Kolaba ,	••			1,820	400	880	410	470	. 50			
Ratnagiri	••		400	1,020		820	480	840	80			
Kanara	••	900	450	1,820		480	560	500	60			

		Names of crops.									
Name of District.		Fagli,	Varts	Kodra.	Cotton.	Sugar- cano.	Ground- nut,	Lin- mod.	Seen-		
	-	Lbs.	Lba.	Lba.	Lba. Lint.	Lbs. Gul.	Lbs.	Lbs.	Lbs.		
Ahmedabad	••	1,440	1,350	1,320	125	6,000	[850	400		
Kaira	••	1,440	1,830	1,275	115	6,000	••	••	400		
Panch Mahale	••	1,420	1,290	1,139	100	5,000		360	400		
Broach	••	1,500	1,025	1,025	130	7,000	2,500	••	450		
Burat		1,200	1,140	1,120	120	7,000	2,500	••	400		
West Khandesh	••	900	800	908	110	7,000	2,500	360	400		
Bast Khandesh	••	900	800	900	110	7,000	2,500	360	400		
Nasik		850	840	806	100	7,000	2,500	360	400		
Ahmednagar		900	8 2 0	800	90	7,000	2,500	360	400		
Poons.		900	850	800	90	7,000	2,500	360	40		
Sholapur	••	900	800	800	90	7,000	2,500	360	400		
Betera	••	900	820	800	. 80	7,000	2,500	360	40		
Belgaum .		900	840	800	100	7,000	2,500	360	40		
Bljapur	••	900	700	800	90	7,000	2,500	\$60	40		
Dharwar	•• ;	1,200	920	800	120	7,000		360	40		
Thans	••	770	1,050	640		7,000			30		
Kotaba		800	1,140	800		4,000		••	80		
Ratnagiri		680	900	675		4,000			30		
Kanara		1,470	1.530			6,000			80		

54

Statement showing the average outturn per acre of stated crops of each district.

•					Names	of crops	•		
Name of District.		Jowar. Bajri.	Rice.	Wheat.	Gram.	Tur.	Udid Mug and Math.	Maize	
		Average	of 14 ye	are from	1906 to	1920 —	Average	of 10 ye	ars
		Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.
Ahmedahad		601	618	951	385	331	274	284	84
Kaira		661	551	1,135	893	306	435	302	604
Panch Mahals		736	597	1,029	419	352	437	812	855
Broach		725	412	567	438	320	358	428	644
Surat		861	485	1,297	431	364	540	469	880
West Khandesh	••	471	822	776	418	370	770	394	531
East Khandesh		615	318	646	460	386	657	450	475
Nasik	••	281	236	824	317	235	589	877`	35
Ahmednagar	•-	811	209	784	263	194	534	325	400
Poona	••	309	217	929	278	202	430	297	29
Sholapur		845	137	468	240	173	493	209	27
Satara	••	532	228	787	329	248	732	366	47
Belgaum		578	247	. 850	383	260	583	314	58
Bljapur	••	865	184	475	278	217	518	272	48
Dharwar	••	678	285	784	898	839	792	867	57
Tohna	••	423	202	1,073	805	814	274	291	40
Kolaba	••	465	289	1,255	••	286	335	400	43
Ratnagiri	٠.,	600	239	830	••	207	264	214	62
Kanara		437	225	1,007		264	221	285	46

	ì		·		Mazzes of	f erops.			
Name of District.		Nagil.	VarL	Kodra.	Cotton.	Sugar- cane,	Ground- nut.	Lin-	Sesa-
		from 19	09 to 191	8-19.				·;	
		Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.	Lbs.
Ahmedabad	••	676	718	706	84	4,988	2,229	214	264
Kaira	••	985	880	796	70	4,373	1,302	196	228
Panch Mahale	••	1,045	794	865	60	3,104	1,860	224	277
Broach	••	912	543	642	100	4,696	2,400		302
Surat	••	1,082	988	1,000	90	6,130	2,253		322
West Khanderh	••	72	548	495	83	5,540	2,394	290	282
Sast Khandesh	••	609	555	659	77	5,825	2,633	259	318
Nasik		674	629	379	69	6,393	2,101	271	245
Ahmodorgar	••	711	585	. 546	61	6,297	2,107	247	287
Poons.	••	735	613	531	49	6,023	2,425	189	200
Sholapur	••	374	412	347	42	5,776	1,912	223	180
Satara	••	636	554	476	56	5,818	2,341	228	270
Belgaum	••	677	582	476	69	6,084	2,248	259	256
Bijapur	••	458	420	408	58	5,488	1,249	257	200
Dharwas	••	850	632	476	89	5,761	2,214	264	248
Chana		574	825	460	••	6,325		••	351
Cotaba	••	689	973	701	••	8,700		••	827
Satnagiri		493	643	480	••	8,333	1,822	••	**
Canara		917	858		••	5,756		••	201

I.—Statement showing the average grain yield (in the case of sugarcane gul) of different crops in different tracts in Bombay Presidency.

			· 					
No.	Kame of crop.		Gujarat.	Khandesh.	Central Division.	Karnatak.	Konkan including Karwar.	Remarks
1	Jowar		716-8	543-0	355-6	540.3	481 - 25	
2	Bajri		512-6	320 0	205-4	238 7	238 · 75	
3	Wheat		413-2	439.0	285 4	353-0	305.00	
4	Rice	٠.,	995-8	711.0	738-4	703-0	1,041 · 25	
5	Gram		334 · 6	878-0	210-4	272.0	267-75	
6	Groundnut	••	1,000-0	1,000-0	1,000.0	1,000-0	1,000.00	
7	Sugarcane	••	4,658-2	5,682.5	6,061-4	5,761.0	4,778 50	
8	Maize		767-0	501.5	358 6	546.0	479-50	
9	Nagli		940-0	837-5	626.0	661-7	668-25	
10	Tur		408-8	713.5	555 · 6	631.0	273 · 00	
11	Other pulses		859-0	368-4	368-4	317-7	297-50	
12	Other ceresis		792-2	535 1	535-1	499-0	685 · 00	
13	Sesamum		278-6	300-0	226.0	235.3	293.00	
14	Fodder crops		••	ļ	l			

II.—Statement showing the average yield of dry fodder of different crops in different tracts in Eombay Presidency (in the case of sugarcane yield of green tops is taken).

Lbs.

			,			,		
No.	Name of crop.		Gujarat.	Khandesh.	Central Division.	Karnatak,	Konkan including Karwar.	Remarks.
1	Jowar		1,792 · 0	1,221 · 75	800-10	1,782 99	1,082 82	
2	Bajri		720.0	720.0	595-66	595 · 66	658 80	
8	Wheat	• •	909-04	557-58	428-10	529 · 50	887:35	
4	Rice		1,414.08	1,306 - 73	1,305 - 78	1,305 · 73	1,197-44	
5	Gram		361 - 87	275.94	189-36	198-56	198-56	
6	Groundnut		2,200.00	2,300 - 00	2,400-00	1,200.00	1,500.00	
. 7	Sugarcane		6,055 66	7,887 · 25	7,879 · 82	7,849 00	6,212.05	
8	Maize	••	1,150-5	752 - 25	587.9	819-00	719-25	
9	Nagli		2,256.0	810.0	1,502 · 4	1,852 76	1,269 68	
. 10	Tur		725-84	1,070 - 25	777 - 84	757-2	856-55	
11	other pulses		574-4	589-44	589-4	508-82	476-00	
12	Other cereals	[1,584-2	1,070-02	1,070 · 2	998-0	1,871.8	
13	Sesamum		834-82	360-0	271.2	282-36	851-6	
14	Fodder crops		5,000.00	5,000-0	5,000.0	5,000-0	5,000-0	

XVIII.—IRRIGATION.

Water is indispensable for every kind of product of the soil. Most crops require at least 350 times as much water during their growth as the dry material (including straw) they produce. Where there is any doubt of the water which falls in rain being sufficient to supply the necessary quantity, irrigation is resorted to in almost all countries. But in countries like the Bombay Presidency where the rainfall is not only very unequally distributed throughout the year, and even during the period when a crop is grown, but is also very uncertain, irrigation becomes an essential in raising many crops, and of advantage, if available, with nearly all.

Considerable parts of the Bombay Presidency, especially the districts of Ahmednagar, Poona, Sholapur, Bijapur, parts of Nasik, Satara, Dharwar and Belgaum, suffer periodically from famine caused by the failure of rainfall and in these areas irrigation, from whatever source, is of extreme value where it can be obtained.

The great advantage of irrigation has been recognised for many centuries, and seems to have been realized by the early rulers and communities ages ago. The very large number of wells, and the old bandharas, or embankments across the rivers found in the country (especially in the Nasik, Khandesh and Satara districts) give testimony to this. The old works, mostly communal, were small but efficient. The British Government has constructed large irrigation works and have projected still larger canals. Many once arid regions, like the tract between the Nira river and Baramati in the Poona district, now smiling with beautiful crops and a well fed population show the benefit of the money invested in the works, and even though the canals may not be very remunerative directly to the Government the indirect advantages are far reaching, and they warrant the execution of bigger schemes.

The following is the list of canals which supply water for perennial crops like sugarcane:—

1.	Pravara	in the	Ahmednag	ar District
2.	Godavari	99	,	,,
3.	Nira	,,,	Poona	29
4.	Mutha	,,	>>	,,
5.	Krishna	"	Satara	•,
6.	Girna	"	Nasik	**
7.	Ozar	,,	**	,,
8.	Kadva	15	خذ	**
65	8			

D A

- 9. Hatmati in the Ahmedabad District.
- 10. Ekruk ", Sholapur
- 11. Godak` ,, Belgaum

In old times private and communal bandharas served a very useful purpose, and even now the Nasik district den onstrates how communal bandharas are efficiently managed with the least possible waste of water. There are as many as 296 bandharas in that district (185 of them being only on 17 rivers) and they have been accustomed quite recently to supply-water for about 30,000 acres. Similar bandharas are seen in the West Khandesh and the Satara districts, but not in such large numbers.

",

The area in the Bombay Presidency that is irrigated by wells, tanks, private canals, and Government canals is given in the statement which follows (page 60) from which it will be seen that in the year 1914-15 the following acreage was irrigated.

175,419 acres by Government canals.

17,788 , private canals.

490,023 ,, wells.

131,318 , tanks and storage lakes.

88,987 , other sources.

903,535 average total area irrigated.

The table attached shows the total area cropped and the area irrigated from different sources. It also indicates the percentage of the irrigated to the total cropped area. Taking the Bombay Presidency as a whole, this percentage only come to 3.6. Of the irrigated area, about 54 per cent. is served by wells, 26 per cent. by tanks or storage lakes, private canals and other sources, and 19.4 per cent. by Government canals.

Irrigation wells are particularly frequent in the Satara, Poona, Nasik, Ahmednagar, Sholapur, East Khandesh, Ahmedabad and Kaira districts. Kanara, Dharwar, Kaira, Belgaum and Surat have a larger number of tanks than the other districts. Nasik stands first in the number of bandharas, while Government canals serve the largest areas in Poona, Ahmednagar and Nasik districts.

59

Irrigated area in the Bombay Presidency.

						Area :	lrrigated.	•
Name of District.		Net croped area under cultivation.	Total irrigated area.	Percent-	Government Canals.		Private Canals.	
•	_				Area.	Percent- age.	Area.	Percent- age.
		Acres.	Acres.		Acres.		Acres.	
Ahmedabad		1,103,923	78,741	7-13	17,075	21.7	••	•/4-
Kaira		634,427	89,753	6 · 26	6,726	16.9	••	
Panch Mahala	••	420,206	8,761	0.89	731	19.4]
Broach .	••	556,216	448	0.08				
Surat	••	453,181	9,775	2.15	2	0.02	••]
West Khandesh		1,390,912	26,405	1.93	10,977	41.5	2,160	8.1
Rast Khandesh		1,976,077	26,878	1.36	1,705	6.8	154	0.6
Nask		1,986,370	93,499	4.70	83,486	85-8	3,792	4.05
Ahmednagar		2,755,054	85,900	8·11	11,989	13.9		
Poons.	,	1,896,506	132,929	7.08	60,665	45.6	8,538	6.4
Sholapur		2,050,240	100,075	4.88	12,906	12.8		•••
Satara		1,502,464	97,772	6.20	8,452	8.6	620	0.6
Belgaum		1,673,884	51,871	8.09	6,000	11.5	429	0.8
Bijapur		2,947,451	17,774	0.80	140	0.7	266	1.5
Dharwar		2,102,710	94,815	4-50	4,565	4.8	1,451	1.2
Thana		442,281	6,155	1.39			••	
Kolaba		848,841	1,806	0.51		' [••	 .
Ratnagiri		168,793	11,089	6.28	••		56	0.2
Kanara		210,826	24,089	11.42	••		322	1.3
Total		24,619,862	903,535	3.66	175,419	19.4	17,788	1.9

				Area Irriga	ited.		
Name of District		Welli	5.	Tank	38.	Other Sources.	
	• 1	Area.	Percent- age.	Area.	Percent-	Area.	Percent-
		Acres.		Acres,		Acres.	
hmedabad		40,090	50.9	19,378	24 6	2,203	2.8
Kaira		29,814	74.9	2,919	7.3	294	0.7
Panch Mahals		2,891	76.8	135	8.6	4	0.1
Broach .		448	100.0		••	••	
Surat		5,657	57-8	8,945	40.8	171	1.7
West Khandesh		12,300	46-6	243	0.8	725	2.7
East Khandesh		24,126	89-8	58	0.2	840	3.1
Nasik		49,516	52.9	1,217	1.8	5,488	5.8
Ahmednagar		66,137	76.9			7,774	9-05
Poona `		57,666	43.8	8,196	2.4	2,864	2·1
Sholapur		86,655	86-6	••	••	514	0-5
Satara '	•	50,696	51.8	1,904	1.9	36,100	36-9
Belgaum		82,112	61.9	8,792	16.9	4,538	8.7
Bijapur		16,130	90.7	842	4.7	396	2.2
Dharwar		3,202	8.8	76,086	80.2	9,511	10.03
Thans	[5,087	82.6	2	••	1,066	17:8
Kolaba		1,495	82.7	5	0.2	306	16.9
Ratnagiri		5,302	47.8	910	8.2	4,821	43-4
Kanara ·		699	2-9	11,696	48-5	11,872	47.2
Total	ai	490,028	54.2	181,318	14.5	88,987	9.8

XIX.—LIVE STOCK.

The close relationship between the crops and the live-stock makes it necessary to add one or two tables showing the numbers and distribution of the principal domestic animals in the parts of Western India with which we are dealing.

The live-stock of the Bombay Presidency consists of the common farm animals, cattle and buffaloes, sheep and goats, poultry and a comparatively small number of horses, asses and mules.

The following statement shows the total number of each class as returned in the year 1915-16:—

Cattle including buffs	••	91,57,395	
Horses and Ponies	••	••	1,19,619
Asses and Mules	• •	• •	99,877
Sheep and Goats		9-0	40,13,668

Of the total number of cattle about 37 per cent. are working cattle, 32 per cent. cattle kept for milk, the remaining 31 per cent. being young stock, stud bulls, etc.

The proportion of males to females for cattle and buffaloes is shown below:—

		Cattle.	Buffaloes,		
Males	62	per cent.	20	per cent.	
Females	38	per cent.	80	per cent.	

is chiefly kept as a work animal on the farm.

This shows that the cultivator gives more care to the male calf of a cow and the female calf of a buffalo. The excess of male cattle is increased by a very considerable import of bullocks and bull-calves from the Malva plateau in Central India, and from the Mysore plateau in the south. Nevertheless the above figures illustrate how the buffalo is primarily the milch animal of the country, while the bullock

Work cattle.—Oxen and he-buffaloes, as measured by total value and universal service, form the most important class of live-stock on Indian farms. Steam and oil engines are still curiosities and are only found here and there for pumping water. Horses and mules are scarcily used in connection with agriculture. The road traffic is almost entirely carried by means of bullock-carts.

Oxen form about 93 per cent. of the work-cattle and buffaloes in the Bombay Presidency, the remaining being he-buffaloes. In Gujarat the proportion of oxen to he-buffaloes comes to 100 to 1, in the Central Division to 100 to 3, in the Karnatak to 100 to 8, and in the Konkan to 100 to 29. The he-buffalo is found more useful for rice

cultivation, and the small number of he-buffaloes, used in Gujarat, in the Central Division and in the Karnatak are found chiefly in the rice-growing tracts of these divisions.

The attached table shows among other things, the acreage which can be worked by one pair of cattle in each district. The statement shows considerable variation. In the Ratnagiri district the average area per pair of work animals is only about three acres, while in Bijapur it reaches thirty-one acres. It can be roughly said that the Konkan and the rice-tracts near the ridge of the Western ghats require a larger number of working cattle than the mallad tract, while desh tract needs less than the mallad tract. In the Konkan and in the kharif tract of the desh the land is more intensively worked than in the rabi tract. The latter, moreover, having deep soils, does not require such frequent ploughing.

Milch Cattle.—The buffalo is essentially the milk animal of India, but although there are certain milch breeds of buffaloes which give fair quantities of milk, on the whole the buffalo as a class gives very little milk compared with the yields obtained from cows in Europe and America. The cow again, is kept more for her bull calf, though she contributes a scanty supply of milk. Unlike other countries where cattle are valuable as producers of beef as well as producers of milk, cattle in the Bombay Presidency are not stall fed to any considerable extent. Milch cattle usually give milk only a few months in the year, and calve about once in two years. Hence their dry periods are long, and even when in milk they are generally poor milkers, while, in fact, a very large proportion of them are not milked at all, their sole utility being as producers of bull calves and of manure. It is, therefore, very difficult to form any approximation of the average milk yield obtained, but an annual production of about 800 lbs. of milk per buffalo and 300 lbs. per cow (apart from the amount consumed by the calf) may be taken as an average for the Bombay Presidency.

If we take the milch animals recorded in the quinquennial cattle census of 1915-16 and the population of 1911 (since the figures of the census of 1921 are not yet available) the quantity of milk produced per head of population comes to about one-eighth of a pound. Taking into account the large quantities of milk and *ghee* absorbed by cities like Bombay and Ahmedabad, it will be seen what an infinitesimally small quantity is available to the poor in other parts of the country.

Fodder supply.—Mr. Keatinge, the late Director of Agriculture, has recently given the following rough statistics of the land in the Bombay Presidency.

(1) Cultivated area	••		24 1	million	acres.
(2) Uncultivated occupi	ed area		7	"	,,
(3) Unoccupied waste			1	: >	,,
(4) Forest	••		7	,,	•
(5) Not available (road	ds, build	lings,			•
water, etc.)	• •		5	••	

It is not easy to calculate how much fodder (2), (3), (4) and (5) will supply. The statements at the close of this note (page 69) give the estimated average quantity of fodder available annually from important crops in each Division, from which it is seen that the cultivated crops of the Presidency supply about 6 lbs. of fodder daily to each cattle and horse. The daily ration of fodder per head is greater than this, being about 10 pounds in the Karnatak, about 6 6 pounds in the Deccan, 4 4 pounds in Gujarat, and only 2 7 pounds in the Konkan. The balance required has, of course, to be made up,—so far as it is made up at all by grass grazing. It may be remembered that this ration of fodder has to be made up by grazing on privately owned and Government waste lands, on forest grass lands and on communal grazing areas (quracharan) to some extent.

How far these resources are capable of filling the additional requirements of the animals it is, as already noted, difficult to say. But the unequal distribution of these grass and grazing resources brings about the result that while in some areas (particularly in the mallad tracts, in upper Gujarat, and in the Satpuras, there is in normal years plenty of grass, in other parts such as the eastern desh tract, and many parts of the Konkan, there is always a scarcity during a considerable part of the year. There is, in fact, an immediate and urgent need in many of these areas to grow and to conserve more fodder. This is especially the case in view of the uncertainty of the rains in many of the regions where the fodder supply is shortest, and where the future development of agriculture is most closely bound up with the maintenance of the agricultural animals in good condition.

	·	•	Catt	le.	•		
No.	Name of District.	Ploug	h Cattle.	Oxen and buffaloes for other purposes.			
		Oxen.	He-buffaloes.	Oxen.	He- buffaloes.		
1	2	8	4	5	6		
1	Ahmedabad	107,177	358	2,337	350		
2	Kaira	94,290	15	817	26		
3	Panch Mahais	123,642	32	663	43		
4	Broach	56,702	10	837	121		
5	Surat	107,989	4,540	4,652	46		
6	West Khandesh	161,110	2,080	6,167	312		
7	East Khandesh	212,038	417	9,643	1,314		
8	Nasik	249,790	25,698	10,630	. 827		
9	Ahmednagar	280,750	5,152	5,111	575		
10	Poona	247,976	5,411	6,246	659		
11	Sholapur	173,867	3,622	5,241	1,888		
12	Satara	209,770	· 11,261	3,282	690		
13	Belgaum	167,914	25,190	3,745	859		
14	Bijapur	184,521	2,244	3,551	1,486		
15	Dharwar	206,323	13,739	5,046	616		
16	Thana	147,177	43,012	3,669	415		
17	Kolaba	92,376	42,580	2,335	314		
18	Ratnagiri	98,349	17,725	805	130		
19	Kanara	80,514	21,300	4,387	1,799		
	Total	3,000,275	224,336	79,164	12,470		

This statement does not include any cattle or area from the villages for which statistics are not regularly prepared. The area worked by one pair of cattle is therefore not affected any way.

No.	Name of Di	strict.	Total	Net area cultivated in the corresponding year 1914-15.	Area worked by a pair of cattle.	Remarks
1	2		7	. 8	9	
1	Ahmedabad		110,222	1,103,923	20.00	
2	Kaira		95,148	634,427	13.83	
8	Panch Mahala	••.	124,380	420,206	6-75	
4	Broach	•••	57,670	559,216	19-39	
5	Surat	••	117,227	453,181	7.73	
6	West Khandesh	•	169,619	1,390,912	16.40	
7	East Khandesh		223,412	1,976,077	17-68	
8	Nasik	••	286,945	1,986,370	12-84	
9	Ahmednagar	••	291,588	2,755,054	18.89	
10	Poons		260,292	1,896,506	14-57	
11	Sholapur	••	184,618	2,050,240	22 · 21	
12	Satara		225,003	1,502,464	13.35	
18	Belgaum	••	197,708	1,673,384	16.92	
14	Bijapur		191,802	2,947,451	30.73	
15	Dharwar		225,724	2,102,710	18.63	
16	Thens	••	194,273	442,281	4.55 ~	
17	Kolaba	••	137,605	848,841	5.06	-
18	Ratnagiri		115,009	168,793	2.93	
19	Kanara	••	108,000	210,826	3-90	
		Total	8,316,245	24,619,862	Average 14.85	

Statement showing the quantity of fodder available per head of cattle per annum as well as per day.

No.	Names of Districts.	Name of Division.	Total quantity of fodder available dur- ing the period of one year.		Quantity of fodder available per animal per annum.	Quanti- ty of fodder avail- able per animal per day.	Remarks.
1 2 3 4 5	Ahmedabad Kaira Panch Mahals Broach	-Gujarat	2,937,799,308	1,840,066	1,597	4.4	The figures showing animals have been taken from crop and season report of Bombay Presidency for the year 1915-16.
1 2 3 4 5 6 7	East Khandesh West Khandesh Nasik Ahmednagar Poona Sholapur Satara	Deccan	10,439,042,645	4,332,462	2,409	6-6	The figures about the about the areas of crops have been adopted about the crop report for the
1 2 8	Belgaum Bijapur Dharwar	Karnatak	6,339,540,807	1,724,204	3,677	10.07	year 1914-15
1 2 3 4	Thana Kolaba Ratnagiri Karwar	Konkan	1,343,127,918	1,380,282	973	2.7	
	Bombay City			58,808	ĺ		
	Total		21,059,510,678	0 225 817	2,256	6.1	

Statement showing the quantity of fodder from various crops available in various Divisions of Bombay Presidencey.

		Gujara	t Division.	ŀ	Deccan I	Division.
No.	Name of crop.	Area in acres.	Quantity of fod- der available.	Arc	ea in acres.	Quantity of fod- der available,
1	Jowar	583,286	955,648,512	K	432,397	5,28,281,035
		!		C	8,074,589	2,459,978,659
2	Bajri	375,158	270,113,760	ĸ	685,777	493,759,440
				C	3,479,517	2,072,609,096
8	Wheat	269,075	244,559,938	K	192,812	107,498,474
				C	534,654	228,885,577
4	Rice	336,337	475,590,608	K	46,367	60,542,783
1	l 			C	203,798	266,105,163
5	Gram	93,529	83,798,575	K	43,895	12,112,386
	. i			C	237,059	44,889,492
6	Groundnut	2,781	6,118,200	K	6,287	14,460,100
				C	183,158	439,579,200
7	Sugarcane	8,579	26,673,207	K	1,107	8,177,686
ì				C	82,805	258,497,495
8	Maize	115,658	133,064,529	K	8,474	2,613,317
1				C	22,511	12,108,667
9	Nagli, Ragi or Mar-	98,526	25,222,656	K	10,603	8,588,430
	Vå.				209,330	314,497,392
10	Other cereals (Wari, Sava, Kodra, Dengli Berati, etc.)	191,479	803,379,328		206,069	2,205,350,438
11	Tur	59,582	44,842,819		282,581	261,118,978
12	Other pulses (Kul- thi, Mug, Udid, Masur, Lang, Wal etc.)	121,814	69,682,762		879,884	518,638,825
13	Sesamum	78,780	24,649,414	K	53,594 100,539	19,293,840 27,266,177
14	Fodder crops (Jowars, Lucerne, guines grass).	64,891	824,455,000	C C	240 14,598	1,200,000 72,990,000
	Total	2,338,925	2,937,799,308		10,937,645	10,439,042,645

Note.—It was desired to work out figures for the two Khandesh Districts separately; but the acreage under pulses was not returned separately, hence the two Districts also are included in C. D. and the whole Division is called Deccan Division. Since the outturn of fodder per scre of different crops is different in Khandesh and Central Division the total quantity of fodder from important crops is given separately.

K-area in East and West Khandesh Districts.

C— area in Nasik, Ahmednagar, Poona, Sholapur and Satara Districts,

		Karnati	c Division.	Konkan I	division.
No.	Name of crop.	Area in acres.	Quantity of fod- der available.	Area in acres,	Quantitý of fod- der available.
1	Jawar	2,462,014	4,389,746,342	1,254	1,357,844
2	Bajri	578,644	344,675,085	23	15,130
8.	Wheat	525,861	278,443,400	214	82,893
4.	Rice	260,439	340,063,016	831,266	995,391,159
5	Gram	108,693	21,582,082	3,236	. 642,540
6	Groundnut	22,234	26,680,800	452	678,000
7	Sugarcane	12,923	96,780,347	3,855	23,947,414
8	Maize	13,716	11,233,404	498	358,187
9	Nagli, Ragi or Marva.	69,523	128,809,434	130,750	166,010,660
10	Other cereals (Wari, Sava, Kodra, Deng- li, Berati, etc.)	284,546	283,976,508	95,505	131,013,759
11	Tur	164,585	124,628,762	6,825	2,426,629
12	Other pulses (Kulthi, Mug, Udid, Masur, Lang, Waletc.).		141,705,383	39,910	18,997,160
13	Sesamum	22,883	6,461,244	4,683	1.646,543
14	Fodder crops (Jowars, Lucerne, guinea grass).		144,760,000	112	560,000
	Total	4,833,785	6,339,540,807	1,118,583	1,343,127,918

	1	Total	· ·	
No.	Name of crop.	Area in acres.	Quantity of fodder available.	Remarks.
1	Jawar	6,503,540	8,335,012,392	-
2	Bajri	5,119,119	8,181,172,511	
8	Wheat	1,522,616	859,470,282	
4	Rice	1,678,207	2,137,692,729	
5	Gram	486,412	113,025,075	
6	Groundnut	214,912	487,516,800	
7	Sugarcane	54,269	414,076,149	
8	Malze	155,857	159,378,104	
9	Nagli, Ragi or Marva.	518,732	643,128,572	
10	Other cereals (Wari, Sava, Kodra, Dengli Berati, etc.).	777,599	2,923,720,033	
11	Tur	513,573	433,012,183	
12	Other pulses (Kulthi, Mug, Udid, Masur, Lang, waletc.).	1,319,880	749,024,130	
18	Sesamum	255,429	79,317,218	
14	Fodder grops (Jowars, Lucerne, guinea grass).	108,793	548,965,000	Average yield of dry fodder taken to be 5,000 lbs. per acre.
	Total	19,228,938	21,059,510,678	

PART II.

CROPS OF EACH DISTRICT OF THE BOMBAY PRESIDENCY.


.

AHMEDABAD.

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreages.

A

No.	Taluka.	Cotton.	Jowari.	Bajri.	Wheat.	Rice.	Math.	Gram.	Castor.	Seasame.	Sugar- cane.	Other. crops.	Total.
		Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.
1	North Daskroi	18,237	. 23,757	18,563	1,472	5,366	2,964	213	3,377	419	. <u>18</u>	12,382	86,768
2-	South Daskroi	33,801	35,796	24,063	15,323	36,034	2,560	1,198	3,714	606	305 161	18,266	171,660
3	Dholka	89,148	41,235	. 7,252	65,496	15,299	1,321	8,205	1,908	2,540	$\frac{113}{46}$	24,487	201,994
4.	Dhanduka	71,534	97,598	45,709	44,359	11	2,134	8,040	6	8,627	$\frac{108}{442}$	13,976	287,097
5	Viramgaon	1,16,820	58,345	36,877	15,187	1,747	4,717	698	4,716	1,123	···· 55	12,259	251,989
6	Prantij	5,808	2,922	31,239	3,978	10,011	23,929	11,444	2,446	7,290	92 23	80,743	129,402
	Total	284,343	259,648	163,703	145,815	68,468	37,625	24,798	16,162	15,605	636 753	112,113	1,128,916




Statement showing the distribution of important crops by Talukas, the distribution is shown by percentages.

To.	Name of Taluks			Percentage of crops.										
	rome of Islue	••	Cotton.	Jowari.	Bajri.	Wheat.	Rice.	Math.	Gram.	Castor.	Seasame.	Sugar- cane.		
			%	%	%	%	%	%	%	%	%	% 2·83		
1	North Daskroi	- ''	6.41	9.15	11.84	1.009	7.84	7.87	0.86	20.89	2.68	8.46		
2	South Daskrol		11.88	18.78	14.70	10.21	52.63	6.80	4.83	22.98	8.88	47·95 21·88		
8	Dholka		18.77	15.88	4.42	44.92	22-34	8.21	12.02	11.78	16.28	17·77 6·10		
4	Dhanduka		25 15	187.59	27.92	80.42	0.02	5.67	82.42	0.03	28 · 24	16·98 58·70		
5	Viramgaon	••	40 91	22.47	22.52	10.41	2.55	12.53	2.81	29·18	7·19	7:30		
6	Prantij		1.87	1.12	19.08	2·78	14.62	68-60	46-14	15.13	46.71	14·46 8·06		
	١	Гоtạl	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100·00 100·00		

C

	,												
	Name of				<u>.</u>		Percent	ages.					
	Taluka.	Cotton.	Jowari.	Bajri.	Wheat.	Rice.	Math.	Gram.	Castor.	Seasame.	Sugar- cane.	Others.	Total.
		%	%	%	%	%	%	%	%	%	%	%	%
1	North Daskroi	21:02	27:38	21.39	1.69	6.18	3.41	0.24	8.89	0.48	0.02	14.30	100.00
2	South Daskroi	19 69	20.85	14.02	8.92	20.99	1.49	0.69	2.16	0.35	0.18	10.66	100.00
. 8	Dholka	19:38	20.41	8.28	32.42	7.57	0.65	1.59	0.94	1.26	0.05	12.14	100.00
4	Dhanduka	24.92	83.99	15.92	15.45	0.03	0.74	2.80	0.02	1.26	0.03	4.89	100.00
5	Viramgaon	46.16	23:15	14.63	6.03	0.69	1.87	0.27	1.87	0.44	0.00	4.90	100.00
6	Prantij	4.10	2.26	24·14	8.07	7 · 74	18-49	8.84	1.89	5.63	0.07	23.77	100.00
					!	<u> </u>	<u> </u>			<u> </u>			



75

KAIRA.

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreage.

A

No.	Taluka.		Bajri.	Kodra.	Rice.	Bavto or Nagali.	Cotton.	Tobacco.	Varialii.	Jiru.	Potatoes.	Sugarcane.	Others.	Total.
		ļ	A cres.	Acres.	Acres.	Acres.	Acres.	Acres.	A cres.	Acres.	Acres.	Acres.	Acres.	Acres.
1	Mehmedabad	••	15,017	8,452	13,º25	9,207	5,625	6,619	840	13	18	9	21,700	74,905
2	Matar		12,068	6,147	15,288	6,440	•7,735	1,877	21	23	1	20 25	81,015	80,135
8	Thasra		15,541	8,266	17,003	11,150	6,997	1,845	18	3	2		18,011	78,336
4	Kapadwanj		22,675	6,183	13,337	9,946	7,881	201	 196	3	7	2/2	39,772	100,203
5	Nediad		28,727	15,644	14,385	13,212	2,768	7,712	639	191	91	27 6	30,249	113,645
6	Anand		86,797	27,850	9,712	7,598	4,140	4,937	352	503	29	23 8	28,563	120,004
7	Borsad		22,091	21,502	5,672	4,104	3,190	9,846	2	181	3	42 37	25,409	91,992
	Tota	ı	152,916	93,544	89,222	61,657	88,336	26,117	1,568	-867	151	123	194,719	659,220

В

	Name	,				Percentag	e of crops.	•	·			
No.	of Taiuka.	Bajri.	Kodra.	Rice.	Bavto.	Cotton.	Tobacco.	Varialli.	Jiru.	Potatoes.	Sugarcane.	Total
		 %	%	%	%	%	%	%	%	%	%	
1	Mehmadabad	 9.81	9.04	15.49	14.93	14.67	2.68	21 68	1.50	11.92	7.31	
2	Matar	 7.89	6.22	17·13	10.44	20.18	5.27	1.34	2.65	0.66	16·26 28·73	
8	Thesra	 10.16	8.84	19.06	18.08	18.25	5.15	1.15	0.32	1.32	····	
4	Kapadwanj	 14.83	6.20	14.95	16.13	20.55	0.77	12.50	0.32	4.64	1·63 2·30	
5	Nadiad	 18:79	16.71	16.12	21.43	7.22	29.53	40.75	22.03	60.27	21·95 6·89	
6	Anand	 24.06	29.24	10.88	12.32	10.80	18-90	22.45	58.01	19.20	9.20	
7	Borsad	 14.45	22.99	6.86	6.66	8.32	37.70	0.13	15.11	1.98	34·14 42·53	
							<u> </u>					
	Total	 100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	

Statement showing how the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

O

	1		Percentages.											
No.	Name of Taluka.		Bajri,	Kodra.	Rice.	Bavto.	Cotton.	Tobacco.	Varialii.	Jira,	Potatoes.	Sugarcane.	Others.	Total.
			%	%	%	%	%	%	%	%	%	%	%	%
1	Mehmadabad		20.02	11.28	18-46	12.29	7 51	0.88	0.45	0.02	0.25	0.012	28.98	100.00
2	Matar		15 07	7.67	19.08	8.03	9.65	1.72	0.03	0.08	0.001	0.02	88.70	100.00
8	Thasra		19:84	10.55	21 · 70	14.23	8.93	1 72	0.02	0.004	0.003		23.01	100.00
4	Kapadwanj		22.63	6.17	18.31	9.92	7.86	0.20	0.18	0.003	0.007	0.002	39.71	100.00
5	Nadiad		25.28	13.77	12.66	11.63	2.44	6 · 78	0.56	0.17	0.08	0.02	26.61	100.00
6	Anand		80.66	22.79	8.09	6.33	3·45	4.11	0.29	0.42	0.02	0.02	23.82	100.00
7	Borsad .		24.01	23.37	6.16	4.46	3.47	10.70	0.002	0.14	0.003	0.04	27 · 65	100.00

7

PANCH MAHALS.

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreage.

A

No.	Taluka.	Maiz.	Rice.	Gram.	Bajri.	Seasame.	Cotton.	Bavto.	Kodra.	Ragi or Nagali.	Wheat.	Other crops.	Total.
1	Godhra	Acres. 26,366	Acres. 29,002	Acres. 13,400	Acres. 21,764	Acres. 17,009	Acres. 4,906	Acres. 11,675	Acres. 8,632	Acres. 9,368.	Acres. 1,476	Acres. 18,448	Acres. 162,046
2	Kalol and Halol	2,580	29,532	2,587	28,107	3,550	29,336	7,593	6,872	5,074	233	26,732	137,196
ង	Dohad and Zalod	82,082	10,567	37,991	105	13,908	175	772	4,068	1,676	12,251	32,980	196,575
	Total	111,028	69,101	59,978	44,976	34,467	34,417	20,040	19,572	16,118	13,960	78,160	495,817

Statement showing the distribution of important crops by Talukas, the distribution is shown by percentages.

В

	·											
			Percentage of crops.									
No.	Name of Taluka.	Maiz.	Rice.	Gram.	Bajri.	Seasame.	Cotton.	Bavto.	Kodra.	Ragi.	Wheat.	· Total.
1	Godhra	% 23·74	% 41·97	% 24·82	% 48·39	% 49·34	% 14·25	% 58·26	% 44·10	% 58·12	% 10·57	
2	Kalol and Halol .	2.82	42.73	4.79	51.37	10.80	85.23	37.89	35·11	31-48	1.67	
3	Dohad and Zalod .	73.93	15.29	70.38	0.53	40.35	0.21	8.85	20.78	10.39	87.75	
	Total .	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	

	FANCH-MARAL.	Fig.28
MAIZE	RICE	GRAM
BAJRI S DOO ACRES.		
BAJRI	SESAMUM	COTTON
ONE DOT REPRE		
KODRA	RAGI	WHEAT
ONE DOT METALES LATS 100 ACRES.		
Names of Salahas MAIX 6	10 5 1N 7 M	PUSANDS
I GODNRA 2 MALGL 380HADQVIALOO		
I GODHRA 2 MALOLEHALOL 3 DOMADEJMALOB	SCANON.	7.76
J GODHRA 2 MALOLBHAEDL	ECTTON	

Statement showing how the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

(


No.	Name of Taluka.	Percentages.												
		Maiz.	Rice.	Gram.	Bajrl.	Seasame,	Cotton.	Bavto.	Kodra.	Ragi.	Wheat.	Others.	Total.	
		%	%	%	%	%	%	%	%	%	%	%	%	
1	Godhra	16.27	17.90	8-27	13.43	10.49	3.05	7 - 20	5.82	5 · 78	0.91	11-41	100.00	
2	Kalol and Halol	1.81	21.25	1.88	16.84	2.58	21 · 38	5.58	5.00	3.69	0.17	19.58	100.00	
8	Dohad and Zalod	41.75	5.87	19.82	0.05	7.07	0.08	0.89	2.07	0.85	6 · 23	16.81	100	

BROACH.

The following statement shows the distribution of important crops by Talukas. Figures in the following statement show actual acreage.

A

No.	Tajuka.		Cotton.	Jowari	Wheat.	Lang	Rice.	Tur.	Seasame.	Bajri.	Tobacco.	Sugarcane.	Others.	Total.
1	Broach		Acres. 50,142	Acres. 24,968	Acres. 3,317	Acres. 13,281	Acres. 5,821	Acres. 2,496	Acres. 2,263	Acres.	Acres 1,077	Acres.	Acres. 3,341	Acres. 107,261
2	Amod		46,210	17,292	3,115	10,318	686	886	771	591	124		1,254	81,242
. 3	Wagra	• -	84,010	33,249	26,911	12,975	76	112	861	574	169		2,147	110,884
4	Jambusar		57,114	25,760	25,222	728	4,883	4,867	3,047	4,943	258	₁	13,195	139,517
5	Ankleswar		50,356	39,095	11,108	2,330	2, 313	3,888	2,840	1,267	675	12 38	4,539	118,413
	Total		2,37,832	1,40,864	69,678	39,627	13,279	. 12,249	9,782	7,915	2,303	17 50	24,276	557,817



Statement showing the distribution of important crops by Talukas, the distribution is shown by percentages.


						В								
		· · · · · · · · · · · · · · · · · · ·		Percentage of crops.										
No.		Name of Taluka.		Cotton.	Jowari.	Wheat.	Lang.	Rice.	Tur.	Seasame,	Bajri.	Tobacco.	Sugarcane.	
				%	%	%	%	%	%	%	%	%	%	
1	Broach		••	21.08	17.78	4.76	83.21	48.88	20.38	28·18	6.82	46.76	29·41 22·0	
2	Amod			19.42	12-32	4.47	26.02	5.16	7.23	7.88	7-47	5.88		
3	Wagra	••	••	14.80	23 · 68	88-62	32.74	0.57	0.91	8 80	7-25	7.84		
4	Jambusar	••		24.01	18.35	86.50	1.84	88-00	89.73	81.15	62-45	11.20	2.0	
5	Ankleswar	••		21.17	27.85	15.94	5.88	17.42	31.74	29.03	16.00	29.81	70.59 76.0	
			Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	

α

Statement showing how the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

					•	C		•	•	•			
	•					Per	centage.		··				
No.	Name of Taluka.	Cotton.	Jowar.	Wheat.	Lang.	Rice.	Tur.	Seasame.	Bajri.	Tobacco.	Sugarcane.	Other.	Total.
1	Broach	% 46·75	% 23·28	% 8·09	% 12·38	% 5·42	% 2.32	% 2·11	% 0·50	% 1·004	% 0·005	% 8·14	% 100·00
2	Amod .	56.88	21.28	8.83	12.69	0.84	1.09	0.95	0.78	0.12	0.00	1.56	100.00
3	Wagra .	80.67	29.99	24.27	11.70	0.07	0.10	0.77	0.21	0.15	0.00	1.77	100.00
4	Jambusar .	. 40.93	18.46	18.08	0.52	8.14	8.49	2·18	8.54	0.18	0.00	11.48	1 0.00
5	Ankleswar .	. 42.52	80.10	9.88	1.97	1.95	8 · 28	2.89	1.07	•57	0.01	8.76	100.00

.



SURAT.

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreages.

A

No.	Taluka.	Cotton.	Jowari.	Rice.	Wal.	Wheat.	Castor.	Kodra.	Tur.	Nagali.	Sugarcane.	Others.	Total.
-		Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.
1	Chorashi	10,581	15,295	927	924	554	363	1	1,781	••••	10 7	5,635	35,871
2	Olpad	29,313	30,997	3,270	1,136	28,029	964	1	2,508		1	9,988	100,987
3	Mandvai	19,456	19,222	11,961	5,237	1,591	4,478	4,942	4,048	1,299	1 1	8,537	80,767
4	Barodli	37,391	19,507	18,548	10,482	2,411	52,519	586	2,710	660	85	8,052	105,691
5	Jalalpoor	10,864	11,651	11,109	5,785	76	2,554	38	1,119	203	381 1,344	4,570	54,350
6	Chikhali	9,104	4,346	17,945	7,015	59	1,051	2,118	1,274	1,481	555 689	4,362	49,310
7	Bulsar	267	457	19,229	- 7,449	9	3,027	5,073	1,194	3,273	941 794	8,847	44,266
8	Pardi		. 64	13,478	5,242	500	2,459	6,864	1,551	9,265	677	8,136	43,308
	Total	122,976	1,01,539	96,267	43,270	28,229	19,885	19,623	16,185	16,181	2,723 3,833	47,622	514,500

T

			·										
							Percentage	of crops.	·				_
No.	Name of Taluka.	Ì	Cotton.	Jowar.	Rice.	Wal.	Wheat.	Castor.	Kodra.	Tur.	Nagli.	Sugarcane.	
			%	%	%	%	%	%	%	%	. %	%	
1	Chorashi	•	8.60	15.06	0.75	2.13	1.96	1.82		11.00	0.00	0.38	
2	Olpad	••	23.83	30.23	3.40	2.62	81.58	8 · 49		15.49	0.00	0.03	
3	Mandavi		15.82	18.93	12.42	12.10	5.63	22.52	25 · 18	25.01	8.03	0.03	
4	Bardoli		30.40	19.21	19:27	24.22	8.54	26 · 45	2.99	16.74	4.08	3.16	
5	Jalaipur	••	13.71	11.47	11.54	18:37	0.27	12.84	0.19	6.91	1.25	35·06 13·99	
6	Chikhali .	•	7.40	4.28	18-64	16.21	0.20	5 · 28	10.79	7 . 87	9.15	20.38	
7	Bulsar	••	0.51	0.45	19.97	17.22	0.03	15.22	25 85	7:38	20 - 23	34·56 25·93	
8	Pardi	•	••••	0.08	14.00	12.12	1.77	12.87	84.98	9.58	57.26	27·53 17·66	
	Total		100.00	100 00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	-


								Percentages	·				-		
No.	Name of Taluk	ca.	Cotton.	Jowar.	Rice.	Wal.	Wheat.	Castor.	Kodra.	Tur.	Nagal i.	Sugarcane.	Others.	Total.	
			%	%	%	%	%	%	%	%	%	%	%	%	
1	Chorashi		29.49	42.64	2.03	2.58	1.54	1.01	0.00	4.96	• • • •	0.003	15.75	100.00	
2	Olpad		29.04	80.71	8.24	1.12	22·81	0.69		2.48	••••	••••	9.91	100.00	
8	Mandavi		24.09	23.80	14.81	6.48	1.97	5.54	6.12	5.01	1.61		10.57	100.00	
4	Bardoli		85.88	18:45	17.55	9-92	2.28	4.98	0.55	2.56	0.62	0.08	7·63	100.00	
5	Jalalpur		81.08	21.44	20.44	10.64	0.14	4.69	0.07	2.06	0.37	0.07	9.05	100.00	
6	Chikhali		18-46	8.81	86.39	14.28	0.12	2·18	4.29	2.58	8.00	1.18	8.86	.100*00	
7	Bulsar		0.60	1.03	48-44	16.83	0.03	6.84	11-46	2.70	7.39	2.13	7·46	100.00	
8	Pardi		••••	0.12	81.12	12·10	1.15	5.68	15.85	8 58	21.39	1.78	7 · 25	100.00	

WEST KHANDESH.

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreages.

A

No.	Name of Taluka.		Cotton.	Bajri.	Wheat.	Jowar.	Kulith.	Seasame.	Gram.	Math.	Ground- nut.	Sugarcane.	Others.	Total.
	•		Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.
1	Dhulia	$\cdot \cdot $	121,098	1,01,860	8,054	19,586	17,817	6,980	491	2,812	950	52 241	5,192	284,342
2	Sindkheda		112,935	71,954	16,599	15,196	11,067	3,018	1,359	5,469	361	13 26	5,692	243,663
3	Shirpur		64,300	30,362	7,287	8,194	4,652	8,608	1,436	1,821	17	1 10	3,468	125,146
4	Shahada		64,122	32,663	86,733	22,148	4,808	6,958	7,167	2,293	60	55 35	7,894	184,901
5	Taloda	••	7,691	2,760	24,298	10,624	76	5,672	5,739	169	9	3 5	15,249	72,290
6	Nandurbar	•	60,895	38,321	44,488	26,417	7,203	8,327	11,994	1,817	864	28 55	93,253	292,107
7	Sakri		47,658	81,278	7,194	8,664	20,824	8,294	4,753	832	. 802	601 945	36,942	212,342
	Total		478,199	859,198	144,653	110,229	65,947	87,807	82,939	14,718	2,063	753 1,317	167,690	1,414,791



C Percentages. Name of Taluka. Total. No. Cotton. Bajri. Wheat. Jowar. Kulith. Sesame. Gram. Math. Ground-Sugarcane. Others. % % % % % % % Dhulia 6.89 2.43 0.17 0.99 0.33 1.89 100.00 42.55 35.82 2.83 0.018 100.00 Sindkheda 46.34 29.53 6.81 6.23 4.54 1.24 . 0.55 2.24 0.14 0.005 1.14 1.45 0.013 0.00 2.81 100.00 Shirpur 51.38 24.26 5.82 6.24 3.71 2.88 4.31 2.60 3.87 1.24 0.03 0.03 100.00 Shahada 84.67 17.66 19.86 11.97 3.76 Taloda 10:64 8.81 33.61 14 69 0.10 7.84 7.94 0.23 0.01 0.00 21.13 100.00 4.10 0.45 0.12 81.95 100.00 Nandurbar 20.67 13.12 15.23 9.04 2.46 2.85 0.01 100.00 3.38 4.08 9.80 1.55 2.24 0.39 0.14 0.27 17.44 Sakri 22.44 38:27

COTTON	BAJRI	JOWAR
WTS 1000 ACRES.	ior	J. S.
OION ESENTS	KULTHI	WHEAT
SOS	100	ier
SESAMUM	GROUND-NUT	SUGAR-CANE
Be a serve to a ches	ior	in o
Names of Talustys CPT TON		365 (A)
I MALGOON 2 SENDER SENDER 5 ANALOR SENDER 5 ANALOR SENDER 5 C PRANCO 7 CACPER 8 YAVAL 10 CM & USSAON		
IVALGAON ZAMMER A JAPUSAMAL SAMAL MER SAMAL MER SE ANNOL T CHOPON B VAVAL B CORP. JOSAGORA JO		S FOUNT (VI
I VAL CAON 2 JAMNER 3 BHUSANAN 4 RAVER 5 EMANUOL 7 CHOPPA 8 XHVAL 2 PACHOPA OCHALISGAON	WASAT .	SORANGANE

EAST KHANDESH.

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreages.

							A							
No.	Taluka.		Cotton.	Bajri.	Jowar.	Vdid.	Kulith.	Wheat.	Seasame.	Gram.	Groundnut.	Sugarcane.	Other crops.	Total.
			Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.
1	Jalgaon	•	66,210	14,368	23,360	12,975	2,163	10,219	568	2,114	188	28 81	10,598	142,786
2	Jamner		112,787	20,505	46,100	24,507	17,076	186	1,404	154	70	20 48	15,078	237,887
8	Bhusawal	••	112,142	18,456	59,866	21,256	4,946	2,117	1,741	378	871	12 · 46	16,926	238,206
4	Raver	•••	65,299	7,550	24,504	7,082	346	5,071	1,606	503	796	5 56	15,165	127,927
Б	Amainer		136,947	51,766	30,221	4,416	16,495	6,514	2,680	2,436	671	53 845	11,440	263,589
6	Erandol	••	99,890	42,281	38,751	14,095	9,924	8,807	1,363	1,659	178	60 117	16,088	232,546
7	Chopda	••	84,189	22,076	19,866	9,008	1,188	4,752	1,823	908	. 10	9 15	8,142	151,471
8	Yaval	••	68,560	14,499	19,253	13,958	540	8,699	746	864	1,646	€ 22	8,150	181,421
9	Pachora	••	112,672	57,586	32,905	15,756	10,850	8,267	1,752	878	183	84 807	13,434	249,347
. 10	Chalisgaon		80,988	77,542	26,742	1,298	12,622	8,527	2,654	1,567	111	102 231	11,480	218,528
	Total		939,129	326,579	321,568	124,851	76,150	48,159	15,787	10,956	4,224	354 1,218	126,451	993,708

89

Statement showing the distribution of important crops by Talukas, the distribution is shown by percentages.

В

	Name	i				P	ercentage of	crops.				
To.	of Taluka.		Cotton.	Bajri.	Jowar.	Udid.	Kulith.	Wheat.	Seasame.	Gram.	groundnut.	Sugarcane
			%	%	%	%	%	%	%	%	%	%
1	Jalgaon	••	7.05	4.39	7.26	10.41	2.84	21.22	3.29	19.29	4.45	2.54
2	Jamner		12.00	6.27	14.83	19.70	22.42	0.38	8.79	1.40	1.66	3·65 3·94
3	Bhusawal		11.94	5.65	18-61	17:09	6.49	4.39	11.02	3.40	8.78	34·39 3·78
4	Raver		6.94	2.31	7.62	5.69	0.45	10.23	10.17	4.59	18.84	1·41 4·60
5	Amalner	•	14.58	15.79	9.39	3.55	21.66	13.52	16.68	22.22	15 88	14·97 28·32
6	Erandol	· ••	10.58	12.93	12.05	11.33	13.03	18.28	8.63	15.14	4.21	16.95 9.60
7	Chopda	••	8.97	6.76	6·17	7 · 24	1.26	9.87	8.38	8.28	0.53	2.54 1.23
8	Yaval		7:30	4 43	5.98	11.22	0.70	7.68	4.72	3.32	38.96	1.69
9	Pachora ·		11.99	17.63	10.23	12.67	14.24	6.78	11.09	8.01	4.33	18·07 25·20
10	Chalisgaon	••	8.62	23 · 74	8.31	1.04	16.57	7:32	16.81	14.30	2.63	28·80 18·96
		Total	100.00	100.00	100.00	100.00	100.00	100.00	100 00	100.00	100.00	100.00

C

Statement showing how the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

.	Name							Percenta	ges.					I
No.	of Taluka,	_	Cotton.	Bajri.	Jowar.	Udid.	Kulith.	Wheat.	Seasame.	Gram.	Groundnut.	Sugarcane.	Others.	Total.
			%	%	%	%	%	%	%	%	%	%	%	%
1	Jalgaon	••	46.37	10.08	16.36	9.08	1.21	7.15	0.39	1.48	0.18	0.01	7.46	100.00
2	Jamner		47.42	8·19	19·87	10.30	7.17	0.07	0.20	0.08	0.08	0.01	6 79	100.00
8	Bhusawal	••	47.08	7.74	25·18	8.92	2.07	0.89	0.78	0.12	0.12	0.002	7.14	100.00
4	Raver		51.04	5.90	19·15	5.23	0.27	3.96	1.25	0.39	0.62		11.89	100.00
5	Amainer		51.94	19.64	11.46	1.67	6.25	2.47	0.88	0.92	0.25	0.02	4.39	100.00
6	Erandol	••	42.74	18·16	16.66	6.06	4.26	8.78	0.28	0.71	0.07	0.02	6.96	100.00
7	Chopda	٠.	55.58	14.57	13.11	5.94	0.78	8·18	0.87	0.60	0.08	00.05	5.41	100.00
8	Yaval		52·16	11.08	14.65	10.47	0.41	2.81	0 56	0.27	1.25	00.04	6.39	100.00
9	Pachora		45.18	23.09	13·19	6.82	4.85	1.31	0.70	0.32	0.04	0.05	5.45	100.00
10	Chalisgaon		37.03	85.48	12.23	0.59	5:77	1.61	1.21	0.71	0.05	0.04	5.28	100.00


မ

NASIK.

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreages.

Α


							A							
No.	Taluka.		Bajri.	Wheat.	Cotton.	Jowar.	Gram.	Rice.	Ground- nut.	Sugarcane.	Onions.	Methi.	Others.	Total.
1	Nasik		Acres. 27,654	Acres. 45,873	Ac es.	Acres. 1,023	Acres. 9,178	Acres. 8,093	Acres. 2,097	Acres. 877 1,241	Acres. 796	Acres. 501	Acres. 51,831	Acres. 147,433
2	Sinnar		137,196	16,874	111	2,436	3,563	1,714	3,528	809 1,209	552	198	49,645	215,626
8	Igatpuri		847	16,538	••••	808	8,834	25,459	334	27	6	4	64,462	117,301
4	Dindori	.:	11,223	61,471		782	17,538	7,736	1,935	946 1,411	187	240	80,639	182,697
5	Niphad		92,708	61,069	108	2,146	6,684	186	3,824	664 1,960	1,431	484	83,089	202,391
6	Chandwad		98,124	17,996	529	8,308	8,644	182	2,747	288 1,097	458	161	29,128	156,560
7	Yeola		100,481	22,162	1,234	21,112	2,058	2	. 903	118 570	156	35	29,960	178,221
8	Nandgaon		81,139	6,241	18,489	5,683	586	9	85	8 18	88	12	18,445	129,685
9	Malegoan		159,339	6,828	61,245	18,275	2,889	684	1,129	895 590	84	129	46,400	296,892
10	Bagian	••	98,796	6,201	25,383	11,952	4,824	798	692	$\frac{727}{1,147}$	282	56	43,215	192,426
11	Kalwan		50,009	8,799	3,782	9,492	9,065	5,786	603	_580 829	852	104	55,532	144,104
12	Peint			22	••••	24	837	15,214	806	4			68,758	84,661
	Total		856,514	260,069	110,881	77,036	68,700	65,863	18,133	5,431 10,041	4,342	1,924	570,104	2,047,997



E

_						Percen	tage of crop	18.					
No.	Name of Taluk	B.	Bajri.	Wheat.	Cotton.	Jowar.	Gram.	Rice.	Ground- nut.	Sugarcane.	Onions.	Methl.	
Ì		•	%	%	%	%	%	%	%	%	%	%	
1	Nasik.	••	8.28	16.86	••••	1.83	18.86	12.29	11.56	16·38 . 12·85	18-88	26-04	
2	Sinnar	••	16.02	6.27	0.10	. 8 · 16	5·19	2.60	19:46	14·40 12·04	12.71	10.29	
. 8	Igatpuri	••	0.10	6.12	0.0	1.05	12.86	88-65	1.84	0·17 0·26	0.14	0.51	
4	Dindorl		1.81	22.84	0.0	1.02	25.58	11.75	10.67	17·42 14·05	4.81	12.47	
5	Niphad		10.82	22.70	0.10	2.79	9.78	0.58	21.09	12·28 19·51	82.96	25·16	'
6	Chandwad	••	11-46	6.69	0.48	4.29	5.30	0.27	15.15	5·30 10·92	10.55	8:37	
7	Yeola		11.73	8-24	1.11	27:40	2.99	0.08	4.98	2·17 5·07	8.59	1.82	
8	Nandgaon		9.36	2.82	16.67	7.38	0.85	0.01	0.10	0·15 0·15	0.88	0.62	
9	Malegaon	••	18.60	2.85	55.23	28.72	4-20	1.08	6.53	7·27 5·87	1.93	6.70	
10	Bagian		11.28	2.30	22.89	15.20	6-29	1.51	3.81	13·89 11·42	6.49	2.91	
11	Kalwan		5 88	8 · 27	8.41	12.32	18.20	8.78	8.88	10·66 8·25	8-10	5.40	
. 12	Peint			0.08	0.0	0.08	0.49	23·10	1.68	0.03		••••	
		Total	100-00	100.00	100.0	100.00	100.00	100.00	100.00	100.00	100.00	100.00	

	I						Percen	tages.						
No.	Tajuka.		Bajri.	Wheat.	Cotton.	Jowar.	Gram.	Rice.	Ground- nut.	Sugarcane.	Onions.	Methi.	Others.	Total.
			%	-%	%	%	%	%	%	%	%	%	%	%
1	Nasik	ا.:	18.75	30.70		0.69	6.23	5.49	1.42	0.60	0.54	0.34	85 17	100.00
2	Sinnar	İ	63.63	7.83	0.02	1.18	1.65	0.79	1.68	0.87	0.26	0.09	22.57	100.00
8	Igatpuri		0.72	14.09		0.69	7.52	21.71	0.28	0.007	0.002	0.008	54.99	100.00
4	Dindori		6.14	138-64		0.43	9.59	4 · 23	1.05	0.21	0.011	0.18	44.17	100.00
5	Niphad		45.81	80.17	0.08	1.06	8.30	0.09	1.89	0.83	0.70	0.24	16.35	100.00
6	Chandwad		62 - 67	11.49	0.84	2.10	2.32	0.12	1.76	0.18	0.29	0.11	18.62	100.00
7	Yeola	[56.88	12.44	0.69	11.85	1.15	0.01	0.50	0.08	0.09	0.02	16.82	100.00
8	Nandgaon		61.79	4.81	14.26	4.88	6 · 45	0.07	0.027	0.006	0.08	0.008	14.24	100.00
9	Malegaon		58.67	. 2.13	20 63	6.16	0.97	0.28	0.38	0.13	0.03	0.04	15.63	100.00
10	Baglan		51.84	8.22	18:19	6.21	2.25	0.40	0.36	0.38	0.15	0.8	22.47	100.00
11	Kalwan		84.70	6-10	2.62	6.59	8-29	4.02	0.42	0.40	0.24	0.07	88.55	100.00
12	Peint		,	0.02	,	0.02	0.40	17:97	0.36				81 · 23	100.00



AHMEDNAGAR.

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreage.

A

No.	Name of Taluka.	Bajri.	Jowar.	Cotton.	Wheat.	Saflower.	Sesame.	Gram.	Linseed.	Groundnut.	Sugarcane.	Others.	Total area
	Nagar	Acres. 72,509	Acres. 122,778	Acres. 15,361	Acres. 7,255	Acres. 16,224	Acres. 5,948	Acres. 1,860	Acres. 295	Acres. 677	Acres. 82 43	Acres. 82,084	Acres. 274,518
. 2	Parner	1,47,217	76,178	1,310	17,798	13,935	951	2,018	267	434	136 882	63,834	829,508
3	Shrigonda	54,955	121,865	7,862	3,783	20,702	. 2,410	2,247	700	821	42 225	42,228	256,610
4	Karjat	44,434	78,566	17,558	1,862	11,414	4,045	1,588	837	411	14 72	28,582	189,301
5	Jamkhed	44,500	49,675	7,638	2,974	5,055	1,029	3,135	823	1,265	185	20,347	136,450
6	Shevgaon	60,871	55,862	83,948	8,528	8,164	8,944	6,155	1,307	7	178	18,184	241,425
7	Pathardi	88,886	22,308	30,772	1,386	1,584	3,581	1,490	549	91	67 278	19,183	169,899
8	Newasa	80,506	77,085	77,676	14,060	9,309	9,465	7,978	1,085	92	51 176	20,131	297,438
9	Rahuri	90,197	53,284	22,676	12,202	5,320	4,983	3,565	453	539	453 803	19,848	218,820
10	Kopergaon	138,168	81,928	6,035	50,486	4,066	6,063	4,817	1,026	1,651	2,377 6,214	81,085	277,652
11	Sangamner	188,293	16,761	1,911	3,582	667	9,138	1,136	55	552	593 852	46,878	269,566
12	Akola	58,776	6,907		5,418	485	221	5,765	1	290 •	298 416	86,857	165,016
	Total	1,068,812	712,687	272,742	124,324	91,925	56,773	41,244	7,398	6,330	4,079 10,394	428,436	2,814,78

Statement showing the distribution of important crops by Talukas, the distribution is shown by percentages.

B

			·	~									. —
_	Name of Taluka.	ł					Percent	tage of crops	•				
No.	Taluka.		Bajri,	Jowar.	Cotton.	Wheat.	Safflower.	Sesamum.	Gram.	Linseed.	Groundnut	Sugarcan .	Ī
1	Nagar		% 6·78	% 17·28	% 5·63	% 5·84	% 17·65	% 10·47	% 8·30	% 3·99	% 10·70	% 0.78 0.41	-
2	Parner		13.77	10.69	0.48	14.81	15.16	1.67	4.89	3.61	6.85	3·33 8·54	
3	Shrigonda		5.14	17.08	2.88	8.04	22.52	4.24	5.45	9.46	5.07	1.02	
4	Karjat	• •	4-16	11.02	6.43	1.20	12.42	7.12	8.84	11.81	6.49	0.84	
5	Jamkhed	••	4.16	6.97	2.80	2.39	5 50	1.81	7-60	11.12	19.98	0.22	
6	Shevgaon		5.65	7-82	80.78	2.48	8 44	15.75	14.92	17.66	0.11	0.17	
7	Pathardi		8.82	8 · 13	11.28	1.11	1.72	6.30	8.61	7.42	1.44	2.69	
8	Newasa	••	7.58	10.82	28.48	11.80	10.12	16.67	19.33	14.67	1.45	1·25 1·70	
9	Rahuri	••	8.43	7.48	8.81	9 81	5.70	8.76	8.64	6.12	8.21	11·15 7·78	
10	Kopergaon	••	12.93	4.48	2.21	40.80	4.42	10.68	11.68	13.87	26.08	58·27 60·18	
11	Sangamner	••	17.61	2.85	0.70	2 88	0.72	16.09	2.75	0.74	8.72	14·53 8·25	
12	Akola	••	5.20	0.97	••••	4.86	0.23	0.88	13.98	0.00	4.28	7·30 4·03	
		Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	

Statement showing how the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

\ \	27 4 M-1-1-						Perce	ntages.						!
Mo.	Name of Taluk	A.	Bajri.	Jowar.	Cotton.	Wheat.	Safflower.	Sesamum.	Gram.	Linseed.	Ground- nut.	Sugarcane.	Others.	l'otal.
_			%	\ %	%	%	%	%	%	%	%	%	%	%
1	Nagar		26.41	44.72	5.28	2.64	5.91	2.16	0.49	0.10	0.24	0.01	11.7	100.00
2	Parner		45-49	28.54	0.40	5.20	4.80	0.29	0.62	60.08	0.13	0.01	19.6	100.00
3	Shrigonda		21 · 41	47.29	8.06	1.47	8.07	0.94	0.87	0.27	0.12	0.01	16.3	100
4	Karjat		23 · 47	41.50	9.27	0.98	6.08	2.14	0.43	0.44	0.22	0.007	15.1	100
5	Jamkhed		32.61	86-40	5.60	2.19	3.70	0.75	2-30	0.60	0.92	0.008	14.9	100
6	Shevgaon		24.99	23.15	84.77	1.46	1.81	8.70	2.54	0.24	0.003	0.003	7.5	100
7	Pathardi		52-82	13.12	18.11	0.81	0.93	2.11	0.88	0.32	0.05	0.04	11.8	100
8	Nevasa		27.06	25 · 91	26.11	4.72	8.13	3.18	2.68	0.86	0.03	0.01	6.8	100
9	Rahuri		42.28	24.97	10.63	5.72	2.49	2.83	1.67	0.21	0.25	0.21	9.2	100
10	Kopergaon		49.78	4.46	2-17	18-18	1.46	2.18	1.73	0.37	0.28	0.85	11.2	100
11	Sangamner		69.85	6.21	0.40	1.82	0.24	3.39	0.42	0.02	0.20	0.22	17.4	100 .
12	Akola		35.61	4.18		3-28	0.29	0.18	3.49		0.17	0.18	52-6	109

POONA.


The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreage.

A

No.	Name of Taluka		Bajri.	Jowar.	Rice.	Wheat.	Gram.	Cotton.	Ground- nut.	Sugarcane.	Potatoes.	Sesamum.	Others.	Total.
	<u> </u>		Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.
1	Junnar	••	119,664	- 83,580	12,687	23,488	10,259	,	2,236	712	269	81	44,047	247,028
2	Indapur	••	56,081	124,428	450	1,920	3,499	16,115	3,440	1,027 1,954		716	84,328	242,014
8	Khed		113,908	36,669	15,286	16,078	18,161	8	9,416	450 765	5,133	215	76,854	287,178
4	Sirur	••	134,018	88,464	13	6,142	2,011	8,342	985	159 685	617	1,486	49,713	286,945
Б	Purandhar		88,990	64,588	2,478	2,992	4,025	1	421	231 1,098	222	238	84,658	193,839
6	Bhimthadi		145,927	198,194	1,285	4,794	4,641	6,244	1,025	5,941 6,548	6	1,844	56,928	426,824
7	Haveli		78,179	75,658	20,191	7,550	4,469	7	1,087	_4,620 _4,781	44	163	41,584	228,502
8	Mawal		2,173	10,876	21,892	10,722	2,496	3	126	_56 78	54		22,539	70,987
	Total		728,985	632,462	74,282	78,686	44,561	25,720	18,736	13,196 17,207	6,345	4,748	860 596	1,983,262

The second figure given in each case under sugarcane represents the condition in 1917-18.

9




1

_						Pe	rcentage of	crops.					
No.	Name of Taluka.		Bajri.	Jowar.	Rice.	Wheat.	Gram.	Cotton.	Groundnut.	Sugarcane.	Potatoes.	Seasame.	
		-	%		%	%	%	%	%	% 5·89	%	%	
1	Junnar	}	16.42	5.80	17.08	81.87	23 · 02		11.93	7.54	4.24	1.70	
2	Indapur		7*69	19.67	0.60	2.60	7.85	62-65	18.36	7·78 11·35		15.09	
8	Khed		15.62	5 - 79	20.28	21.82	29.58	0.08	50.25	3·41 4·44	80.89	4.58	
4	Sirur		18.88	13.99	0.02	8.88	4.51	12.99	5 25	1·20 3·98	-9.72	81.88	هر ۱
. 5	Purandhar		11.52	10.21	3.33	4.08	9.08	00.04	2.24	1·75 6·38	8-49	5.02	
6	Bhimthadi		20.02	31.85	1.78	6.50	10.41	24.27	5.47	45·02 88·05	0.08	38.88	
7	[[aveli		10.04	11.98	27·18	10.54	10.03	0.027	5.80	35·01 27·78	0.69	3.43	
8	Maval		0.29	1.72	29 · 47	14.55	5.59	0.011	0.67	0:42 0:45	0.85		•
		Cotal	100.00	100.00	100.00	100.00	100-00	100.00	100.00	100-00	100.00	100:00	

C

	Name	-					F	ercentage.						
No.	of Taluka.		Bajri	Jowar,	Rice.	Wheat.	Gram.	Cotton.	Ground nuts.	Sugarcane.	Potatoes.	Sesamum.	Others.	Total.
1	Junnar		% 48·44	% 13·59	% 5·13	% 9·51	% 4·15	%	% 0·90	% 0·29	% 0·10	% 0·03	% 17·86	% 100·00
2	Indapur		23 · 17	51.42	0.18	0.79	1.44	6.66	1.42	0.42	Nil	0.59	14-21	100.00
3	Khed		89.66	12.77	5.82	5.28	4.28	0.008	3.58	0.16	1.79	0.07	26.78	100.00
4	Sirur		46.70	80.83	0.004	2.14	0.40	1.16	0.34	0.02	0.22	0.20	17:36	100.00
5	Purandhar .		43.88	83.82	1.28	1.24	2.07		0.51	0.12	0.11	0.11	17:91	100.00
6	Bhimthadi		34.19	46-43	0.80	1.12	1.08	1.46	0.24	1.39	0.001	0.48	13.36	100.00
7	Haveli	•	32.02	83·11	8.84	8.80	1.86	0.003	0.48	2.02	0.05	0.07	18˙-18	100.00
8	Mawal		8.06	15.83	80.86	15.11	8.52	0.004	0.18	0.08	0.07		81.79	100.00

100



10

SHOLAPUR.

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreage.

A

₹o.	Name of Taluk	3.	Jowar.	Bajri.	Cotton.	Safflower.	Tur.	Groundnut.	Math.	Hemp.	Wheat.	Sugarcane.	Others,	Total.
		_	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.
1	Sholapur		298,600	42,201	9,417	13,057	31,017	7,446	1,643	8,006	8,815	734	20,554	486,175
2	Barsi		159,472	13,617	7,509	1,692	17,227	34,529	294	5,329	5,296	236 616	81,081	276,282
8	Madha		144,492	60,807	25,884	13,331	18,654	4,437	5,878	9,001	2,083	695 502	19,852	800,124
4	Karmala	••	175,654	53,281	52,710	21,806	7,212	660	2,278	4,652	2,600	22 115	25,155	845,530
5	Pandharpur		135,823	48,652	11,792	10,668	1,778	639	4,469	1,089	2,918	48 518	12,676	230,547
6	Sangola		85,795	117,651	2,493	10,518	1,815	834	14,773	1,960	4,155	143 820	21,167	261,804
7	Malsiras	••	128,785	67,307	8,085	15,664	1,086	199	5,803	299	1,221	260	6,912	235,394
	Total		1,128,601	403,516	117,840	86,286	73,784	48,744	35,218	30,336	22,088	1,566 3,065	187,407	2,085,856

Statement showing the distribution of important crops by Talukas, the distribution is shown by percentages.

В

	 					,				r			
No.	Name of Taiu	ka.	Jowar.	Bajri.	Cotton.	Safflower.	Tur.	Groundnut.	Math.	Hemp.	Wheat.	Sugarcane.	
			%	%	%	%	%	%	%	%	%	%	
1	Sholapur		26-46	10.46	7.99	15.14	42.04	15.28	4.66	. 26.89	17.27	23.98	
2	Barsi	••	14·13	3.37	6.87	1.96	23.35	70.84	0.83	17:57	23-97	15·07 20·09	
3	Madha		12.80	15.07	21.97	. 15.45	18.51	9.10	16.69	29.67	9-43	16·37	
4	Karmala	••	15.26	13.20	44.73	24.70	9.76	1.35	6.47	15.33	11.77	3·75	
5	Pandharpur		12.03	12.06	10.00	12.37	2.40	1.31	12.69	3.28	13.21	3.06	
6	Sangola		7:60	29.16	2.11	12.19	2.46	1.71	41 95	6.46	18.81	9·18 10·44	İ
7	Malsiras	••	11.40	16-68	6.82	18-16	1.47	1.41	16.70	0.88	5.23	8.48	
		Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	


ļ		Ì					1	Percentages.						
to.	Name of Taluk	.	Jowar.	Bajri.	Cotton.	Safflower.	Tur.	Groundnut.	Math.	Hemp.	Wheat.	Sugarcane.	Others.	Total.
			Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent
1	Sholapur	••	68-46	9.68	2·16	2.99	7-11	1.71	0.38	1.84	0.87	0.09	4.71	100.0
2	Barei		57.72	4.98	2.72	0.61	6-28	12.49	0.11	1.98	1.92	0.08	. 11-24	100.0
8	Madha		48-14	20.26	8.62	4.44	4.22	1.48	1.96	2.99	. 0.69	0.58	6.62	100.0
4	Karmala		50.84	15.42	15.26	6.14	2.09	0.19	0.66	1.82	0.75	0.008	7.28	100.0
5	Pandharpur		58.91	21·10	5.11	4.68	0.77	0.58	1.94	0.47	1.27	0.02	5.49	100.0
6	Sangola		82.83	45.02	0.95	4.03	0.60	0.35	5.65	0.75	1.59	0.02	8 10	100.0
7	Malsiras		54.71	28.59	3.41	6.65	0.46	0.08	2.50	70.13	0.52	0.001	2.94	100.0

SATARA.

The following statement shows the distribution of important crops by Talukas.

Figures in the following statement show actual acreage.

						A							
No.	Name of Taluka.	Jowar.	Bajari.	Groundnut.	Math.	Gram.	Wheat.	Kulta.	Sugarcane.	Tobacco.	Turmer c.	Other crops.	Area.
		Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.
1	Satara	48 500	18,169	11,968	474	3,772	3,624	2,125	1,228	56	172	24,660	114,178
2	Jawali	19,605	1,634	4,485	33	1,199	3,879	701	- 170 376		56 .	37,037	68,799
3	Wai	40 138	34,494	2,172	3,403	3,447	3,809	4,428	376 825	9	352	25,476	118,104
4	Koregaon	47 848	40,098	5,547	1,447	7,103	4,069	4,850	251 527	59	239	20,432	131,938
5	Man	04 984	139,327	425	27,135	834	2,981	4,367	74 508	41	4'	11,577	211,,029
6	Whate-	44.802	97,204	3,125	11,262	- 7,526	6,476	8,400	483 2,171	180	72	21,565	201,095
7	77	66 004	56,018	5,899	11,178	6,336	6,834	6,732	796 1,508	148	1,342	26,316	187,603
•	-	66,608	14,449	6,880	4,417	8,782	2,754	6,237	472 1,031	1,693	717	27 793	140,862
8	Tasgaon	78,130	8,441	14,702	2,556	13,679	7,329	2,533	2.433 3,774	4,748	887	71,478	201,911
P	Walwa	61,419	15,718	18,007	1,763	4,986	1,952	2,051	2,294	1,221	1,644	24,257	135,312
10	Karad		1,188	18,005	5	1,572	1,780	246	535	16	200	57,931	107,959
11	Patan	26,481	1,180								<u>'</u>		
	Total	523,803	421,740	91,215	63,733	59,236	45,487	42,670	8,533 16,742	2,166	5,685	848,522	1,618,790




SATARA—continued.

Slatement showing the distribution of important crops by Talukas; the distribution is shown by percentages.

						- ;	B			.,	,g	
No.	Name of Taluk	a. :				•	Percentages of	Crops.				
			Jowar.	Bajri.	Ground nut.	Math.	Gram,	Wheat.	Kulthi.	Sugarcane.	Tobacco.	Turmeric.
1	Satara		Per cent.	Per cent.	Per cent.	Per cent	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.
2	Jawali		8.74	0.89	4.92	0.05	2.02	8.53	1.64	7·82 1·99 2·24		0.58
3	Wai	••	7.66	8·18	2.88	5.84	5 · 82	8 - 37	10.87	4.41	0-11	6·19
4	Koregaon	••	9·13	9.50	6.08	2-27	11.99	8.94	11.87	2·94 8·14	0 72	4.20
5	Man	••	4.63	88.08	0.46	42.57	1.40	6.55	10.53	3.02	0.20	0.07
6	Khatav	••	8.55	23 · 61	8 · 43	17.67	12.72	14.24	19.69	5·66 12·96	2.50	1.27
. 7	Khanapur	••	12:60	13.58	6.47	17.54	10.68	15.02	15.77	9.02	1.81	23 · 61
8	Taegaon	••	12.72	8.42	7.54	7.04	14.83	6.02	14.62	6.53	20.78	12.61
9	Walwa	••	14.92	0.84	16.11	4.02	28.01	16.12	5 · 94	28.52	58.08	15.60
10	Karad	••	11.72	8 . 78	19.75	2.77	8 -44	4.29	4.82	26.88	14.97	28.92
11	Patan	••	5.07	0.28	19.74	0.09	2.65	3.02	0.57	6 · 27 7 · 66	0.19	3.2
	Total		100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

SATARA—concluded.

				•				Percentag	es.					
No.	Name of Ta	duka.	Jowar.	Bajri.	Ground- nuts.	Math.	Gram.	Wheat.	Kulthi.	Sugarcane.	Tobacco.	Turmeric.	Others.	Total.
			Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent
1	Satara	••	42-49	15.91	10.48	0.43	8.30	8.17	1.86	0.57	0.05	0.15	21.59	100.00
2	JawaJi		28 · 50	2.37	6.52	0.06	7 - 74	5.64	1.02	0.24	0.00	0.08	53 · 83	100.00
8	Wai		83.98	29 · 21	1.84	2.88	2.92	8.22	8.75	0.82	. 0.08	0.30	21.60	100.00
4	Koregaon		86 · 26	80.39	4 · 20	1.10	5.38	8.08-	3.67	0.19	0.04	0.18	15.61	100.00
5	Man	••	11.50	66 · 02	0.20	12.86	0.39	1.41	2.07	0.08	0.02	0.002	5.50	100.00
6	Khatav		22 · 28	48.34	1.55	5.60	8 • 74	8.22	4.18	0.24	0.09	0.03	10.73	100-00
7	Khanapur		85.18	29 86	8.14	5.98	8.37	8 64	8 59	0.42	0.08	0.72	14.04	100-00
8	Tasgaon		47 · 29	10.26	4.88	8.16	6 - 23	1.95	4.48	0.88	1.20	0.51	17.76	100.00
9	Waiwa	••	88 - 69	1.70	7 · 28	1.27	6.77	3.63	1.25	1.20	2.85	0.44	85.42	100.00
10	Karad		45 · 39	11-62	13.31	1.30	3.68	1.44	1.51	1.69	0.90	1.21	17 - 95	100.00
11	Patan		24 · 53	1.10	16 · 68	0.005	1.46	1-65	0.23	0.50	0.01	0.18	53.66	100.00



=

BELGAUM.

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreage.

A

No.	Name of Taluka.	Jowar.	Cotton.	Bajri.	Rice.	Wheat.	Kulthi.	Tur.	Tobacco.	Safflower.	Sugarcane.	Others.	Total.
_		Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres,	Acres,	Acres.	Acres.
1	Belgaum	14,276	4,079	4,927	82,902	1,458	3,148	1,513	21	204	792 1,487	44,252	107,567
2	Hukerl	48,274	9,047	18,953	8,363	977	5,804	6,017	4,802	1,875	2,698 3,613	45,879	142,189
8	Chikodi	66,793	18,690	27,186	2,507	8,025	7,407	11,060	21,629	4,812	4,214 5,062	54,106	2 20,929
4	Athani	158,172	74,394	59,691	263	14,508	7,065	4,021	311	7,666	738	85,782	862,356
5	Gokak	95,219	58,780	55,455	267	20,700	10,245	5,289	1,025	3,860	756 1,278	50,860	802,456
6	Parasgad	92,125	66,739	25,638	101	47,218	11,243	10,129	34	7,466	159 294	43,436	804,282
7	Sampgaon	64,886	26,041	8,587	19,256	11,478	8,654	6,603	70	2,246	508 1,718	42,324	185,646
8	Khanapur	2,972	227	481	44,127	16	4,160	265 •	0	9	502 937	23,669	76,878
	To al	537,717	257,997	195,863	102,786	99,165	57,726	44,897	27,892	27,188	10,364 16,060	8,40,258	1,701,803

Statement showing the distribution of important crops by Talukas; the distribution is shown by percentages.

В

No.	Name of Taluka.		Percentages of crops.											
	Heme or yearn	-	Jowar.	Cotton.	Bajri.	Rice.	Wheat.	Kulthi.	Tur.	Tobacco.	Safflower.	Sugarcane.		
			Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.		
1	Belgaum		2.66	1.28	2.21	32.01	1.47	5-45	8 · 37	0.08	0.75	7.64		
2	Hukeri		8*05	8.21	9.68	8 27	0.99	10.02	13.42	17 · 22	5.08	26·03 22·49	0	
3	Chikodi	••	12-42	7 · 24	13.88	2.44	8.05	12.83	24.63	77 55	15.89	44 · 66 31 · 51		
4	Athani		29.41	28 - 83	30 · 47	0.58	14.42	12.24	8-93	1.11	28 25	7.12		
5	Gokak		17.71	22.78	28:31	0.58	20 87	17.75	11.78	8.67	14.22	7:30		
6	Parasgad		17.13	25 · 87	13.09	0.10	47.62	19:48	22.56	0.12	27.51	1.52		
7	Sampgaon		12.06	10.10	1.83	18 73	11.57	14-99	14.71	0.25	8.28	4·88 10·68		
8	Khanapur		0.55	0.07	0.55	42.93	0.01	7.21	0.59		0.03	4·84 5·83		
	Total		100.00	100*00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00		

BELGAUM—concluded.

No.	Name of Taluka.		Percentages.												
NO.	Name of Taru	ка.	Jowar.	Cotton	Bajari. Per cent.	Rice.	Wheat.	Kulthi.	Tur.	Tobacco. Per cent.	Safflower. Per cent.	Sugarcane. Per cent.	Others. Per cent.	Total.	
			Per cent.												
1	Belgaum		18.27	8.79	4.58	80.28	1.85	2.93	1.41	0.02	0.19	0.78	41.14	100.00	
2	Hukerl	٠.	80.48	6.36	18.33	2.86	0.69	4.08	4.28	3.38	0.97	1.90	82.26	100.00	
8	Chikodi		80.28	8.46	12.30	1.18	1.37	3.32	5.00	9.79	1.95	1.91	24.51	100.00	
4	Athani		43.65	20.23	16.47	0.07	8.95	1.95	1.11	0.09	2.12	. 0*20	9.86	100.00	
5	Gokak		81.48	19.43	18.88	0.09	6.84	3.39	1.76	0.84	1.28	0.52	16.81	100.00	
6	Parasgad		80.28	21.93	8.42	0.08	15.2	3.69	8.83	0.01	2.45	0.02	14.29	100.00	
7	Sampgaon	٠.	84.95	14.03	1.98	10.37	6.18	4.66	8.56	0.04	1.21	0.27	22-80	100.00	
8	Khanapur		3.89	0.30	0.56	57.77	0.02	5.45	0.85	0.00	0.01	0.66	80.88	100.00	


BIJAPUR.

The following statement shows the distribution of important crops by Talukas.

[The figures in the following statement show actual acreage.]

A

No.	Name of Talu	ka.	Jowar.	Cotton.	Bajri.	Wheat.	Safflower.	Tur.	Linseed.	Kulthi.	Gram.	Sugarcane.	Others.	Tótal.
			Acres.	Acres.	Acres.	Acres,	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres,
1	Indi	••	226,627	57,605	108,538	9,547	16,109	7,668	5,674	1,662	4,149	142 380	16,820	44,954
2	Sindagi		222,317	88,681	69,224	24,034	13,097	11,033	11,183	3,570	4,449	<u>31</u>	22,269	469,888
3	Bijapur		189,688	106,946	66,797	26,658	12 526	3,233	4,456	3,508	5,933	97 422	15,238	435,070
4	Bagewadi	••	171,795	112,728	39,202	46,254	11,422	7,030	11,930	8,752	5,657	102 298	14,636	429,308
5	Mudebihal		187,887	98,780	80,412	16,277	4,000	10,285	4,842	5,176	4,070	<u>18</u>	9,940	321,687
6	Bagalkot		154,883	89,488	15,926	17,288	5,440	10,275	5,166	4,428	4,863	88	16,811	324,606
7	Badami		89,654	87,702	44,609	10,706	2,562	11,449	1,656	6,953	8,155	180 252	33,453	242,079
8	Hundgund	••	118,829	101,628	4,086	12,572	2,902	6,508	4,529	4,936	5,884	- 27 31	21,131	283,082
i	, Total	••	1,311,680	693,558	873,594	163,831	68,058	67,481	49,486	88,980	88,160	635	150,298	2,955,211



BIJAPUR—continued.

Statement showing the distribution of important crops by Talukas; the distribution is shown by percentages.


B.

No.	Name of Taluka.	Jowar.	Cotton.	Bajri.	Wheat.	Satilower.	Tur.	Linseed.	Kulthi.	Gram.	Sugarcane.	
		Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	
1	Indi	17:28	8.81	27 · 71	5.84	23 · 67	11.36	11-48	4 - 24	10.87	22.80	
2	Sindagi	16*95	12-79	18.28	11.71	24	16.85	22.62	9.16	11.66	9.48	,
3	Bijapur	14.46	15.42	17:88	16.82	18.40	4.79	9.01	8.99	15 55	15·27 24·77	
4	Bagewadi	18-10	16 · 25	10.44	26.32	16.78	10.42	24 · 18	22.45	14.82	16.06	
5	Mudebihal	10.21	14 . 24	8.14	9.97	5.88	15 - 24	9 - 79	18 · 28	10.67	2.83	
ð	Bagalkot	11.81	12.90	4 • 26	10.28	7.99	15 • 22	10.45	11.36	12.74	-5·98 5·18	
7	Badami	6.83	5-44	11 - 94	6.22	8.76	16 - 97	3.35	17.84	8 27	28·35 14·80	•
8	Hundgund	9.06	14.65	1.09	7.69	4 · 26	9.64	9·16	12.66	15.42	1 82	
						i						
٠.	Total	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	

Statement showing how the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

C

	Name of Tal	nka					:	Percentages -		_				
0.	Name of Tar	una.	Jowar.	Cotton.	Bajri.	Wheat.	Safflower.	Tur.	Linseed.	Kulthi.	Gram.	Sugarcane.	Others.	Total.
			Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent .	Per cent.
1	Indi		50.41	12.81	23 . 08	2.12	3.28	1.71	1.26	0.87	0.92	0.03	8 . 74	100.00
2	Sindgi	• •	47.31	18 - 87	14.78	5.11	2.79	2.35	2.38	0.75	0.94	0.007	4.74	100.00
8	Bijapur		48.60	24.58	15 85	6.18	2.88	0.74	1.02	0.81	1.36	0.02	8.50	100.00
4	Bagewadi		40.02	26 - 26	9.08	10.77	2.66	1.64	2.78	2.04	1.82	0.02	8 • 41	100.00
5	Mudebihal		42.86	80.17	9.45	5.06	1.24	3 · 20	1.21	1.61	1.26	0.008	8.09	100.00
6	Bagalkot		47.71	27.57	4.91	5.82	1.67	8.17	1.59	1.36	1.50	0.01	5 · 18	100.00
7	Badami		87.08	15.57	18.43	4.42	1.06	4.78	0.68	2.87	1.30	0.07	13.82	100.00
8	Hundgund		41.98	85.91	1.44	4.44	1.08	2.30	1.60	1.74	2.08	0.01	7.46	100.00



DHARWAR.

The following statement shows the distribution of important crops by Talukas.

The figures in the following statement show actual acreage.

A

•	•					A								
No.	Name of Taluka.	Jowar.	Cotton.	Wheat.	Rice.	Rala.	Kulthi.	Tur.	Chilles.	Mug.	Sugarcane.	Others.	Total.	
		Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres,	
-1	Dharwar	42,078	29,261	18,911	19,187	6,265	6,879	3,986	689	2,089	116 388	16,252	145,719	
2	Hubli	44,599	55,449	86,798	8,709	2,112	3,189	4,149	508	8,861	15	15,591	169,975	
8	Navalgund	83,601	102,859	112,104		1,568	1,548	6,358	66	8,978	-8-9-	83,668	845,758	
4	Gadag	109,091	118,608	51,870	136	19,288	6,337	9,279	870	9,945	140 357	88,077	863,141	113
5	Bankapur	47,063	39,028	5,004	17,861	7,294	6,642	4,020	614	8,853	278 428	18,080	149,232	မ
6	Ranibenner	72,528	46,197	401	509	14,590	14,116	6,614	9,881	1,637	—54 99	11,404	177,428	
7	Hungul	18,441	10,365	82	45,889	5,146	6,728	1,602	1,331	8,275	244 1,115	14,045	102,148	
9	Karajgi	81,242	65,189	1,256	2,966	14,190	12,246	6,624	8,994	7,458	- 198 877	15,499	210,862	
9	Kalghatgi	12,599	4,308	88	86,501	. 1,078	1,915	1,424	464	1,173	158 458	9,669	69,317	
10	Kod	38,522	12,332	38	27,381	5,728	. 18,014	4,002	31,185	3,840	1,978	30,608	167,365	
11	Ron	67,858	83,270	86,863	78	4,448	782	4 ,149	. 79	3,539	8 34	22,089	223,156	
	Total	612,617	566,866	263,365	158,717	81,697	73,396	52,207	48,676	44,643	1,924 5,284	224,980	2,124,088	

DHARWAR—continued.

Statement, showing the distribution of important crops by Talukas; the distribution is shown by percentages.

o. Na							Average of (горя.				
J. Na	ime of Talul	ka.	Jowar.	Cotton.	Wheat.	Rice.	Rala.	Kulthi.	Tur.	Chilles.	Mug.	Sugarcane.
1 Dba	arwar		Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.
Dna	arwar		6.83	5.14	7.18	12.48	7.66	9.87	7.58	1.41	4.67	6.03
Hut	bli	•	7.23	9.69	18 . 97	2 41	2.57	4.34	7.94	1 03	8.65	7·34 0·77 0·77
Nav	valgund		13.66	18:49	42.60		1.91	2·10	12:38	0.13	8.90	0.48
Gad	lag		17.85	20.72	19:69	0.08	23 · 69	8.63	17.80	0.76	22:27	9.27
Ban	nkapur	••	7:68	6 · 78	1.90	11.22	8.92	9.04	7.71	1.26	8.63	14.13
	nibennur	••	11.83	8·14	0.12	0.33	17 85	19 23	12.60	19-27	3.66	2.80
	ngul	••	2.19	1.83	0.03	29 · 85	6 · 29	9·16	. 8*06	2.73	7.33	12.68
	rajgi		13.25	11.21	0.47	1-93	17:36	16.68	12.68	8.10	16.95	10·29 7·13
	ighatgi	••	2.05	0.76	0.01	23 78	1 31	2.60	2.72	0.95	2.52	7.95
Kod	·		6 · 28	2 16	0.01	17.80	7 01	17.18	7 56	64.06	8.60	87·15 87·46
Ron	a		11.07	14.68	13.99	0.05	5.42	1.08	7.94	0.10	7.82	0·41 0·64
	Total		100.00	100.00	100:00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

DHARWAR—concluded.

Statement showing how the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

C

. \							Percentag	ge.					
0.	Name of Taluka.	Jowar.	Cotton.	Wheat.	Rice.	Rala.	Kulthi.	Tur.	Chilles.	Mug.	Sugarcane.	Others.	Total.
1	Dharwar	Per cent. 28 80	Per cent, 20 08	Per cent. 12.98	Per cent. 13·16	Per cent. 4 23	Per cent.	Per cent. 2.73	Per cent. 0 · 47	Per cent. 1'43	Per ce nt. 0.07	Per cent. 11.33	Per cent.
2	Hubli	26 · 28	82.62	21.64	2·18	1.24	1.87	2:44	0.29	2.51	0.00	9.28	100.00
8	Navalgund	24.18	29.74	82.45	٠.,	0.45	0.44	1.83	0.01	1.14	0.00	9.76	100.00
4	Gadag	80.04	82.66	14.28	0.08	5 .81	1.74	2.55	0.10	2.78	00.03	10.28	100.00
5	Bankapur	81.28	26 · 21	8 85	11.63	4.82	4.45	2 69	0.41	2.40	0.18	12.32	100.00
6	Ranibennur	40 87	26 04	2.22	0.28	8.22	7.95	8.72	5 · 28	0.92	0.08	6.47	100.00
7	Hangul	18.15	10.14	0.08	44.92	5:03	6.58	1.26	1.30	8.50	0.53	18.81	100.00
8	Karajgi .	38 - 52	80.91	0.59	1.40	6-72	5.70	8.14	1.87	8.53	0.09	7.58	100.00
9	Kalghatgi .	18-17	6.21	0.02	52.65	1.24	2.76	2.05	0.66	1.68	0.55	14.01	100.00
10	Kod .	28.01	7.80	0.02	16.36	3.42	7.82	2.89	18.63	2.29	0.42	18.84	100.00
11	Ron .	80.40	87.81	16.20	0.08	1.90	0.85	1.85	0.03	1.28		10.05	100:00

115

The following statement shows the distribution of important crops by Talukas.

Figures in the following statement show actual acreage,

A

	•						Д	•							
No.	Name of Taluka		Rice.	Nagli.	Varai.	Udid.	Kođra.	Niger seed.	Tur.	Wal.	Gram.	Sugar cane.	Others.	Total.	
1	Shahapur		Acres. 31,673	Acres. 9,252	Acres. 4,578	Acres. 2,097	Acres.	Actes. 1,440	Acres. 98	Acres.	Acres.	Acres.	Acres. 1,289	Acres. 50,458	
2	Murbad		29,646	9,325	2,923	2,643		1,364	102	46	188		1,693	47,930	
3	Salsette		31,629	317	79	65	1	25	2	18			3,327	35,443	1
4	Bassein	••	. 32,789	1,433	192	591	259	55	175	. 787	188	136	8,532	40,143	
. 5	Kalyan	إ	36,853	2,040	658	817		264	67	489	473		928	41,589	
6	Dahanu	••	51,725	6,845	1,364	2,133	7,544	594	1,948	1,978	221	27 48	4,461	78,820	
7	Wada		26,541	16,738	6,038	2,493	••	2,799	1,714	18	281	·	1,162	57,784	
8	Bhivandi	••	41,463	2,335	176	412		277	81	210	499	••	1,150	46,603	
9	Mahim	•	44,731	1,825	489	1,013	306	111	285	835	174	230 326	985	50,484	
	Total		326,550	50,110	16,492	11,764	8,110	6,929	4,472	8,892	2,049	899	18,487	449,254	

THANA—continued.

Statement showing the distribution of important crops by Talukas, the distribution is shown by percentages.

В

Ì						P	ercentages of	crops.				
»	Name of Talul		Rice.	Nagli.	Varai.	Vdid.	Kodra.	Niger seed .	Tur.	Wal.	Gram.	Sugarcane.
1	Shahapur		Per cent.	Per cent. 18.45	Per cent. 27 · 78	Per cent. 17:83	Per cent.	Per cent. 20 78	Per cent. 2·19	Per cent .	Per cent. 1.22	Per cent.
2	Murbad		9.07	18.61	17.72	22 · 47	••	.19.68	2.28	1.18	9 17	
8	Salsette		9.69	0.68	0.48	0.22	0.01	0.86	0.04	0.46		
	Bassein		10.04	2.86	1.16	5.02	8.19	0.79	8.90	20 · 22	9.17	35·50 26·66
5	Kalyan		11.13	4 07	8.99	2.69	••	8.81	1.20	12.26	28 08	••
	Dahanu		15 84	18.66	8.27	18.13	93.02	8 57	43.26	50.82	10.79	-6·76 9·41
.	Wada		8.13	88-40	86.61	21 · 19		40.40	38 33	0.46	13.71	••
۱	Bhivandi		12.69	4.66	1.07	8.20	••	4.00	1.81	5 39	24.85	
,	Mahim		18.70	3.64	2.96	8.61	8.78	1.60	6 · 87	8-61	8 48	57·65 68·92
	Total		100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

THANA—concluded.

Statement showing how the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

 \mathbf{C}

No.	Name of Ta	lako						Percentages	i .		•			*
NO.			Rice.	Nagli.	Varai.	Vdid.	Kodra.	Niger seed.	Tur.	Wal.	Gram.	Sugarcane.	Others.	Total.
1	Shahapur		Per cent. 62.77	Per cent. 18 ⁻ 84	Per cent. 9.06	Per cent.	Per cent.	Per cent.	Per cent. 0·19	Per cent.	Por cent.	Per cent.	Per cent. 2.55	Per cent. 100 '00
2	Murbad		61.85	19 · 46	6.09	5.21		2.84	0.21	0.09	0.89		8 · 54	100.00
8	Salsette	••	89.24	0.89	0.55	0.18	0.003	0.07	0.005	0.02	••	l	9.33	100.00
4	Bassein		81.68	8.57	0.48	1.47	0.84	0.14	0.48	1.96	0.47	0.35	8.80	100.00
5	Kalyan		87 - 41	4.90	1 58	0.77		0.68	0.16	1.18	1.14		2 23	100.00
6	Dahanu		65 · 62	8.68	1.73	2.70	9.57	0.75	2.47	2.50	0.28	0.08	5.66	100.00
7	Wada	••	45 . 93	28 - 97	10.45	4.81	••	4.84	2.97	0.08	0.49		2.01	100.00
8	Bhivandi		88 . 96	5.01	0.39	0.88		0.59	0.17	0.45	1.07		2 · 47	100.00
P	Mahim		88 -60	8.61	0.97	2.00	0 61	0.22	0.56	0.66	0.84	0.46	1.95	100.00

KOLABA

	KULABA	Fig 4â
ONE DOT REPRESENTS , 1000 ACRES.	NAGLI	WARI
WTS 100 ACRES.	WAL	VOID
SESAMUM SESAMUM	TUR	MUG
I ALIDAG 2 PEN 3 PENYEL 4 MARJAT 3 ROHA 1 MARGADN 7 MANAD 1 ALIDAG 2 PEN 3 PANYEL 4 MARJAT 5 ROMA 6 MARJAT 5 ROMA 6 MARJAT 7 MANAD	WAA.	7UN
ALIBAG 2 PEN 3 PANYEL 4 NARMT 5 ROM 6 MANGAON 17 MAMAO	<i>U9:0</i>	

KOLABA.

The following statement shows the distribution of important crops by Talukas.

The figures in the following statement show actual acreage.

A

0.	Name of Talul	- 1	Rice.	Nagli.	Varai.	Kodra.	Wal.	Vdid.	Sesamum.	Tur.	Mug.	Niger seed.	Others.	Total.
1	Allbag		Acres. 40,252	Acres. 366	Acres. 213	Acres.	Acres. 1,601	Acres. 83	Acres.	Acres.	Acres. 21	Acres,	Acres. 2,093	Aeres. 44,653
2	Pen		38,245	2,602	1,800	1	680	88	99	93	107	100	167	43,982
8	Panvel		5 2 ,342	913	555		789	80	207	71	171	13	920	55,961
4	Karjat		35,910	3,983	2,272	20	743	219	244	56	115	213	824	44,099
5	Roha		21,641	8,479	3,262	678	1,899	29	194	83	171	99	76	87,611
6	Mangaon]	32,398	11,066	10,371	5,229	1,699	1,163	461	868	575	728	820	64,878
7	Mahad		84,156	19,238	12,655	3,054	1,025	2,011	1,223	862	550	162	89	75,025
	Total		254,944	41,647	81,128	8,982	8,386	3,623	2,428-	2,045	1,710	1,327	3,989	860,209

KOLABA—continued.

Statement showing the distribution of important crops by Talukas; the distribution is shown by percentages.

В

No.	Name of Ta	duka.	Rice.	Nagli.	Varai.	Kodra.	Wal.	Udid.	Sesamum.	Tur.	Mug.	Niger Seed.
1]	Alibag		Per cent.	Per cent.	Per cent. 0.68	Per cent.	Per cent. 19:09	Per cent. 2·29	Per cent.	Per cent. (Per cent. 1.23	Per cent.
2] 1	Pen		15 001	6.25	5.78	0.01	8.11	2.43	4.08	4.55	6 · 26	7:54
8 1	Panvel		20 58	2.19	1.78		8.81	0.83	. 8.52	3.47	10.00	0.88
4]	Karjat	••	14.09	9.54	7.80	0.22	8.86	6.04	10.05	2.74	6.72	16.02
5 1	Roha.		8.48	8.35	10.48	7.54	22:64	0.80	7.99	4.06	10.00	7.46
6 1	Mangaon		12.71	26.57	88-82	58.21	20.26	82.10	18.98	42.44	38.62	54.86
7 1	Mahad		18.40	46.19	40.65	84.001	12.22	55.50	50.87	42.15	32.16	12.20

KOIABA—concluded.

Statement showing the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

C

о.	Name of Ta	e Imba						Perce	ntages.					
٥.	Wante OT 18		Rice.	Nagli.	Varai.	Kodra.	Wal	Vdid.	Setamum.	Tur.	Mug.	Niger seed.	Others.	Total.
1	Alibag		Per cent. 90·14	Per cent. 0.82	Per cent.	Per cent.	Per cent. 8.58	Per cent. 0·18	Per cent.	Per cent.	Per cent. 0.04	Per cent.	Per cent.	Per cent. 100·00
2	Pen		86.95	5.91	4.09	0.002	1.55	0.20	0.23	0.21	0.24	0.23	0.88	100.00
3	Panvel		98 58	1.63	0.99		1.82	0.05	0.86	0.18	0.81	0.02	1.64	100.00
ı	Karjat		81-43	9.03	5:15	0.05	1·68	0.50	0.55	0.18	0.26	0-48	0-78	100.00
5	Roha		68-46	11.005	10.82	2.14	6.007	0.09	0.61	0.26	0.54	0.81	0.24	100.00
В	Mangaon	••	49.94	17.06	15.99	8.06	2.62	1.79	0.71	1.82	0.89	1.12	0.49	100.00
7	Mahad		45.52	25 - 64	16.87	4.07	1.37	2.68	1.63	1.15	0.78	0.21	0.12	100.00

The following statement shows the distribution of important crops by Talukas. The figures in the following statement show actual acreages.

[Black figures represent the estimated acreage of the area for which figures represent the estimated acreage of the area for which figures represent the estimated acreage of the area for which figures represent the estimated acreage of the area for which figures.

A

о.	Name of Tale	oka.	Rice.	Nagli.	Kodra.	Varai.	Kulthi.	Niger Seed.	Udid.	Wal.	Tag.	Sugarcane.	Other crops.	Total cropped area.
1	V engur ia	••	Acres. 8,275	Acres. 1,862	Acres. 949	Acres. 369	Acres. 1,199	Acres.	Acres. 775	Acres.	Acres. 52	Acres.	Acres, 4,125	Acres. 18,365
ļ			25,514	7,894	3,140	1,221	8,968	2,468	2,564	••	172	43 0	13,650	60,634
2	Malvan		23,064	6,575	4,314	1,667	5,392	1,891	1,122	25	275	871	4,439	49,185
			71,113	27,876	14,275	5,516	17,842	6,257	8,713	83	909	1,229 371	14,687	1,63,500
3	Devgad		9,067	4,156	1,502	1,000	1,665	883	478	94	1,581	103	886	• 21,415
		•	27,956	17,620	4,970	8,309	5,509	2,922	1,581	811	5,232	343 486	2,932	72,685
4	Rajapur		6,617	2,444	969	1,360	602	416	268	168	483	26	457	13,810
			20,402	10,362	3,206	4,500	1,992	1,377	887	556	1,598	86	1,512	48,478
5	Ratnagiri		9,298	4,518	2,839	2,256	898	1,386	204	268	85	15	1,136	22,403
			28,669	19,155	9,394	7,465	1,316	4,586	675	887	281	48 21	3,759	76,235

122

	RAINAGIRI	Fig.43
RIGE STATES OF THE STATES OF T	NAGLI	KODRA
WA ARES	KULTHI	NIGER-SEED
ONE DOT REPRESENTS	WAL	TAG
Momes of Tables 1 ATT ATT ATT ATT ATT ATT ATT ATT ATT A		7 M O U S A Apple O S II
I VENGUELA 2 MAI MAN 3 OEDGAD 4 RANAPUR 5 SANTMGINI 6 SANTMGINI 7 CHIPLUM 8 RANED	No. Tr.	
/ VENGURLA 2 MALWAN 3050CAD 4 RAUAPUR 5 SANTMAGIRI 5 SANGMESHMAR 7 CHIDE UN 6 MARBO 30APOLI		

6	Sangameshvar	••	3,209	2,719	1,460	749	20	158	38	490	2	6	89	8,935	
			9,894	11,528	4,831	2,478	66	523	109	1,620	7	- 19	295	31,370	
7	Chiplun	••	2,809	1,796	979	647	11	7	85	754	8	1	433	7,480	
			8,661	7,614	8,240	2,141	87	23	116	2,497	. 27	- 8	1,433	25,792	
. 8	Khed		7,181	6,451	2,281	2,175	1	· 216	. 97	892	19	••	93	18,906	
			22,141	27,350	7,548	7,197	3	715	821	1,296	63	:	308	66,942	
9	Dapoli	••	6,971	5,347	2,553	2,452	21	840	820	664	88	٠.	952	20,212	
			21,494	22,669	8,448	8,114	70	2,780	1,059	2,197	291	<u> 13</u>	8,150	70,285	
													[
	Total		76,491	85,868	17,846	12,675	9,809	6,543	8,832	2,855	2,598	539	12,610	1,80,661	F
			2,85,844	1,52,068	59 ,052	41,941	80,803	21,651	11,025	9,447	8,580	1,784	41,726	6,13,921	2

RATNAGIRI—continued.

Statement showing the distribution of important crops by Talukas, the distribution is shown by percentages.

 \mathbf{B}

No.	Name of Taluka.	Rice.	Nagii.	Kodra.	Varai.	Kulthi.	Niger seed.	Udid.	Wal.	Tag.	Sugarcane.
_		Por cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent .	Per cent.	Per cent.	P r cent.
1	Venguria	10.82	5.19	5.82	2.91	12.88	11.40	23 - 26	0.00	2.00	2·41 0·91
2	Maivan	80.12	18.84	24·17	13 · 15	57.98	28.90.	83.68	0.88	10.59	83.85
8	Devgad	11 ·85	11 · 59	8-48	7 · 89`	17.85	13.50	14.34	8 · 29	60.95	19·23 44·47
4	Rajapur	8 - 65	6.81	5.43	10.72	6 · 47	6.36	8.02	5.89	18 63	9.69
5	Ratnagiri	12·16	12-60	15 - 91	17.80	4 · 28	21 · 14	6.12	9.39	8.58	2·69 1·92
6	Sangameshvar	4.19	7-59	8·18	5.80	0.22	2-42	0.89	17.15	0.09	1.06 1.47
7	Chiplun	3 · 67	5.00	5.48	5.11	0.13	0.11	1.05	26 · 43	0.33	0·17 0·82
8	Khed	9.89	17 - 98	12.78	17·16	.0.01	8.83	2.91	13.72	0.73	0.00
9	Dapoli	9.12	14-90	14.80	19:36	0·23	12.84	9-60	23 - 25	8.40	0.48
	Total	100.00	100.00	100 00	100.00	100.00	100.00	100.00	100.00	100.00	100.00

		Fig 44
OME DOT REPRESENTS 1001 ACRES	BETLE-NUT	PEPPER
CANTO 100 ACRES	SUGAR-CANE	KULTHI
ONE DOT REPRESENTS	CARDAMUM	CHAVLI
HOMES OF TRANSPORT I RAPINEAR 2 ANNOL A 3 NUMBER 4 STORAMURA 5 STORAMURA 5 STORAMURA 6 ST	WAGAS AND THE STREET	0 4 5 A 7 0 3.
I RAMMAR I RAMMAR I RAMMA 3 RUMMA 4 HOMMAR 5 GIODAPUR IS SIRS! YYELLAPUR 8 MALIYAL	97	CARDAMAN
IRARRIAG EANKOLA 3 RUMTA HOMANIAR ESIDOAPUR LERKSI TYELLARUR GMALIMAL		

RATNAGIRI—concluded.

Statement showing how the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

С

.	N							Percent	sges.					
io.	Name of Talul		Rice.	Nagli.	Kodra.	Varai.	Kulthi.	Niger seed.	Vdid.	Wal.	Tag.	Sugarcane.	Other crops.	Total.
1	Vengurla		Per cent. 42.08	Per cent. 18.02	Per cent. 5 18	Per cent. 2·01	Per cent. 6 · 55	Per cent.	Per cent. 4·28	Per cent.	Per cent. 0.28	Per cent. 0.07	Per cent. 22·51	er cent 100 00
2 1	Malvan		43.50	17.05	8 . 78	8 · 87	10.91	3.83	2.27	0.02	0.28	0.75	8.88	100.00
8 1	Devgad		88 - 46	24 · 24	6.84	4.55	7.58	4.02	2.18	0.43	7.20	0.47	4.08	100.00
4 3	Rajapur		48.90	22.29	6.90	9.68	4 · 29	2.96	1 . 91	1.20	8.44	0.13	8 · 25	100.00
5 1	Ratnagiri		87.61	25 · 18	12.82	9 - 79	1.78	6.02	0.88	1.16	0.87	0.08	4.98	100.00
6 8	Sangameswar	٠.,	81 54	36 75	15 · 40	7.90	0.21	1.67	. 0.85	5.16	0.02	0.08	0.94	100 .00
7 0	Chiplun		83.58	29.51	12.56	8 · 80	0.15	0.09	0.45	89.8	0.11	0.01	5.26	100.00
8 1	Khed		88 07	40.86	11.28	10.75	0.00	1.07	0.48	1.94	0.08	0.00	0.46	100.00
9] 1	Dapoli		80.28	82 · 25	12.02	11.24	0.10	8.98	1.21	3.18	0.41	0.02	4 - 41	100 .00

12

12

KANARA.

The following statement shows the distribution of important crops by Talukas.

The figures in the following statement show actual creage.

A

					•			А	•						-	
No.	Name o	of Taluk	а.	Rice.	Betlenuts.	Miri pepper.	Nagli.	Sugar- cane.	Kulthi.	Udid.	Cardamom.	Chavli.	Mug.	other.	Total cropped Area.	
			_ i	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	Acres.	
1	Karwar			23,997	38	••	49	203 181	3	219	••	21	197	2,610	27,832	
2	Ankola			16,363	880	26	165	262 242	10	846	7	119	181	2,876	20,735	
3	Kumta			19,754	654	58	277	214 218	5	389	••	440	107	8,710	25,603	
- 4	Honavar	,		25,455	1,686	89	834	672 765	818	518		589	320	6,319	86,75	
5	Siddapu	•		12,475	6,383	2,123	42	886 858	80	97	736	••	8	2,159	24,439	
6	Sirai			19,245	5,795	1,820	89	467 644	208	90	550	8	12	2,462	80,541	
7.	Yellapur	•		20,588	2,419	297	174	463 800	195	70	100	6	80	2,193	26,480	
8	Haliyal			35,459	169	2	1,495	218 226	1,265	10	2	2	86	2,876	41,534	
	-	Total	•	178,281	18,019	4,160	3,125	2,885 2,934	2,084	1,789	1,895	1,180	891	24,705	2,33,414	

KANARA-continued.

Statement showing the distribution of important crops by Talukas, the distribution is shown by percentages.

B

o.	Name of Taluka.					Percentages of	f Crops.					
0.		Rice.	Betlenute.	Mirt.	Nagli.	Sugarcane.	Kulthi.	Vaid.	Cardamom.	Chavli.	Mug.	
_		Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	Per cent.	
1	Karwar	13.85	0.18		1.26	7·04 6·16	0.12	12.59		1.78	22.11	
2	Ankola	9.44	4.88	0.62	5.28	9·08 8·24	0.49	19.89	. 0.50	10.08	20.81	
8	Kumta	11.40	8.68	1.27	8.86	7.41	0.54	22.37		87 · 29	12.01	į
4	Honavar	14.69	8.8¢	0.94	26 · 69	28 · 29 26 · 07	15 · 68	29.79		49.91	85 - 91	
5	Siddapur	7.20	35 · 42	51.08	1.34	13.88	1.48	5.58	52.76	••••	0·89 +1	
6	Sirsi	11,11	82·16	88.94	2.85	16·19 21·94	10.22	5.17	39.43	0.25	1 · 34 +1	٠
7	Yellapur	11.85	13.42	7.18	5 . 57	16.04	9.58	4.02	7.17	0.21	8 · 87	
8	Haliyal	20.46	0.94	0.02	47.84	7·55 7·66	62 19	0.57	0.14	0.17	4.01	
	Total	100.00	100.00	100.00	100.03	100.00	700.00	100.00	100.00	100.00	100.00	

KANARA—concluded.

Statement showing how the gross cropped area of each Taluka is allotted to different crops, the gross cropped area of each Taluka being represented by 100.

O

								Percentag	e s.					,
No.	Name of T	aluka.	Rice.	Betelnut.	Miri.	Nagli.	Sugarcane.	Kulthi.	Udid.	Cardamom.	Chavli.	Mug.	others.	Total.
1	Karwar		Per cent. 87.79	Per cent. 0.12	Per cent.	Per cent. 0.17	Per cent.	Per cent. 0.01	per cent. 0.80	Per cent.	Per cent. 0.08	Per cent. 0'72	Per cent. 9 54	Per cent.
2	Ankola		78 · 91	4.24	0.12	0.80	1.26	0.02	1 67	0.03	0.28	0.87	11-47	100 .00
8-	Kumta	٠.	77 · 15	2.55	0.21	1.08	0.84	0.02	1.52		1.72	0.42	14 · 49	100.00
4	Honavr	••	69-27	4.59	0.11	2.27	1.83	0.86	1.41		1.60	0.87	17:17	100.00
5	Siddapur	••	51.05	26 · 12	8.69	0.17	1.58	0.12	0.39	8.01		0.08	8.83	100.00
6	Sirsi		63 · 01	18 - 97	5.30	0.29	1.23	0.68	0.29	1.80	0.01	0.04	8 06	100.00
7	Yellapur	••	77.53	9.18	1.12	0.66	1.75	0.74	0.26	0.38	0.02	0.11	8 · 28	100.00
8	Haliyal		85 · 87	0.41	0.002	8.60	0.58	8.04	0.02	0.005	0.002	0.09	6.92	100.00
							1			ļ!		<u> </u>		

APPENDIX.

Glossary of Botanical, English and Vernacular names of crops.

Botanical Name.	_	English.		Marathi.		Gujarati.		Kanarese,
I.—Cerclas.						•	.	
Triticum		Wheat		Gahu		Gehu		Godhi.
Triticum Spelta		Spelta Wheat .		Khapli Jod				
Sorghum vulgare		Great millet		Jowari		Jowari		Jola.
Pennisetum typhoideum		Bulrush millet .		Bàjri		Bàjri		,.
Zea mays	••	Maize .	·	Maka		Makai		Mekkejol Goingol.
Orvza sativa		Rice ; Paddy .	\cdot	Bhàt]	Danga		Bhatta.
Eleneine coracana	••			Nagli or chani	Na-	Bavto	٠.	Ragi.
Panicum miliaceum				Vari		Gadio, Kuri		Save, Baragu
Panicum miliare				Sava		Cheno		
Hordeum vulgare		Barley .		Satu		Do.		Jave godhi.
Paspalum scrobiculatum	٠			Kodra		Kodra		Harik.
Setaria italica				Rala		Kang		Navani, Rai
Panicum crusgalli			-	Berti		Banti		Do.
II.—Pulses.		i						
Cajanus indicus		Pigeon-pea .		Tur		Tuver		Togari.
Vigna catiang		Cow-pea .		Chavall		Chola		Alsandi.
Dolichos biflorus		Horse-gram .	-	Hulga		Kulthi		Hulke.
Dolichos lab-lab		Indian bean .	··	Wal or Pawte	٠.,	Wa		Avare.
Cyamopsis psoralioides		Ciuster bean .	٠٠	Guwar		Guwar		Chivali kal.
Phaseolus mungo		Green gram .		Mug		Mag		Hesaru.
Phaseolus aconitifolius		Kidney bean .		Matki		Math		Madaki.
Phaseohus mungo var radiatus		Black gram .		Udid		Adad		Vdđu.
Lathyrus sativus		Chickling vetch	١.	Lakh		Lang		
Lens esculenta		Lentil .		Masur		Masur		Channangi.
Pisum sativum		Peas (white) .	إ	Vatana	٠.	Vatana.		Batagadle
Cicer arietinum		Gram .		Harbara	••	Chana		Kadle.
Oil Seeds.								
Arachis hypogæa		Groundnut .		Bhuimug		Bhoising	•	Bhaimug
Guizotia abyssynica		Niger seed .		Karla Khurasani.	0 T	Ramtal	•	Gurellu.

APPENDIX—continued.

<u> </u>				•				
Botanical name.	_	English.		Marathi.		Gujarati.	į	Kanarese.
Oil Seeds—continue	d.		_				_	
Sesamum indicum	• •	Sesamum		Til		Tal		Yellu.
Ricinus communis		Castor		Erand		Divel .		Audla.
Linum usitatissimum	٠!	Linseed	٠.,	Javas		Alsi		Agashi
Carthamus tinctorius		Safflower		Kardai		Kasambi		Kusubi.
Gossypium	••	Cotton		Kapas				
Others								
Nicotiana Tabacum	••	Tobacco	••	Tambakhu		Tamakhu		Hogesoppu Tabaku.
Saccharum officinarum	••	Sugarcane		Us		Sherdi		Kabbu-
Vegetables.					•			
Dioscorea bulbifera		Yam		Konfa Gora	đn	Goradu		Heggenasu.
Curcuma longa		Turmeric		Halad	••••	Haldar	••	Arshana Beru.
Zingiber officinale	•••	Ginger		Ale	•	Adu	••	Alla
Allium cepa		Onion		Kanda	•••	Dungri Kand		Ullegaddi.
Allium sativum		Garlic	•	Lascon		Lasan	•	Belloli.
Daucus carota	•••	Carrots		Gajar	••	Gajar	••	Gajjar
Raphanus sativus		Radish		Mula		Mulangi		Muri.
Solanum melongena	••		egg	Vangi	••	Vengan		Badnikai.
Capsicum sp.	••	Chillies		Mirchi		Marcha.		Menashinakal
Hiblscus esculentus		Lady's finger		Bhendi		Binda		Bhendekal.
Cucurbitacease sp.		Cucurbits			•		•	••••
Trigonella fœi græcum.	num	••••		Methi		Methi	••	Menthia.
Amarantus paniculatus			ł	Rajgira		Rajagro		Rajgiri.
Amarantus mangos tant	us	••••		Pokla.	•			••••
Rumex vesicariu		••••		Chuka				****
Chenopodium album			•	Chakwat				••••
Brassica oleracea		Cabbage		Kebi		Pangoli		Kobi gaddi.
Brassica caulocarpa		Cauliflower		Fulvar				••••
Brassica botrytis	••	Knol khol K	hol	Kaval nol	••			••••
Beta vulgaris	••		eet-	••••				••••
Lycoperaicum esculentu	ım .	Tomatoes		Bailwangl	••	Vilaiti vangar	١	Chappa badnikai,
Solanum tuberosum		Potato		Batata		Batata		Batata.

131
APPENDIX—concluded.

Musa paradisiaca Banana plan- Kele Balehannu. Psid'um Guyava Guava Peru Jamrukh Peral hann	Botanical name.		English.		Marathi.		Gujarati.		Kanarese.	
Citrus medica V a r Aurantium Narangi Narangi Kittale Ficus Carica , Fig Anjir Anjir Anjuri Mangifera Indica Mango Amba Ambo Ma v in h a n n Mavu Musa paradisiaca Banana plantain. Psid'um Guyava Guava Peru Jamrukh Peral hann Shebe hannu Punica Granatum Pomegranate Dalimb Dadum Dalimba	Fruil Trees.	Í	i			į			•	
Aurantium Narangi Narangi Kittale Ficus Carica , Fig Anjir Anjir Anjuri Mangifera Indica Mango Amba Ambo Ma v in h a n n Mavu Banana plan tain. Peru Jamrukh Peral hann Shebe hannu Punica Granatum Pomegranate Dalimb Dadum Dalimba	Citrus medica Var a	cida	••••		Limbu		Limbu		Limbe.	
Mangifera Indica Mango Amba Ambo Ma v in h a n n Mavu. Musa paradisiaca Banana plantain. Psid'um Guyava Guava Peru Jamrukh Peral hann Shebe hannu Punica Granatum Pomegranate Dalimb Dadum Dalimba			••••		Narangi		Narangi#	••	Kittale	
Musa paradisiaca Banana plankele Balehannu. Psid'um Guyava Guava Peru Jamrukh Peral hannu Punlea Granatum Pomegranate Dalimb Dadum Dalimbus	Ficus Carica	,.	Fig	••	Anjir		Anjir		Anjuri	
Psid nm Guyava Guava Peru Jamrukh Peral hann Shebe hannu Punica Granatum Pomegranate Dalimb Dadum Dalimbus	Mangifera Indica		Mango	••	Amba	••	Ambo	. ••	Mavinhannu Mavu,	
Punica Granatum Pomegranate Dalimb Dadum Dalimbus	Musa paradisiaca			lan-	Kele		••••		Balehannu.	
	Psid'um Guyava		Guava	••	Peru		Jamrukh	••	Peral hannu. Shebe hannu,	
Vitis Vinifera Grape Draksha Draksh. Draksh.	Punica Granatum		Pomegranate		Dalimb		Dadum		Dalimbu	
	Vitis Vinifera		Grape		Draksha	••	Draksh.	••	Drakh.	

132

Glossory of technical agricultural terms.

Term.			Explanation.				
	•	,	•				
Kharif	••		The term is applied to such crops as are grown in rainy season.				
Rabi	••	• •	The term is applied to such crops as are grown in winter.				
Jirayat	·· .		The term is applied to crops grown without irrigation.				
Bagayat	••	• •	The term is applied to crops grown with irrigation.				
Phusa	٠,	. •	Chaff left after the trodding out of the grain under the feet of bullocks on threshing flour.				
Rab	••		It is a term expressing the preparation of seed bed for Rice or other millets in the heavy rainfall tract by burning cowdung, tahal and grass arranged in layers one over the other.				
Tahai	• •	•	Branch loppings of trees.				

SOO3-OH