

Tables of Representative
Values of foods commonly
Used in Tropical countries

PRIVY COUNCIL

MEDICAL RESEARCH COUNCIL

SPECIAL REPORT SERIES No. 253

**TABLES OF REPRESENTATIVE
VALUES OF FOODS COMMONLY
USED IN TROPICAL COUNTRIES**

by

B. S. PLATT

LONDON: HIS MAJESTY'S STATIONERY OFFICE

1945

Price 9d. net

Universal Decimal Classification
613.2 (213)

MEDICAL RESEARCH COUNCIL

The Rt. Hon. **LORD BALFOUR OF BURLEIGH, D.L.** (*Chairman*).

Sir **WILLIAM M. GOODENOUGH, Bart., D.L., J.P.** (*Treasurer*).

Sir **CHARLES G. MACANDREW, T.D., M.P.**

Sir **HENRY H. DALE, O.M., G.B.E., M.D., F.R.C.P., F.R.S.**

Professor **D. KEILIN, Sc.D., F.R.S.**

Professor **F. C. BARTLETT, C.B.E., F.R.S.**

Professor **L. J. WITTS, M.D., F.R.C.P.**

Professor **J. R. LEARMONT, C.B.E., Ch.M., F.R.C.S.E.**

A. N. DRURY, C.B.E., M.D., F.R.S.

Professor **J. C. SPENCE, M.C., M.D., F.R.C.P.**

Professor **P. A. BUXTON, M.R.C.S., F.R.S.**

Professor **Sir ALEXANDER FLEMING, M.B., F.R.C.S., F.R.C.P., F.R.S.**

Sir EDWARD MELLANBY, K.C.B., M.D., F.R.C.P., F.R.S. (*Secretary*).

TABLES OF REPRESENTATIVE VALUES OF FOODS COMMONLY USED IN TROPICAL COUNTRIES

PREFACE

In the course of its work during the years 1936-9, the Committee on Nutrition in the Colonial Empire, which had been set up by the Economic Advisory Council, appointed a sub-committee on research, under the chairmanship of Sir Edward Mellanby.* The sub-committee's main recommendation was that there should be an organisation for co-ordinating a series of field surveys in the Colonial Territories, and the Medical Research Council undertook to appoint a small staff to serve as a scientific nucleus for this purpose. The first survey was made accordingly in Nyasaland under the direction of Dr. B. S. Platt, to determine the relationship of economic, sociological and agricultural matters to nutrition, as well as the correlation of health with the nutritive value of the diet.

This survey took place in 1938-40 and the first draft of a report on its results was made, but the final stages of preparation for publication could not be undertaken at the conclusion of the survey (June, 1940). The diversion of personnel to work on wartime problems prevented more than the maintenance of part-time advisory assistance to the Colonial Office, and only recently has it been possible to resume active preparation for the work for which the organisation was intended. In setting up their Human Nutrition Research Unit in 1944, the Council had the needs of the Colonial Territories in mind.

One of the objects of the original survey was to establish a technique, and it at once became clear that the provision of data on the nutritive value of the foods consumed was a basic need. For the purpose of studies on groups of individuals, representative values can be applied in the evaluation of dietaries in different areas, and one of the first tasks undertaken was the compilation of a table of food values for use in Nyasaland and areas consuming similar foods. This was the foundation from which the present tables have been built up. The latter thus represent the first published results of the Nyasaland Survey.

An illustrated glossary of terms is in preparation, to meet a similar need in relation to the recognition, nomenclature and interpretation of signs of ill-health due to nutritional deficiencies. A handbook on the methods of conducting nutritional surveys in Colonial Territories is also being prepared, and it is hoped that this will be followed by a report on the data collected during the Nyasaland Survey.

MEDICAL RESEARCH COUNCIL,
c/o London School of Hygiene and Tropical Medicine,
Keppel Street,
Gower Street, London, W.C.1.

29th October, 1945.

TABLES OF REPRESENTATIVE VALUES OF FOODS COMMONLY USED IN TROPICAL COUNTRIES

BY

B. S. PLATT, C.M.G., M.B., M.Sc., Ph.D.

(from the Human Nutrition Research Unit of the Medical Research Council,
National Hospital, Queen Square, London)

1. INTRODUCTION

A widespread demand exists for tables of food composition designed for use in tropical countries. In the tables given here, values have been selected after scrutiny of an extensive field of literature, including analyses done on samples collected by the Nyasaland Nutrition Survey (p.3), and other unpublished data.

An attempt has been made to assess the weight to be attached to the various analyses, and to arrive at representative values for a suitable range of foodstuffs. These values are considered to be sufficiently well founded for the purpose for which they have been compiled, namely, the evaluation of dietary data based on records of group intake. They are not suitable for surveys by the individual method, nor for precise nutritional investigations on individual human and animal subjects. Many of the values are at present only tentative or provisional, and revision will be undertaken from time to time as further data become available. Any communications or suggestions for new values will be welcomed by the Human Nutrition Research Unit.

2. Food Categories

In classifying the foodstuffs, two considerations have been kept in mind —firstly, the type of dietary commonly found in tropical countries and secondly, the nutritive value of the foods. They have been grouped as follows :—

1. Cereals and Starchy Roots, Fruits, etc.
 - (a) Cereals
 - (b) Starchy Roots, Fruits, etc.
2. Legumes
3. Oilseeds and Nuts
4. Vegetables
 - (a) Leafy, Green and Yellow
 - (b) Other

Food Categories—contd.

5. Fungi and Yeasts
6. Fruits
7. Stems
8. Fats and Oils
9. Animal Products (except Fats)
 - (a) Fish and Fish Products (including Molluscs and Crustacea)
 - (b) Insects
 - (c) Meat, Meat Products and Eggs
 - (d) Milk and Milk Products
10. Condiments and Spices
11. Syrups, Sugars and Preserves
12. Beverages.

3. Explanations and Definitions

(a) *Carbohydrate* values are conventionally expressed either as available carbohydrate by direct chemical analysis (AC) or as carbohydrate estimated arithmetically by difference (CD). For certain foodstuffs these values are similar and in some cases identical, but in others there are considerable differences. The values for carbohydrates (C) given here represent a compromise between the two, arrived at by assuming that, after deducting the fibre (F), one-half the carbohydrate, not allowed for in the chemical analysis but included in the total carbohydrates estimated by difference, is utilised in the body. This may be expressed in the formula $C = \frac{1}{2} (AC + CD - F)$. Where values for total carbohydrate only could be found, this figure (excluding fibre) has been given in italics and the calorie value is then shown in brackets.

(b) *Calories*—The factors 4, 9 and 4 Calories per gram of protein, fat and carbohydrate respectively have been employed in computing calorie values. The values obtained are therefore only approximations, but their order of accuracy is sufficient for the purpose described above.

(c) *Minerals*—The figures are for total mineral content and take no account of availability.

(d) *Vitamins*—Vitamin values of any given food may vary widely and are affected by such factors as variety, maturity, storage and processing, as well as locality and, in the case of certain animal products, the feeding of the animal. Where the range is so wide that the value given in the tables cannot be regarded as representative, fuller information is given in a footnote.

Vitamin A values are expressed as International Units (I.U.) of vitamin A, either as vitamin A itself or as the equivalent of β -carotene, denoted by the symbol “-c” (1 I.U. = 0.6 mcg. β -carotene). In computing requirements and intake, 1 I.U. of vitamin A may for the present be taken as the equivalent of 3 I.U. in the form of a precursor. It should be noted that the allowance of 5,000 I.U. recommended by the National Research Council (U.S.A.) is based on the assumption that 40 per cent. is derived from vitamin A and 60 per cent. from precursors. It is thus the equivalent of 3,000 I.U. in terms of vitamin A only. The ratio in biological value of vitamin A to its precursor, carotene, varies, and when knowledge permits, separate factors may have to be used for different foodstuffs.

Vitamin B₁ (aneurin, thiamine) is shown in mg., the relation to International Units being 1 I.U. = 3 mcg.

(e) Abbreviations.

0 = contains none

§ = contains so small a quantity, if any, that it can be neglected in the dietary evaluation

* = average nutrient factors which should be used if there is no detailed information on the type of the commodity

[]_x = taken as Item No. x

() = tentative or provisional value

-c = I.U. vitamin A as carotene

w = white varieties

y = yellow varieties.

4. "Edible Portion" and "As Purchased"

All values refer to the portion usually eaten (E.P.) and, where possible, a waste figure has been shown as a percentage of the food as purchased (A.P.). The proportion of waste may vary widely, e.g. in mangoes from 10 to 52 per cent. according to the size of seed, amount of fibre, etc.; or, in fish, according not only to size and species but to the consumer's method of handling it. In a survey it is therefore advisable to determine the waste values locally for at least the important foods. Where dietary records are kept on an A.P. basis, the nutrient values to be applied may be calculated from the E.P. figures as follows:—

$$\text{A.P. value} = \text{E.P. value} \times \frac{(100 - \text{Waste percentage})}{100}$$

5. Importance of Water Content

Many tables of food composition omit values for water content. This varies considerably from one sample to another, affecting the values for all nutrients, and scrutiny of the literature reveals that many apparent divergencies are eliminated when allowance is made for differences in water content. The effect of this factor is, of course, most marked in foods of high water content, such as fruits and fresh vegetables. Where necessary, the nutrient values shown here should be adjusted to a value for water determined independently.

6. Importance of Accurate Evaluation of Staple Foods

In the case of foods consumed in large quantities, and supplying a substantial proportion of the total diet, e.g. rice, maize, cassava (depending on the food habits of the community), quite small differences in composition may give rise to very large differences in the final results of a survey and its estimate of nutrient intake. If, therefore, there is any reason to doubt the values given for a community's staple foodstuff, special analyses should be done.

7. Plant "Pot-ashes", Edible Earths and Drinking Water

In some localities, native "salt" made from the ashes of marsh or other plants is used. Such "pot-ashes" may contribute substantial amounts of minerals to the dietary, and so may edible earths and drinking water. No representative values can, however, be given for any of these items, because the range of variation is too great; special analyses would be required in each case.

8. Use of Common and Botanical Names

Where several common names are given for one item, they are not necessarily strictly alternatives but may refer to several species taken as a group for the purpose of these tables, e.g. taro, eddoes, cocoyams, yautia. Only the most widely used common names are given, but with the help of the botanical names it is possible to consult H. C. Sampson's monograph, *Cultivated Crops of the British Empire and the Anglo-Egyptian Sudan* (Sampson, 1936), in which many additional local names will be found. He gives, for example, as many as twenty names for egg-plant.

Alternative botanical names given by Sampson are shown at the end of the tables (p. 35), followed by a list of the literature consulted.

9. Acknowledgments

Acknowledgments for help in regard to the collection and presentation of these data are due to members of the Human Nutrition Research Unit (1944-5) and of the Nyasaland Nutrition Survey (1938-9); and to Mrs. G. M. Culwick and Miss M. W. Grant, B.Sc.

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
	L.—CEREALS and STARCHY ROOTS, FRUITS, Etc.		6-	6-		6-
	A. Cereals					
1	Buckwheat, flour	Fagopyrum esculentum Moench.	—	14	(337)	6
2	Job's tears (adlay), seed ..	Coix lachryma-jobi Linn.	—	23	(304)	11
3	Maize (Indian corn) Whole, white	Zea mays Linn.	6-12	12	349	10
4	Whole, yellow		6-12	12	349	10
5	Meal*, about 96 per cent. extraction		—	12	342	9.5
6	Meal*, highly refined, about 60 per cent. extraction		11-15	12	337	8
7	Hominy (samp)		—	11	(353)	8.5
8	Maizestarch (cornflour), commercial		—	12	352	0.5
9	Germ meal—germ, some endosperm and fine bran		24-27	25(b)	249	8
10	Immature (green mealies, sweetcorn), on cob in- cluding sheath		37-84	80(e)	(79)	3
11	Millets Bulrush, seed	Pennisetum typhoides Stapf & Hubbard	9-14	12	(365)	11
12	Finger, seed	Eleusine coracana Gaertn.	10-14	12	(338)	6
13	Various (broomcorn, China, common, French, hog, Italian, little, proso)	{ Panicum spp. (cultivated) Setaria spp. (cultivated)}	8-16	12	(336)	10
14	Oatmeal, 65 per cent. extrac- tion Paste—see under Wheat	Avena sativa Linn.	—	9	386	12
15	Rice Home pounded or lightly milled.	Oryza sativa Linn.	10-16	12	354	8
16	Milled, i.e. highly milled ..		11-13	12	349	7
17	Parboiled, milled		11-13	12	352	7.5
18	Rye, 85-90 per cent. extrac- tion	Secale cereale Linn.	—	12	327	8
19	Sorghum (guineacorn, kaffir- corn) Whole seed	Sorghum spp. (cultivated)	8-14	12	(356)	10.4
20	"Straight run" meal		—	12	(352)	9.6
21	Table meal, bran removed		—	12	(356)	9.5

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Aneurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
1	76	10	1	s	0.13	0.11	(4)	0	0	1
4	56	200	10	—	0.14	—	—	—	0	2
4.5	67	12	5	{ w 100-c	0.33	0.13	1.5	0	—(a)	3
4.5	67	12	5	{ s 100-c	0.33	0.13	1.5	0	—(a)	4
4	67	12	5	{ y 100-c	0.3	0.13	1.5	0	0	5
0.5	75	9	2	{ w 100-c	0.05	0.03	0.6	0	0	6
0.6	78.5	11	4	{ y 100-c	0.05	0.02	0.2	0	0	7
0.2	87	15	1.4	0	0	0	0	0	0	8
5	43(c)	15	6	{ w 100-c	0.6	0.26	3.0	0	—(d)	9
0.8	15	9	0.5	{ w 250-c	0.15	0.05	1.7	10	60(e)	10
5	69	(25)	(6)	s	0.50	—	3.2	0	0	11
2	74	350	5	s	0.28	0.05	1.4	0	0	12
4	65	20	5	s	0.6	—	(0.3)	0	0	13
8.7	65	60	5	s	0.50	0.14	1.3	0	0	14
2	76	10	2	s	0.24	0.1	2	0	0	15
1	78	5	0.5	0	0.06	0.05	1	0	0	16
1.5	77	8	1	s	0.21	0.1	4	0	0	17
2.5	68	50	3.5	s	0.27	0.10	1.2	0	0	18
3.4	70.9	32	6.6	s	0.50(f)	0.12	3.5	0	0	19
2.7	72.2	34	(6.6)	s	0.47(f)	0.10	3.0	0	0	20
2.7	73.3	36	8.7	s	0.44(f)	0.08	2.6	0	0	21

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
	L.—CEREALS and STARCHY ROOTS, FRUITS, Etc.—contd.					
	A. Cereals—contd.					
	Wheat	Triticum vulgare Host Cultigen.				
22	Whole*	—	13	316	11.5(g)
23	Flour, 85 per cent. extraction	—	13	346	11
24	Flour, 70 per cent. extraction (white)	—	13	351	10
25	Bread	9-11			
26	Paste (macaroni, noodles, spaghetti)				

Additional Names

- Adlay, *see* Job's tears
 Cornflour, *see* Maizestarch
 Green mealies, *see* Maize, immature
 Guineacorn, *see* Sorghum
 Indian corn, *see* Maize
 Kaffircorn, *see* Sorghum
 Macaroni, *see* Paste
 Millet, pearl, *see* Millet, bulrush
 Noodles, *see* Paste
 Samp, *see* Maize, hominy
 Spaghetti, *see* Paste
 Sweetcorn, *see* Maize, immature.

For significance of asterisks and other abbreviations used throughout Tables, see p. 6.

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Aneurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
2.4	62	30	3.5	0	0.40	0.17	5	0	0	22
1.6	72	20	2.5	0	0.30 ^(h)	0.10	1.5 ⁽ⁱ⁾	0	0	23
1	75.5	16	2	0	0.06	0.05	1	0	0	24
Take as 72 per cent. of corresponding flour										
Take as corresponding flour corrected for water content										
										25
										26

Footnotes

- (a) For ripe maize on cob, with sheath, waste = 25 per cent.
 (b) Water content is relatively high because the grain is moistened to facilitate pounding and this fraction is usually eaten immediately without re-drying.
 (c) "Available" carbohydrate. The data are insufficient for the formula described in the Introduction to be applied.
 (d) This fraction represents approximately 5 per cent. of the original grain.
 (e) Water content and percentage of waste decrease with maturity.
 (f) Value given for aneurin in whole sorghum is from unpublished data, and the values for sorghum meals have been calculated from this on the basis of the loss shown by Fox & Golberg, *South African Food Tables*. Wide variation found.
 (g) Protein may vary from 8-15 per cent., but calorie value remains the same: U.K. 11.5, Manitoba 13.9, Australian 10-15, S. African 13.6.
 (h) Atta, various grades, 0.4-0.6 mg.
 (i) Atta, various grades, 4.0-6.5 mg.

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
	I.—CEREALS and STARCHY ROOTS, FRUITS, Etc.—contd.		g.	g.		g.
	B. Starchy Roots, Fruits, etc.					
27	Arrowroot	Maranta arundinacea Linn.	11-17	15	(337)	0·2
28	Banana (a)	Musa sapientum Linn.	61-84	70	103	1
29	Breadfruit, pulp (a)	Artocarpus communis Forst.	66-74	70	(109)	1·5
	Cassava (manioc)	Manihot utilissima Pohl.				
30	Flour	11-13	12	314	1·5
31	Fresh tuber	49-74	60	131	0·7
32	Tapioca	—	11	346	0·5
33	Jak fruit, pulp (a)	Artocarpus integer (Thunb.) Merr.	—	86	47	2·5
34	Plantain, green (a)	Musa paradisiaca Linn.	61-84	70	103	1
	Potato Irish	Solanum tuberosum Linn.	75-81	78	82	2
36	Potato Sweet	Ipomoea batatas (Linn.) Lam.	58-79	70	121	2
37	Sago, flour	Metroxylon sago Rottb.	—	12	(341)	0·3
38	Taro (coco, cocoyam, dasheen, eddo, tannia, yautia)	Colocasia, Alocasia & Xanthosoma spp.	56-83	75	88	2
39	Yam	Dioscorea spp.	62-80	73	100	2

Additional Names

Coco, *see* Taro
 Cocoyam, *see* Taro
 Dasheen, *see* Taro
 Eddo, *see* Taro
 Manioc, *see* Cassava
 Tannia, *see* Taro
 Yautia, *see* Taro.

For significance of asterisks and other abbreviations used throughout Tables, *see* p. 6.

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Aneurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
ø	84	7	1	0	0	(0)	(0)	0	0	27
0·3	24	7	0·5	350-c ^(b)	0·05	0·06	0·5	10	33	28
0·3	(25)	25	1	35-c	(0·2)	—	—	(15)	—	29
ø	77	55	4	0	ø	(0)	[(1·2)] _{ss}	0	—(c)	30
ø	32	25	1	0	0·02	0·1	—	30	15(c)	31
ø	86	10	1	0	0	(0)	(1·2)	0	0	32
0·3	8·5	20	0·5	400-c	[(0·2)] _{ss}	—	0·4	10	—	33
0·3	24	7	0·5	100-c	[0·05] _{ss}	[0·06] _{ss}	[0·5] _{ss}	[10] _{ss}	[33] _{ss}	34
ø	18·4	10	0·7	35-c	0·1	0·05	1	15	15	35
1	26	20	0·7	{ w ø y 4,000- 7,000-c }	0·1	0·06	0·7	20	15	36
ø	85	10	1	0	0	(0)	(0·2)	0	0	37
ø	20	25	1	40-c	(0·05)	[0·06] _{ss}	0·4	ø	20	38
ø	23	10	1·2	—	0·05	[0·06] _{ss}	[0·4] _{ss}	ø	[15] _{ss} ^(d)	39

Footnotes

(a) Flours derived from breadfruit, bananas, etc., should be calculated from values for fresh foodstuff, bearing in mind that water-soluble factors may be washed out and accessory food factors like vitamin C may be lost.

(b) 30-700 I.U. (carotene) depending on variety.

(c) Thin outer peel only, removed after cooking, waste = 5 per cent.

(d) For large varieties, waste may be as low as 5 per cent.

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
II.—LEGUMES						
40	Bambara groundnut (earth pea, jugo bean)	<i>Voandzeia subterranea</i> Thou.	8-13	10	346	18
41	Bonavist bean (dolichos bean, field bean, hyacinth bean)	<i>Dolichos lablab</i> Linn.	7-10	8	325	24
42	Carob (locust bean), seed ..	<i>Ceratonia siliqua</i> Linn.	8-12	10	291	16
43	Chickpea (Bengal gram) ..	<i>Cicer arietinum</i> Linn.	8-10	9	353	22
44	Cowpea	<i>Vigna unguiculata</i> (Linn.) Walp.	8-14	10	281	24
45	Goa bean (asparagus pea, winged bean), seed	<i>Psophocarpus tetragonolobus</i> DC.	—	14	396	33
	Groundnut (monkey nut, peanut)	<i>Arachis hypogaea</i> Linn.				
46	Dry	4-8	6	555	23
47	Fresh (b)	38-46	45	(328)	(13)
48	Horse bean (broad bean) ..	<i>Vicia faba</i> Linn. ..	—	8	(338)	31
49	Horse gram	<i>Dolichos biflorus</i> Linn. ..	9-12	10	293	22
50	Kidney bean (French, haricot, navy, pinto, snap)	<i>Phaseolus vulgaris</i> Linn.	8-15	10	306	24
51	Lentil (red dhall), split pea	<i>Lens esculenta</i> Moench.	6-12	10	305	24
52	Lima bean (butter bean, curry bean, Madagascar bean)	<i>Phaseolus lunatus</i> Linn.	8-14	12	305	20
53	Locust bean (<i>not</i> Carob), seed	<i>Parkia</i> spp. ..	9-14	12	243	15
54	Mung bean (black gram, woolly pyrol)	<i>Phaseolus mungo</i> Linn.	10-14	11	317	24
55	Mung bean (green gram) ..	<i>Phaseolus aureus</i> Roxb.	10-12	12	309	22
56	Pea, garden or field	<i>Pisum sativum</i> Linn.	7-22	10	309	22
57	Pigeon pea (Congo pea, red gram, yellow dhall)	<i>Cajanus cajan</i> Linn.	7-15	10	306	22
	Soya bean	<i>Glycine max</i> (Linn.) Merr.				
58	Curd	—	84	81	7.5
59	Milk	—	93	35	3.4
60	Sauce	—	72	(55)	5.9
61	Seed, edible varieties	6-13	7	382	35
62	Sword bean (Jack bean), kernel	<i>Canavalia ensiformis</i> DC.	—	8	391	38

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Aneurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
6	55	90	4	—	0.30	—	—	0	0	40
1	55	60	2	—	0.18	—	1.8	0	0	41
3	50	—	—	—	—	—	—	—	—	42
5	55 1	200 44	9 4	300-c 50-c	(0.35) 0.89	(0.15) 0.14	(1.4) 1.3	12 0	0 0(a)	43 44
16	30	—	—	—	0.03	—	—	0	—	45
47 (28)	10 (6)	50 (29)	2 (1)	s s	0.9 (0.5)	0.14 (0.08)	17.4 (10.2)	0 10	30 35	46 47
1.5	50	95	6	—	—	—	—	—	—	48
0.5	50	280	8	100-c	—	—	—	0	—	49
2	48	110	8	0	0.45	0.34	2.3	0	0a	50
1	50	40	7	200-c	0.65	0.18	2.8	0	0	51
1	54	70	6	s	0.53	0.14	1.4	0	0	52
7	30 1	200 53	9	[0]so	0.42	[0.34]so	[2.3]so	0	0	53 54
1	53	140	8	[0]so	0.46	[0.34]so	[2.3]so	0	0	55
1 2	53 50	60 140	4 4	200-c [0]so	0.45	0.18 [0.34]so	2.8 [2.3]so	0 0	0 0	56 57
4.2	3.4	34(c)	1.8	—	—	—	—	—	—	58
1.5	2.0	12	—	—	—	—	—	—	—	59
1.2	5.2	59	4.9	s(d)	1.12	0.32	2.1	0	0	60 61
18 3	20 53	220 100	7 —	—	0.15	—	—	0	(10)(e)	62

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
	II.—LEGUMES—contd.		g.	g.		
63	Tepary bean	Phaseolus acutifolius A.	—	10	341	24
64	Velvèt bean	Mucuna utilis hort.	9-15	10	231	24

Additional Names

Asparagus pea, *see* Goa bean
 Bengal gram, *see* Chickpea
 Broad bean, *see* Horse bean
 Butter bean, *see* Lima bean
 Congo pea, *see* Pigeon pea
 Curry bean, *see* Lima bean
 Dolichos bean, *see* Bonavist bean
 Dhall, red, *see* Lentil
 Dhall, yellow, *see* Pigeon pea
 Earth pea, *see* Bambara groundnut
 Field bean, *see* Bonavist bean
 French bean, *see* Kidney bean
 Gram, red, *see* Pigeon pea
 Gram, black, *see* Mung bean
 Gram, green, *see* Mung bean
 Haricot bean, *see* Kidney bean
 Hyacinth bean, *see* Bonavist bean
 Jack bean, *see* Sword bean
 Jugo bean, *see* Bambara groundnut
 Locust bean, *see* Carob, also Item 53
 Madagascar bean, *see* Lima bean
 Monkey nut, *see* Groundnut
 Navy bean, *see* Kidney bean
 Peanut, *see* Groundnut
 Pinto bean, *see* Kidney bean
 Snap bean, *see* Kidney bean
 Winged bean, *see* Goa bean
 Woolly pyrol, *see* Mung bean.

LEGUMES, FRESH

See under Vegetables, leafy,
 green and yellow, also Item
 47 above.

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Aneurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
1	59	—	—	—	—	—	—	0	—	63
3	27	130	—	50-c	0.3	—	—	0	—	64

Footnotes

- (a) Seed coat is sometimes removed, averaging 10 per cent.
- (b) Nutrient values (except for vitamin C) are provisional only, calculated from values for dry nuts.
- (c) Depends on agent used for curdling the soya milk and may be as high as 270 mg.
- (d) 0-400 I.U. (carotene), depending on variety.
- (e) Taken as seed coat of kidney bean.

For significance of asterisks and other abbreviations used throughout Tables, see p. 6.

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
	III.—OIL SEEDS and NUTS		g.	g.		g.
65	Arecanut	<i>Areca catechu</i> Linn.	—	31	(248)	4.9
66	Avocado pear, flesh	<i>Persea americana</i> Mill.	67-82	72	(208)	2
67	Cashew nut	<i>Anacardium occidentale</i> Linn.	4-6	5	571	20
	Coconut	<i>Cocos nucifera</i> Linn.				
68	Kernel, ripe fresh	20-52	43	404	4
69	Milk, ripe nut	—	92	22	0.5
70	Dika nut, kernel, sun-dried ..	<i>Irvingia gabonensis</i> Baill.	—	—	—	—
71	Goa bean (asparagus pea, winged bean), seed	<i>Psophocarpus tetragonolobus</i> DC.	—	14	396	33
72	Groundnut (monkey nut, peanut), dry	<i>Arachis hypogaea</i> Linn.	4-8	6	555	23
73	Kola nut	<i>Cola acuminata</i> Schott & Endl.	—	12	(309)	9
74	Oilbean, kernel	<i>Pentaclethra macrophylla</i> Bth.	—	10	(561)	22
75	Olive, flesh	<i>Olea europaea</i> Linn.	—	30	(528)	5
76	Oyster nut, kernel	<i>Telfairia</i> spp.	—	—	—	—
77	Pumpkin, seed, kernel	<i>Cucurbita</i> spp.	—	3	(594)	26
78	Sesame (beniseed, gingelly, simsim), seed	<i>Sesamum indicum</i> Linn.	4-8	5	570	20
79	Shea nut, kernel	<i>Butyrospermum parkii</i> Kotschy	—	(7)	(545)	(9)
80	Soya bean, seed, edible varieties	<i>Glycine max</i> (Linn.) Merr.	6-13	7	382	35
81	Sunflower, seed, kernel ..	<i>Helianthus annuus</i> Linn.	4-8	6	(569)	27

Additional Names

- Asparagus pea, *see* Goa bean
 Beniseed, *see* Sesame
 Fluted pumpkin, *see* Oyster nut
 Gingelly, *see* Sesame
 Monkey nut, *see* Groundnut
 Peanut, *see* Groundnut
 Simsim, *see* Sesame
 Winged bean, *see* Goa bean.

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Anceurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
4·4 20	47 5	50 10	1·5 1	— 200-c ^(a)	0·1	0·15	1·0	20 ^(b)	— 30	65 66
45	21·5	50	5	—	—	—	—	0	—	67
40 —	7 5	10 30	2 —	0 0	0·11 0	—	(0·4) —	1 2	35 ^(c) — ^(c)	68 69
54	—	—	—	—	—	—	—	—	—	70
16	30	—	—	—	0·03	—	—	0	—	71
47	10	50	2	— —	0·9	0·14	17·4	0	30	72
1	66	—	—	—	—	—	—	—	—	73
45	17	90	—	—	—	—	—	—	—	74
52 48 50 50 45	(10) — 10 10 (26)	100 — 60 1,500	10 — 4 15 —	(500-c) — — — —	— — — 0·42 —	— — — — —	— — — — —	— — — 0 —	25 — 25 — —	75 76 77 78 79
18	20	220	7	— ^(d)	1·12	0·32	2·1	— — — — —	0 — — — 50	80 81
45	14	100	7	—	—	—	—	—	—	

Footnotes

(a) 100-850 I.U. (carotene) depending on variety.

(b) 7-37 mg. depending on variety.

(c) For kernel and milk together, waste = about 20 per cent., i.e. kernel averages 65 per cent. and milk 15 per cent. of ripe nut as purchased; by "milk" is meant fluid in coconut, and not a preparation made from grated coconut flesh.

(d) 0-400 I.U. (carotene) depending on variety.

For significance of asterisks and other abbreviations used throughout Tables, see p. 6.

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
	IV.—VEGETABLES		g.	g.		g.
82	A. Leafy, Green and Yellow Beans, all, eaten green in pod		78-94	90	30	2
83	Beans and peas, all, eaten green, shelled		70-80	76	73	6.5
84	Betel leaves	Piper betle Linn.	—	85	(36)	3
85	Carrots, without tops, med. maturity	Daucus carota Linn.	86-95	90	30	0.9
86	Leaves, fresh High carotene—dark green, e.g. leaves of amaranths, beet, cassava, cowpea, kale, mustard, pumpkin, spinach, sweet potato, turnip, watercress	65-95	85	44	4
87	Low carotene—pale green, e.g. brussels sprout, cabbage	65-95	90	26	3
88	Lettuce, mainly green	—	94	14	1.4
89	Leaves, sun-dried, high caro- tene (c)	7-10	10	(264)	(24)
90	Leek	Allium porrum Linn. and A. ampelo- prasum Linn.	85-92	88	25	1.0
91	Okra (Ladies' finger) ..	Hibiscus esculentus Linn.	80-94	90	29	1.8
92	Onion, spring	Allium fistulosum Linn.	—	88	33	1.8
93	Pepper, sweet (sweet chilli, paprika), fresh Green	Capsicum annuum Linn.	82-93	88	37	2
94	Red	82-93	88	37	2
95	Pepper, sweet, sun-dried (k) Green	10-14	12	(268)	(15)
96	Red	10-14	12	(268)	(15)
97	Persimmon leaves Fresh	Diospyros kaki Linn.	—	—	—	—
98	Dried	—	—	—	—
99	Pumpkin (squash), yellow ..	Cucurbita spp.	79-95	90	28	1

Additional Names

- Chilli, sweet, *see* Pepper, sweet
 Ladies' finger, *see* Okra
 Paprika, *see* Pepper, sweet
 Squash, *see* Pumpkin.

For significance of asterisks and other abbreviations used throughout Tables, see p. 6.

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Aneurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
g	5.5	64	1.4	600-c	0.08	0.12	0.5	20	10	82
g	11.8	40	2	400-c	0.20	0.15	0.5	25	55	83
g	6	150 ^(a)	5.7	9,500-c	—	0.70	5	—	—	84
g	6.7	40	0.7	13,000-c	0.07	0.06	0.56	6	10	85
0.4	6	210	3	13,000-c	0.15	0.25	0.85	100	20 ^(e)	86
0.2	3	85	1	500-c ^(b)	0.10	0.10	0.35	50	— ^(e)	87
g (2.4)	2 (36)	35 (1,260)	0.7 (18)	2,000-c ^(c) (7,800-c) ^(d)	0.05 —	0.04	0.18	7 (42)	—	88 89
g	5.3	50	1	700-c	0.03	(0.1)	(0.1)	20	48	90
g	5.5	70	1	1,000-c	0.1	0.1	0.7	25	12	91
g	6.5	40	0.9	3,000-c	0.05	(0.12)	(0.11)	20	48	92
0.5 0.5	6 6	20 20	1 1	1,000-c ^(f) 18,000-c ⁽ⁱ⁾	0.06 [0.06] _{ss}	0.04 [0.04] _{ss}	0.40 [0.40] _{ss}	200 ^(g) 350 ^(j)	16 ^(h) 20 ^(h)	93 94
(3.5) (3.5)	(44) (44)	(150) (150)	(7) (7)	730-c ^(f) 13,000-c ⁽ⁱ⁾	— —	— —	— —	100 ^(j) 180 ^(j)	— —	95 96
—	—	—	—	—	—	—	—	2,726 3,585 ^(m)	—	97
g	6	20	0.8	1,100-c	0.04	0.05	0.7	5	25	98 99

Footnotes

(a) Chewed with lime. Amount and utilisation of calcium ingested with 6 betel leaves similar to 10 oz. cow's milk.

(b) For white cabbage, take vitamin A = 50 I.U. (carotene).

(c) Nutrient values are provisional only, calculated from values for fresh leaves, and assuming vitamin losses as for sweet peppers, see note (i).

(d) On this basis, value for low-carotene group would be 450 I.U. and for white cabbage 45 I.U.

(e) For 16 kinds of leaves used in Nyasaland, waste varied from 0-60 per cent.

(f) 200-5,200 I.U. (carotene), depending on variety.

(g) 100-300 mg., depending on variety.

(h) This figure includes seeds and cores as well as stem ends.

(i) 5,700-41,000 I.U. (carotene), depending on variety.

(j) 250-560 mg., depending on variety.

(k) Values for protein, fat, carbohydrate and minerals are provisional only, calculated from values for fresh peppers.

(l) Allows for 90 per cent. loss of vitamin A and 93-per cent. loss of vitamin C during drying. Further losses would be sustained during storage. See Adams & Smith, *The vitamin content and the preservation of foods*, p. 50.

(m) Even the tea made from dried persimmon leaves is stated to be a potent source of vitamin C.

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
	IV.—VEGETABLES—contd. B. Other					
100	Beetroot, without tops ..	<i>Beta vulgaris</i> Linn.	82-94	87	41	1.8
101	Cucumber, various ..	<i>Cucumis sativus</i> Linn.	91-97	96	10	0.6
102	Egg plant (aubergine, brinjal, garden egg)	<i>Solanum melongena</i> Linn.	89-94	93	20	1
103	Gourd	<i>Lagenaria vulgaris</i> Ser.	86-96	90	26	0.5
104	Leek	<i>Allium porrum</i> Linn. and <i>A. ampeloprasum</i> Linn.	85-92	88	25	1.0
105	Lotus root	<i>Nelumbium nucifera</i> Gaerth.	83-96	96	(12)	0.4
106	Maize, immature (green mealies, sweetcorn), on cob including sheath	<i>Zea mays</i> Linn. ..	37-84	80 ^(a)	(79)	3
107	Onion and shallot	<i>Allium cepa</i> Linn. & <i>A. ascolonicum</i> Linn.	70-95	87	37	1
108	Onion, spring	<i>Allium fistulosum</i> Linn.	—	88	33	1.8
109	Radish	<i>Raphanus sativus</i> Linn.	—	94	16	1.2
110	Tomato, with skin	<i>Lycopersicum esculentum</i> Mill.	90-96	92	18	1
111	Turnip and swede	<i>Brassica</i> spp. ..	—	87	36	1.2
112	Water calthrop Large	<i>Trapa natans</i> L.	—	45	(214)	5
113	Red (Jesuits' nut)	—	81	(68)	3
114	Water chestnut	<i>Scirpus tuberosus</i> Roxb.	79-85	85	(54)	0.8

Additional Names

Aubergine, *see* Egg plant
 Brinjal, *see* Egg plant
 Garden egg, *see* Egg plant
 Jesuits' nut, *see* Water calthrop, red
 Mealies, green, *see* Maize, immature.

V.—FUNGI and YEASTS							
	Fungi, mixed						
115	Fresh	76-93	90	7	1	
116	Dried	9-15	10	73	10	
	Mushrooms*	<i>Agaricus</i> and <i>Morchella</i> spp.					
117	Fresh	—	91	—	—	
118	Dried	—	12	—	—	
	Yeast						
119	Bakers'	<i>Saccharomyces cerevisiae</i>	—	5	—	45	
120	Brewers'	—	5	—	50	
121	Food	<i>Torula utilis</i> ..	—	8	206	43.1	

For significance of asterisks and other abbreviations used throughout Tables, see p. 6.

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Aneurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
0	8.5	30	1	0	0.03	0.04	0.4	10	25	100
0	1.8	10	0.3	200-c	0.04	0.05	0.18	10	30	101
0	4	15	1	0	0.05	0.03	0.79	5	13	102
0	6	20	0.8	0	[0.04] _{ss}	[0.05] _{ss}	[0.7] _{ss}	[5] _{ss}	[25] _{ss}	103
0	5.3	50	1	700-c	0.03	(0.1)	(0.1)	20	48	104
0	2.6	35	1.4	—	—	—	—	—	6	105
0.8	15	9	0.5	{w y 250-c}	0.15	0.05	1.7	10	60 ^(a)	106
0	8.2	30	0.4	250-c	0.03	0.1	0.1	15	5	107
0	6.5	40	0.9	3,000-c	0.05	(0.12)	(0.11)	20	48	108
0	2.8	38	1.6	0	0.05	0.03	0.15	25	39	109
0	3.5	9	0.4	1,200-c	0.05	0.04	0.7	25	2	110
0	7.8	57	0.5	0	0.03	0.06	0.7	34	15	111
0.7	47	36	1.6	—	—	—	—	—	57	112
0	14	20	—	—	—	—	—	—	55	113
0	12.7	4	0.8	—	—	—	0.6	—	26	114

Footnote

(a) Water content and percentage of waste decrease with maturity.

0.3	0	5	1.2	0	[0.12] ₁₁₇ [(1.2)] ₁₁₈	[0.52] ₁₁₇ 0	[5.8] ₁₁₇ —	[9] ₁₁₇ 0	[9] ₁₁₇ —	115 116
0.3	0	14	0.7	0	0.12 (1.2)	0.52 0	5.8 —	9 0	9 0	117 118
2	—(a)	40	25	0	3	7	30-40	0	0	119
1	—(a)	80	20	0	16	4	30-40	0	0	120
2.4	3.0(b)	127	20	0	2	6	30-40	0	0	121

Footnotes

(a) The only figures available are for total carbohydrate, including cellulose.

(b) "Available" carbohydrate in terms of starch.

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
VI.—FRUITS						
122	Avocado pear, flesh	<i>Persea americana</i> Mill.	67-82	72	(208)	2
123	Banana	<i>Musa sapientum</i> Linn.	61-84	70	103	1
124	Breadfruit, pulp	<i>Artocarpus communis</i> Forst.	66-74	70	(109)	1.5
125	Cape gooseberry (ground cherry)	<i>Physalis peruviana</i> Linn.	—	83	55	2
126	Citrus Lime	<i>Citrus aurantiifolia</i> (Christm.) Swingle	89-91	90	31	0.8
127	Orange and tangerine (mandarin orange, naartje)	<i>Citrus sinensis</i> Osbeck and C. <i>nobilis</i> Lour.	83-90	86	43	0.8
128	Other	<i>Citrus</i> spp.	86-94	89	30	0.5
129	Custard apple	<i>Annona reticulata</i> Linn.	—	74	(81)	2
130	Dates, dried	<i>Phoenix dactylifera</i> Linn.	15-23	20	284	2
Fig						
131	Fresh	<i>Ficus carica</i> Linn.	50-88	78	76	1.4
132	Dried	17-24	20	269	4
133	Grape	<i>Vitis vinifera</i> Linn.	72-88	82	64	1
134	Grenadilla (passion fruit) ..	<i>Passiflora edulis</i> Sims.	—	72	(94)	2.3
135	Guava	<i>Psidium guajava</i> Linn.	75-85	80	36	1
136	Indian gooseberry	<i>Phyllanthus emblica</i> Linn.	—	—	—	—
137	Jak fruit, pulp	<i>Artocarpus integer</i> (Thunb.) Merr.	—	86	47	2.5
138	Mango, ripe	<i>Mangifera indica</i> Linn.	74-90	83	59	0.7
Melon						
139	Sweet (cantaloupe, honeydew, muskmelon).	<i>Cucumis melo</i> Linn.	—	96	(10)	0.5
140	Water	<i>Citrullus vulgaris</i> Schrad.	—	93	26	0.5
141	Orange—see under Citrus					
142	Papaya (pawpaw)	<i>Carica papaya</i> Linn.	85-94	89	38	0.6
	Peach, flesh	<i>Prunus persica</i> Stokes	82-90	87	42	0.5

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Aneurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
20	5	10	1	200-c ^(a)	0.1	0.15	1.0	20 ^(b)	30	122
0.3	24	7	0.5	350-c ^(c)	0.05	0.06	0.5	10	33	123
0.3	(25)	25	1	35-c	(0.2)	—	—	(15)	—	124
0.8	10	10	1.8	5,000-c	0.01	—	—	30	6	125
—	—	—	—	—	—	—	—	—	—	—
ø	7	22	0.6	ø	(0.02)	ø	0.2	25	23	126
ø	10	30	0.3	250-c	0.08	0.03	0.2	45	25	127
ø	7	20	0.5	ø	0.04	ø	0.2	40	35	128
1	16	26	1	—	—	—	—	15	26	129
ø	69	70	2	100-c	0.07	0.03	2.2	0	13	130
—	—	—	—	—	—	—	—	—	—	—
ø	17.5	50	1	80-c	0.05	(0.05)	0.6	ø	0	131
1	61	200	4	100-c	0.2	—	1.7	ø	0	132
ø	15	20	0.3	50-c	0.03	0.01	0.3	ø	5	133
3.3	13.7	14	1.7	200-c	—	—	—	20	43 ^(d)	134
ø	8	15	1	200-c	0.06	0.03	0.5	200 ^(e)	20 ^(f)	135
—	—	—	—	—	—	—	—	600	—	136
0.3	8.5	20	0.5	400-c	[0.2] ¹³⁴	—	0.4	10	—	137
ø	14	10	0.5	1,000-c ^(g)	0.06	0.05	0.2	30 ^(h)	34	138
ø	2	6	0.3	{ w ø y 2,400-c ⁽ⁱ⁾ }	0.05	0.05	0.05	35 ^(j)	20	139
ø	6	6	0.2	500-c	0.05	0.07	0.05	ø	50	140
ø	9	10	0.5	2,500-c	0.02	0.02	0.1	60 ^(k)	30	141
ø	10	8	0.4	{ w ø y 1,200 }	0.02	0.04	0.5	8	12	142

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
VI.—FRUITS—contd.						
143	Persimmon	<i>Diospyros kaki</i> Linn.	78-80	79	72	1
144	Pineapple	<i>Ananas comosus</i> Linn.	81-90	85	54	0.4
145	Plantain, green	<i>Musa paradisiaca</i> Linn.	61-84	70	103	1
146	Pomegranate, pulp	<i>Punica granatum</i> Linn.	76-84	81	68	1
147	Prickly pear	<i>Opuntia</i> spp. ..	—	81	(56)	1
148	Small fruit (based on blackberry, raspberry, strawberry)	—	85	26	1
149	Stone fruit (based on peach, apricot, plum)—see also Item 142	<i>Prunus</i> spp. ..	—	90	35	0.7
150	Tamarind	<i>Tamarindus indica</i> Linn.	20-26	21	277	3.5
151	Tomato, with skin	<i>Lycopersicum esculentum</i> Mill.	90-96	92	18	1

Additional Names

- Cantaloupe, *see* Melon, sweet
 Ground cherry, *see* Cape gooseberry
 Honeydew melon, *see* Melon, sweet
 Mandarin orange, *see* Tangerine
 Musk melon, *see* Melon, sweet
 Naartje, *see* Tangerine
 Passion fruit, *see* Grenadilla
 Pawpaw, *see* Papaya.

VII.—STEMS							
152	Maize	<i>Zea mays</i> Linn. ..	73-78	75	58	0.5	
153	Sorghum	<i>Sorghum</i> spp. ..	[73-78] ₁₅₁	[75] ₁₅₁	[58] ₁₅₁	[0.5] ₁₅₁	
154	Sugar cane	<i>Saccharum officinarum</i> Linn.	70-87	82	58	0.4	

For significance of asterisks and other abbreviations used throughout Tables, see p. 6.

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Aneurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
ø	17	15	1	2,000-c	—	—	—	4(l)	—	143
ø	13	20	1	200-c	0.09	0.02	0.1	30(m)	40	144
0.3	24	7	0.5	100-c	[0.05] ₁₂₂	[0.06] ₁₂₂	[0.5] ₁₂₂	[10] ₁₂₂	[33] ₁₂₂	145
ø	16	10	0.7	—	—	0.01	—	10	44	146
ø	13	26	0.3	—	—	—	—	(15) —	54	147
ø	5.6	40	1	100-c	0.02	0.02	0.16	30(n)	1	148
ø	8	33	0.3	670-c(o)	0.02	0.04	0.55	9	9	149
1	63.5	170	11	100-c	—	—	0.10	10	—	150
ø	3.5	9	0.4	1,200-c	0.05	0.04	0.70	25	2	151

Footnotes

- (a) 100–850 I.U. (carotene), depending on variety.
- (b) 7–37 mg., depending on variety.
- (c) 30–700 I.U. (carotene), depending on variety.
- (d) Rind only.
- (e) 10–450 mg., depending on variety and ripeness.
- (f) Skins and seeds.
- (g) May contain as much as 2,500–5,000 I.U. (carotene), depending on variety and ripeness.
- (h) May contain as much as 100 mg.
- (i) May contain as much as 32,000 I.U. (carotene), depending on variety.
- (j) May contain as much as 100 mg.
- (k) 30–120 mg., depending on variety.
- (l) Native American varieties (*D. Virginiana*), 100 mg.
- (m) May contain as much as 60 mg.
- (n) Strawberries may contain two or three times this amount.
- (o) For apricots take vitamin A = 2,500 I.U. (carotene), and for yellow peaches 1,200 I.U. (carotene).

ø	14(a) [ø] ₁₂₂	25 [25] ₁₂₂	1 [1] ₁₂₂	0 [0] ₁₂₂	ø [ø] ₁₂₂	—	—	ø [ø] ₁₂₂	— (b) — (b) — (b)	152 153 154
ø	14	10	1	0	ø	—	—	ø		

Footnote

- (a) The sugars present as such may be as high as 10 per cent. in one sample and as low as 5 per cent. or even less in another.
- (b) Commonly chewed to extract the juice; the material ejected after chewing includes saliva and therefore cannot be used as a basis for calculating the edible portion. Fibre in cane, by analysis = 10–15 per cent.

REPRESENTATIVE

No.	PRODUCT	SCIENTIFIC NAME				
			Water Content		Selected Value	Calories
			Range	g.		
	VIII.—FATS and OILS			g.	g.	g.
155	Butter	—	15	745	0.5
156	Ghee (clarified butter)	—	8	828	0
157	Lard and other animal fats and oils. (b)	—	1	891	0
158	Margarine	—	14	768	0
159	Red palm oil	Elaeis guineensis Jacq.	—	0	900	0
160	Vegetable oils, other, taken as free from water.	—	0	900	0

*Additional Names*Clarified butter, *see* Ghee.

	IX.—ANIMAL PRODUCTS (except FATS)						
	A. Fish and Fish Products (including Molluscs and Crustacea)						
	Fish						
161	Dressed, fresh*	—	78	98	19	
162	Lake, mixed, fresh. (a)	63-82	75	95	17	
163	Lake, mixed, dried. (b)	10-55	30	272	50	
164	Freshwater, lean, fresh fillet	—	77	94	18	
165	White, sea, fresh fillet	—	80	73	16	
166	Fat, fresh	—	69	163	16	
167	Cod, salt wet	—	55	93	22	
168	Salt, hard dry	—	40	182	42	
169	Shell, fresh	—	74	93	18	
170	Canned sardines, in oil	—	50	305	20	
171	Canned salmon, not in oil	—	67	170	20	
172	Sea slug (bêche de mer), soaked	Stichopus japonicus Selenka	—	76	94	22	
173	Snail, river	Vivipara quadrata Benson	—	78	77	12	
174	Turtle	Trionyx chinensis T. et S.	—	80	79	16	

*Additional Names*Bêche de mer, *see* Sea slug.*Footnotes*(a) Values for protein, fat, carbohydrate and minerals are provisional only, calculated from those for dried fish, *q.v.*

(b) Unpublished data on samples from Lake Nyasa.

(c) Large hard bones only, fins, scales and sometimes guts.

	B. Insects						
175	Larval—Caterpillars, fresh	75-85	80	83	14	
176	Mature Termites (white ants), winged, fresh	—	75	148	10	
177	Various, including locusts	60-78	70	134	20	

*Additional Names*White ants, *see* Termites.

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE per cent. of Food as purchased	No.
				A	Ancurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
82.5	0	15	0.2	2,700	0	0	0	0	0	155
92	0	0	0	2,500 ^(a)	0	0	0	0	0	156
99	0	0	0.1	0 ^(b)	0	0	0	0	0	157
85.3	0	4	0	— ^(c)	—	—	—	0	0	158
100	0	0	0	40,000-80,000-c	0	0	0	0	0	159
100	0	0	0	0	0	0	0	0	0	160

Footnotes

(a) May be much lower, depending on the method of processing.

(b) Fish liver oils contain variable amounts of vitamins A and D.

(c) No vitamin A unless artificially added.

+	2.5	—	20	1.0	—	0.06	0.15	3	—	33	161
3 ^(d)	—	—	700 ^(e)	3 ^(f)	—	[0.05] ₁₆₄	[0.13] ₁₆₄	[3] ₁₆₄	[σ] ₁₆₄	25 ^(c)	162
8 ^(d)	—	—	2,000 ^(e)	8 ^(f)	—	[0.06] ₁₆₄	[0.31] ₁₆₄	[4.6] ₁₆₄	[σ] ₁₆₄	15 ^(g)	163
2.5	—	—	20	0.7	—	0.05	0.13	3	—	—	164
q 1	—	—	20	0.7	—	0.05	0.13	3	—	—	165
11	—	—	20	0.7	150	0.05	0.13	3	—	—	166
0.5	—	—	35	1.5	—	0.05	0.36	5	—	45	167
1.5	—	—	65	2.5	—	0.06	0.31	4.6	—	35	168
1.5	2	—	100	5	200	0.05	0.10	4	—	—	169
25	—	—	400	3	250	0.02	0.20	4	0	—	170
10	—	—	100	1.0	100	0.02	0.20	6	0	—	171
—	1.4	—	120	1.4	—	—	—	—	—	0	172
1.4	4	—	1,360	20	—	—	—	—	—	53	173
1	1.5	—	110	1.4	—	—	—	—	—	70	174

(d) Variable, depending on species. Whole dried fish of 4 species, range 2-13 per cent., average 9 per cent; flesh and skin only, 16 samples cured by various methods, range 0.8-9 per cent., average 5 per cent.

(e) In view of local cooking methods, allows for a proportion of the bones in the edible portion. Whole dried fish, 4 species, range 2,000-4,400 mg., average 3,000 mg.; flesh and skin only, 16 samples, range 120-1,100 mg., average 500 mg.

(f) Allows for some bone marrow for same reason as above. Whole dried fish, 4 species, range 7-24 mg., average 13 mg.

(g) Large hard bones only, fins and scales. Range found 0-15 per cent.

3	—	20	0.5	—	—	—	—	—	—	—	175
12	—	25	1	—	—	—	—	—	—	—	176
6	—	30	1	—	—	—	—	—	—	—	177

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
	IX.—ANIMAL PRODUCTS —contd.	—	g.	g.		g.
	C. Meat, Meat Products and Eggs	—				
	Bacon	—				
178	Fat back	—	7	817	4
179	Whole side	—	—	449	11
	Beef	—				
180	Whole carcase, lean	—	66	189	18
181	Whole carcase, mod. fat	—	61	253	16
182	Canned, corned	—	53	244	25
	Eggs	—	74	153	13
183	Goat, carcase	—	73	145	16
	Mutton	—				
185	Average carcase	—	63	241	13
186	Lean carcase	—	73	145	16
	Offal	—				
187	Heart	—	71	163	16
188	Kidney	—	75	127	16
189	Liver	—	70	142	17
	Pork	—				
190	Fresh, fat	—	28	599	8
191	Fresh, lean	—	35	526	10
192	Salt, fat	—	—	781	4
193	Poultry, plucked and drawn	—	73	139	19
194	Rabbit, skinned and gutted	—	67	134	20
	D. Milk and Milk Products	—				
	Cheese	—				
195	Cow, whole milk, hard	—	37	415	25
196	Cow, skimmed milk, soft	—	73	108	20
197	Curds, including whey	—	—	—	—
	Milk, fresh	—				
198	Whole, cow, 2·5 per cent. fat	—	89	53	3
199	Whole, cow, 3·5 per cent. fat	—	88	62	3·1
200	Whole, goat	—	86	71	4·4
201	Whole, sheep	—	81	104	6·5
202	Skimmed, cow	—	91	34	3·2
	Milk, processed	—				
203	Whole, condensed unsweetened (evaporated)	—	74	140	7
204	Whole, condensed sweetened	—	28	314	8·2
205	Whole, powder	—	5	485	25·6
206	Skimmed, condensed sweetened	—	29	266	9·5
207	Skimmed, powder	—	5	360	35·6

For significance of asterisks and other abbreviations used throughout Tables, see p. 6.

VALUES PER 100 gm. EDIBLE PORTION

Fat g.	Carbo- hydrate g.	Calcium mg.	Iron mg.	Vitamins					WASTE per cent. of Food as purchased	No.
				A I.U.	Aneurin mg.	Riboflavin mg.	Nicotinic Acid mg.	Ascorbic Acid mg.		
89	0	5	1	—	0.21	0.10	(0.6)	0	4	178
45	0	11	2	—	0.57	0.26	(1.8)	0	12	179
13	0	12	4	(40)	0.09	0.25	5	0	23	180
21	0	11	4	(40)	0.09	0.25	5	0	20	181
16	0	18	(4)	(30)	0.01	0.20	2.5	0	0	182
11	0.9	55	2.8	1,000	0.14	0.34	0	0	12	183
9	0	11	2.7	0	0.20	0.27	5	0	26	184
21	0	10	2.4	0	0.19	0.24	5	0	16	185
9	0	11	2.7	0	0.20	0.27	5	0	26	186
11	0	10	8	200	0.6	0.9	(7)	0	—	187
7	0	3	10	1,000	0.30	2	7	—	—	188
6	5	11	10	27,000	0.30	2.50	15	30	—	189
63 ^(a)	0	10	1.7	0	0.98	0.26	5	0	13	190
54 ^(b)	0	11	1.8	0	1.04	0.28	5	0	18	191
85	0	2	0.6	—	(0.18)	0.04	0.9	0	4	192
7	0	15	3	0	0.15	0.15	6	0	33	193
6	0	10	2.4	0	0.04	0.12	5	0	18	194
35	0	850	1.0	1,400	0.04	0.50	0.03	0	0	195
2	2.5	350	(0.2)	(20)	0	(0.10)	0	0	0	196
		Take as corr	esponding milk.							197
2.5	4.6	120	0.1	100 ^(c)	0.04	0.15	0.1	1	0	198
3.5	4.6	120	0.1	140 ^(c)	0.04	0.15	0.1	1	0	199
4.1	4.2	140	0.7	80	0.05	0.10	4	1	0	200
6.7	4.5	140	0.7	80	0.05	0.10	4	1	0	201
0.2	4.9	124	0.1	0	0.05	0.21	0.1	1	0	202
8	10	240	0.2	400	0.10	0.36	0.20	1	0	203
9.2	49.6	290	0.2	400	0.10	0.36	0.20	1	0	204
26.7	35.6	900	0.8	1,400	0.30	1.15	0.70	6	0	205
0.5	55.8	320	0.3	20	0.10	0.36	0.20	1	0	206
1	52	1,240	1.0	60	0.39	1.60	1.10	7	0	207

Footnotes

(a) 58 per cent. when kidney and visceral fat removed.

(b) 50 per cent. when kidney and visceral fat removed.

(c) For stall-fed cattle take 70 I.U.

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content		Calories	Protein
			Range	Selected Value		
	X.—CONDIMENTS AND SPICES		g.	g.		g.
208	Asafoetida	Ferula narthea ..	—	16	(297)	4
209	Cardamom	Elettaria cardamomum Maton.	—	20	(226)	10
	Chillies Sweet, see under Vegetables, leafy, green and yellow					
210	Very hot, fresh	Capsicum frutescens Linn.	—	69	85	4
211	Cloves	Eugenia aromatica Linn.	—	23	(293)	5
212	Coriander	Coriandrum sativum Linn.	—	11	(288)	14
213	Cumin	Cuminum cyminum Linn.	—	12	(359)	19
214	Fenugreek, seed	Trigonella foenum-graecum Linn.	—	14	(334)	26
215	Garlic	Allium sativum Linn.	—	63	(140)	6
216	Ginger, root, fresh	Zingiber officinalis Rosc.	—	81	(65)	2
217	Mace	Myristica fragrans Houtt.	—	16	(432)	6
218	Mustard, seed	Brassica nigra (Linn.) Koch.	—	8	(544)	22
219	Nutmeg	Myristica fragrans Houtt.	—	14	(468)	8
220	Omum	Carum copticum ..	—	9	(378)	15
221	Pepper, dry	Piper nigrum Linn.	—	13	(311)	12
222	Tamarind	Tamarindus indica Linn.	20-26	21	277	3.5
223	Turmeric, root, dry	Curcuma longa Linn.	—	14	(341)	4

	XI.—SYRUPS, SUGAR and PRESERVES						
224	Honey	—	23	306	0.4	
225	Jam*	—	29	248	0	
226	Molasses, cane, medium	—	—	(240)	—	
	Sugar						
227	Brown	—	8	400	0	
228	White	—	0	400	0	

For significance of asterisks and other abbreviations used throughout Tables, see p. 6.

VALUES PER 100 gm. EDIBLE PORTION

VALUES PER 100 gm. EDIBLE PORTION										WASTE per cent. of Food as purchased	No.		
Fat	Carbo- hydrate	Calcium	Iron	Vitamins									
				A	Ancurin	Riboflavin	Nicotinic Acid	Ascorbic Acid					
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.					
1	68	700	22	—	—	—	—	0	—	208			
2	42	130	5	—	—	—	—	0	—	209			
5	6	—	—	450-c	—	—	—	50	—	210			
9	48	740	5	—	—	—	—	0	—	211			
16	22	630	18	1,570-c	—	—	1.1	8	—	212			
15	37	1,080	3	870-c	—	—	2.6	3	—	213			
6	44	160	14	8	—	—	1.1	0	—	214			
8	29	25	1.5	0	—	—	0.4	13	—	215			
1	12	20	2.5	8	—	—	0.6	6	—	216			
24	48	180	12.5	—	—	—	—	0	—	217			
40	24	490	18	270-c	—	—	4.0	8	—	218			
36	28	120	4.5	8	—	—	—	0	—	219			
18	39	1,420	14.5	—	—	—	—	—	—	220			
7	50	460	17	—	—	—	1.4	—	—	221			
1	68.5	170	11	100-c	—	—	—	10	—	222			
5	70	150	18	—	—	—	2.3	0	—	223			

<i>s</i>	76	5	0·4	—	<i>s</i>	0·06	0·11	2	0	224
<i>s</i>	62	12	—	—	<i>s</i>	(0·20)	1·5	—	—	225
—	60	273	6·7	—	0·05	—	—	—	—	226
0	100	75	1	—	0	0	0	0	0	227
0	100	0	0	—	0	0	0	0	0	228

No.	PRODUCT	SCIENTIFIC NAME	REPRESENTATIVE			
			Water Content	Calories	Alcohol	Protein
XII.—BEVERAGES						
229	Fermented (a) Beer, African maize (b)	—	23-65	1-7	1
230	European (b)	—	45-56	4-5-6-0	0-3
Unfermented						
231	Cocoa powder	Theobroma cacao Linn.	3	429	—
232	Coffee, ground (d)	Coffea spp.	42	—	4-5
233	Tea leaves (d)	Thea sinensis	—	40	10

For significance of asterisks and other abbreviations used throughout Tables, see p. 6.

VALUES PER 100 gm. EDIBLE PORTION

Fat	Carbo-hydrate	Calcium	Iron	Vitamins					WASTE Per cent. of Food as purchased	No
				A	Ancurin	Riboflavin	Nicotinic Acid	Ascorbic Acid		
g.	g.	mg.	mg.	I.U.	mg.	mg.	mg.	mg.		
o	3(c)	4	0.2	ø	0.03	0.03	0.58	1	—	229
o	3.3	12	0.1	—	—	0.05-0.17	—	—	—	230
25	31	51	14	150-c	0.12	0.27	1	—	—	231
o	6	58	ø	—	—	0.1	13.2	—	—	232
—	—	30	—	—	—	1	6.1	—	—	233

Footnotes

(a) Other fermented beverages are consumed, e.g. other cereal beers, palm wine and coconut toddy; the alcoholic content ranges from 1-7 per cent., depending on length of time of fermentation.

(b) g. per 100 c.c.

(c) "Available" carbohydrate in terms of starch.

(d) Diffusible material per 100 g., extracted by hot water in 5-10 minutes.

ALTERNATIVE BOTANICAL NAMES

Name in Tables	Alternative Name(s)
Ananas comosus (Linn.) Merr.	Ananas sativus Schult. f.
Artocarpus communis Forst.	Artocarpus incisus Linn. f.
Artocarpus integer (Thunb.) Merr.	Artocarpus integrifolius Linn. f.
Cajanus cajan (Linn.) Millsp.	Cajanus indicus Spreng.
Citrus aurantiifolia (Christm.) Swingle	Citrus medica Linn. var. acida hort.
Citrus sinensis Osbeck	Citrus aurantium Linn. var. sinensis Linn.
Cola acuminata Schott. & Endl.	Sterculia acuminata Beauv.
Fagopyrum esculentum Moench	Fagopyrum sagittatum Glib. ; Polygonum fagopyrum Linn.
Glycine max. (Linn.) Merr.	Glycine soya Sieb. & Zucc. ; Soja hispida Maxim.
Manihot utilissima Pohl	Includes the doubtful species M. palmata Muell.-Arg.
Mucuna utilis hort.	Mucuna deeringiana (Bort) Holland ; Stizolobium deeringianum Bort.
Pennisetum typhoides Stapf & Hubbard.. ..	Pennisetum spicatum Koern. ; Pennisetum typhoideum L. C. Rich.
Persea americana Mill.	Persea gratissima Gaertn. ; Persea drymifolia Schlecht. & Cham.
Phaseolus aureus Roxb.	Phaseolus radiatus Linn.
Phaseolus lunatus Linn.	Phaseolus inamoenus Blanco.
Physalis peruviana Linn.	Physalis edulis Sims.
Pisum sativum Linn.	Pisum arvense Linn.
Prunus persica Stokes	Amygdalus persica Linn.
Sesamum indicum Linn.	Sesamum orientale Linn.
Triticum vulgare Host Cultigen.	Triticum sativum Lam. ; Triticum aestivum Linn. ; Triticum hybernum Linn. ; Triticum pilosum Dalz. & Gibbs.
Vigna unguiculata (Linn.) Walp.	Vigna sinensis Endl. ; Vigna catjang Walp.

REFERENCES

- ACCESSORY FOOD FACTORS COMMITTEE OF MEDICAL RESEARCH COUNCIL AND LISTER INSTITUTE (1945) : The Nutritive Values of War-Time Foods. *M.R.C. War Memorandum No. 14.* London : H.M. Stationery Office.
- ADAMS, G., and SMITH, S. L. (1944) : The vitamin content and the preservation of foods. *U.S. Dep. Agric. misc. Pub. No. 536.*
- ADOLPH, W. H. (1928) : Analyses of Chinese food materials. *Philipp. J. Sci., 30, 3.*
- ANDREWS, J. S., et al. (1942) : The riboflavin content of cereal grain and bread. *Cereal Chem., 19, 1.*
- ARNOLD, A., et al. (1941) : Chemical determination of nicotinic acid. *Indust. & Eng. Chem. (Anal. Ed.), 33, 62.*
- ARNOLD, H. C. (1940) : Soya Beans. *Rhod. agric. J., 38, 10.*
- ATWATER, W. O. and BRYANT, A. P. (1906) : The chemical composition of American food materials. *U.S. Dep. Agric. Bull. No. 28.*
- AYKROYD, W. R. (1938) : The nutritive value of Indian foods and the planning of satisfactory diets. *Gout. of India, Hlth. Bull. No. 23, 2nd ed.* Delhi.
- , W. R., et al. (1940) : The rice problem in India. *Indian med. Res. Mem. 32.*
- , W. R., and SWAMINATHAN, M. (1940) : The nicotinic acid content of cereals and pellagra. *Indian J. med. Res., 27, 667.*
- AXTMAYER, J. H. and COOK, D. H. (1942) : *Manual de Bromatologia.* Washington.
- BACHARACH, A. L. (1940) : Nutrient analyses. *Food, 10, 28.*
- , A. L. (1941) : The distribution of nicotinic acid in human and animal foods. *Nutr. Absr. Rev., 10, 459.*
- , A. L. (1942) : The Nation's Food. V—Meat as Food. 4. The mineral and vitamin content of meat. *Chem. & Indust., 61, 404.*
- BAILEY, M. I., and THOMAS, A. W. (1942) : The thiamine and riboflavin content of citrus fruits. *J. Nutrit., 24, 85.*
- BAKER, A. Z., and WRIGHT, M. D. (1935) : The vitamin B₁ content of foods. *Biochem. J., 29, 1802.*
- BARTON-WRIGHT, E. C. (1944) : The microbiological assay of nicotinic acid in cereals and other products. *Biochem. J., 38, 314.*
- BASU, K. P., et al. (1942) : Studies in human nutrition : IV—Availability of calcium ingested in the process of chewing betel-leaves with lime. V—The bones of small fish as a source of nutritionally available calcium and phosphorus. *Indian J. med. Res., 30, 309 ; 30, 417.*
- , K. P., and GHOSH, D. (1943) : Availability of calcium in lady's finger (*Hibiscus esculentus*), cabbage (*Brassica oleracea capitata*), drumstick (*Moringa olifera*) and amaranth tender (*Amaranthus gangeticus*). I and II. *Indian J. med. Res., 31, 29, 37.*
- BAUERNFEIND, J. C., et al. (1942) : Growth stimulants in the microbiological assay for riboflavin and pantothenic acid. *Indust. & Eng. Chem., 14, 667.*
- BECKLEY, V. A., and NOTLEY, V. E. (1943) : Ascorbic acid content of sweet peppers. *J. Soc. chem. Ind., Lond., 62, 14.*
- BEESON, K. C. (1941) : The mineral composition of crops, with particular reference to the soils in which they were grown. *U.S. Dep. Agric. misc. Pub. No. 369.*
- BERRYMAN, G. H., and CHATFIELD, C. (1943) : A short method of calculating the nutritive value of diets. *J. Nutrit., 25, 1.*
- BHAGVAT, K., and NARASINGA RAO, K. K. P. (1942) : Vitamin C content of dry Bengal gram (*Cicer arietinum*). *Indian J. med. Res., 30, 505.*
- BILLINGS, F. L., et al. (1941) : The riboflavin content of fish products. *J. Nutrit., 22, 425.*
- BISWAS, H. G., and DAS, K. L. (1939-40) : Vitamin C content of the chillies, onion and garlic, both in the raw state and when boiled with water. *Indian J. med. Res., 27, 136.*
- BLACK, M. A. (1939) : Soya beans. *N.Z. J. Sci. Tech., 21, 1A.*
- BOMBAY DEPT. OF AGRIC. (1936) : Mineral constituents of food grains, vegetables and cattle feed. *Bull. No. 179.*
- BOOHER, L. E., and HARTZLER, E. R. (1939) : The vitamin B₁ content of foods in terms of crystalline thiamin. *U.S. Dep. Agric. tech. Bull. No. 707.*
- , L. E., HARTZLER, E. R., and HEWSTON, E. M. (1942) : A compilation of the vitamin values of foods in relation to processing and other variants. *U.S. Dep. Agric. Circ. No. 638.*
- , L. E., and MARSH, R. L. (1941) : The vitamin A values of 128 foods as determined by the rat growth method. *U.S. Dep. Agric. tech. Bull. No. 802.*
- BOYER, W. W., and VILLIERS, D. J. R. (1942) : The vitamin C content of guavas. *Fmg. S. Afr., 17.*
- BRIDGES, M. A. (1935) : Food and beverage analyses.
- BRITISH HONDURAS, GOVT. OF. (1937) : Report of the Committee on Nutrition. London (Crown Agents).

- BUCK, J. L. (1937) : Land utilisation in China. Shanghai and Oxford.
- BURKILL, I. H. (1935) : A dictionary of the economic products of the Malay Peninsula. London (Crown Agents).
- BURRELL, R. C., and WOLFE, A. C. (1940) : The chemical composition of 5 varieties of soybeans. *Food Res.*, 5, 109.
- CEYLON, GOVT. OF. (1937) : Further report on Nutrition in Ceylon. *Sessional Paper No. 29.*
- CHATFIELD, C., and ADAMS, G. (1931) : Proximate composition of fresh vegetables. *U.S. Dep. Agric. Circ. No. 146.*
- , —. (1940) : Proximate composition of American food materials. *U.S. Dep. Agric. Circ. No. 549.*
- , C., and McLAUGHLIN, L. I. (1928) : Proximate composition of fresh fruit. *U.S. Dep. Agric. Circ. No. 50.*
- CHI, Y. F., and READ, B. (1935) : The vitamin content of Chinese foods and drugs. *Chin. J. Physiol.*, 9, 47.
- CHUNG, H. L., and RIPPERTON, J. C. (1929) : Utilisation and composition of oriental vegetables in Hawaii. *Hawaii agric. exp. Sta. Bull.* No. 60.
- COLLIER, B. C., et al. (1941) : Ascorbic acid content of some maritime province foods. *Proc. N.S. Inst. Sci.*, 20, 121.
- COLONIAL FOOD YEAST, LTD. (1944) : Food Yeast—A venture in practical nutrition. London.
- COMBINED U.K./U.S.A./U.N.N.R.A. WORKING PARTY ON EUROPEAN FOOD SUPPLIES. (1945) : Nutrient Values (*unpublished*).
- Comparison of tables of food values of one pound of specified food materials. (1943) : Canadian, American and British (*unpublished*).
- CONNER, R. T., and STROUB, G. J. (1941) : Combined determination of riboflavin and thiamin in food products. *Indust. & Eng. Chem.*, 33, 389.
- DALZIEL, J. M. (1937) : The useful plants of west tropical Africa. London.
- DANIEL, E. P., and MUNSELL, H. E. (1937) : The vitamin content of foods. *U.S. Dep. Agric. misc. Pub. No. 275.*
- DANN, W. J., and HANDLER, P. (1942) : Nicotinic acid content of meat. *J. Nutrit.*, 24, 153.
- DARBY, W. J., and DAY, P. L. (1938) : The riboflavin content of meats. *J. Nutrit.* 16, 209.
- DONATE, W. F. (1934) : De voedingswaarde van het inheemsche menu te Koetownangoen. *Landbouw*, Nos. 4 and 5.
- , W. F., KOOHLHAAS, D. R., and VAN VEEN, A. G. (1935) : Voedingstabellen. *Geneesh. Tijdschr. Ned.-Ind.*, 75, 426.
- , W. F., and SPRUYT, J. P. (1936) : Het vitamin B₁ gehalte en de voedingswaarde van eenige belangrijke Indische voedingsmiddelen. *Geneesh. Tijdschr. Ned.-Ind.*, 78, 642.
- DONOVAN, W. O'B., and COCHRANE, E. (1943) : West Indian foods. Grenada.
- EDDY, W. H. (1933) : The nutritive value of the banana. New York.
- FELLERS, C. R., et al. (1940) : The vitamin B₁ and B₂ content of vegetables as influenced by quick freezing and canning. *Food Res.*, 5, 495.
- FENTON, F., and TRESSLER, D. K. (1938) : Losses of vitamin C during the cooking of certain vegetables. *J. Home Econ.*, 30, 717.
- FIXSEN, M. A. B. (1938) : The vitamin content of human foods as affected by processes of cooking and canning. *Nutr. Abstr. Rev.*, 8, 281.
- , M. A. B., and ROSCOE, M. H. (1938) : Tables of the vitamin content of human and animal foods. *Nutr. Abstr. Rev.*, 7, 823.
- FLOYD, W. W., and FRAPS, G. S. (1939) : The vitamin C content of some Texas fruits and vegetables. *Food Res.*, 4, 87.
- FOX, F. W. (1938) : Notes on methods of preparation and nutritive value of certain kaffir beers. *J.S. Afr. chem. Inst.*, 21, 21.
- , F. W., and GOLBERG, L. (1944) : South African Food Tables. *S. Afr. Inst. med. Res.*, No. 46, 9, 123.
- , F. W., and STONE, W. (1938) : The antiscorbutic value of kaffir beer. *S. Afr. J. med. Sci.*, 8, 7.
- , F. W., and WILSON, C. (1936) : Lucerne as a food for human consumption. *S. Afr. Inst. for Med. Res., Laboratory Rept.* No. 3 of Enquiry into Native Diets.
- FRENCH, M. H. (1943) : The composition and nutritive values of Tanganyika feeding stuffs. *E. Afr. Agric. J.*, 8, 3.
- FUJITA, A., and AJISAKA, M. (1941) : Über die Verteilung des Provitamins A I—Verteilung des Carotins in Pflanzen. *Biochem. Z.*, 308, 430.
- GOLBERG, L., and LEVY, L. F. (1942) : The antiscorbutic value of guavas. *J.S. Afr. chem. Inst.*, 25, 1.
- , L., and THORP, J. M. (1942) : Survey of vitamins in African foodstuffs. I—The thiamin content of wheat. *S. Afr. J. med. Sci.*, 7, 95.

- GOMOLYAKO, L. G. (1937) : Antiscorbutic vitamin in some fruits and vegetables of the North Caucasus. *Bull. appl. Bot. Gen. & Plant Breeding* (U.S.S.R.), Suppl. 84, 200.
- GUTIERREZ, M. (1940) : The vitamin B₁ content of Philippine foods. *Acta med. Philipp.*, 1, 415.
- VON HAHN, F. U. (1931) : Vitaminstudien III. Der Vitamingehalt des Obstes. *Z. Untersuch. Lebensmitt.*, 61, 369.
- HALE, E. B., et al. (1942) : Nicotinic acid in feeds. *J. biol. Chem.*, 148, 2.
- , E.B., et al. (1942) : The chemical determination of nicotinic acid in plant materials. *J. biol. Chem.*, 148, 556.
- HALVERSON, J. O., and SHERWOOD, F. W. (1940) : The vitamin A activity and the vitamin B₁ content of soybeans and cowpeas. *J. agric. Res.*, 60, 2.
- HAND, D. B., and SHARP, P. F. (1939) : The riboflavin content of cow's milk. *J. Dairy Sci.*, 22, 779.
- HARDY, G., and RICHEZ, C. (Fils). (1933) : *L'Alimentation indigène dans les colonies françaises-Paris*.
- HARRIS, L. J. (1938) : Vitamins and vitamin deficiencies, Vol. I. London,
- HARRIS, P. L., and GRANGE, L. P. (1937) : Vitamin studies on bananas. I—The vitamin A, B₁, and C contents of ripe bananas. *Food Res.*, 2, 4.
- HARTLEY, K. T., and ROSS, S. D. (1938) : Feeding of Nigerian stock. *Papers of the Third W. African Agricultural Conference*, p. 85.
- HODSON, A. Z. (1940) : The riboflavin content of some common vegetables and fruits. *Food Res.*, 5, 395.
- HOOVER, A. A. (1941) : The mineral analyses of some local yams and vegetables. One of a series of articles under the general title, "The analyses of Ceylon foodstuffs". *Trop. Agriculturist*, 97, 4, 185.
- HORVATH, A. A. (1938) : The soybean industry. London.
- HSU, P., and ADOLPH, W. H. (1935) : On the distribution of calcium in leafy vegetables. *Chin. med. J.*, 49, 325.
- IEDE, A. T., and SCHUETTE, H. A. (1941) : Thiamine, nicotinic acid, riboflavin and pantothenic acid in rye and its milled products. *J. Nutrit.*, 22, 529.
- IMPERIAL BUREAU OF ANIMAL NUTRITION, ABERDEEN. (1936) : Composition of some African foodstuffs. *Tech. Commun.* No. 6.
- (1938) : Table of composition of foods. *Tech. Commun.* No. 10.
- IMPERIAL COUNCIL OF AGRICULTURAL RESEARCH, INDIA. (1938) : The nutritive value of Indian cattle foods. *Misc. Bull.* No. 25.
- IMPERIAL INSTITUTE, LONDON. *Bulletins* (various).
- INTERNATIONAL INSTITUTE OF AFRICAN LANGUAGES AND CULTURES. (1937) : The food and nutrition of African natives. *Memo. No. 13*.
- ITTNER, N. R., and HUGHES, E. H. (1941) : The riboflavin content of pork muscle. *Food Res.*, 6, 3.
- JACQUOT, R., and NATAF, B. (1936) : Le manioc. Actualités scientifiques et industrielles 364, Nutrition. Faculté des Sciences, Strasbourg.
- JOACHIM, A. W. R., et al. (1938) : The analysis of Ceylon foodstuffs, I-V. *Trop. Agriculturist*, 90, 1.
- KERLY, M. (1944) : Riboflavin content of canteen meals. *Biochem. J.*, 38, 428.
- KHORANA, M. L., et al. (1943) : Investigations on the food value of fish and other marine products. II—The protein and mineral contents. *Indian J. med. Res.* 31, 25.
- KIRK, M. M., and TRESSLER, D. K. (1941) : The ascorbic acid content of pigmented fruits and vegetables and their fruit juices. *Food Res.*, 6, 4.
- KODICEK, E. (1940) : Estimation of nicotinic acid in animal tissues, blood and certain foodstuffs. *Biochem. J.*, 34, 5.
- KOKFORD, H. G., POSTMUS, S., and VAN VEEN, A. G. (1939) : De Voedingswaarde van Voedingsmiddelen. *Landbouw*, 15, 193.
- KOHMANN, E. F., et al. (1926) : Vitamins in canned food. *Industr. Eng.*, 18, 302.
- KÖNIG, J. (1903 and 1923) : Chemie der menschlichen Nahrungs- und Genussmittel. Chemie der menschlichen Nahrungs- und Genussmittel (Nachtrag).
- KWOH, T. H., et al. (1941) : The carotene content of some Chinese fruits and vegetables. *J. Chin. chem. Soc.*, 8, 54.
- LANFORD, C. S., et al. (1941) : The riboflavin content of some typical fruits. *J. Nutrit.*, 21, 175.
- LEGGETT, E. L., and ROBINSON, K. W. (1939) : The chemical estimation of vitamin C, with analyses made upon some Queensland products. *Med. J. Aust.*, 2, 241.
- LEITCH, J. N. (1930) : Dietetics in warm climates. London.
- LEONG, P. C. (1940) : Nicotinic acid content of foods. *J. Malaya Br. Brit. med. Ass.*, 4, 3.
- , P. C. (1939) : Vitamin A content of Malayan bananas. *J. Malaya Br. Brit. med. Ass.*, 3, 2.
- LEVY, L. F. (1937) : The antiscorbutic value of S. African mangoes. *Fmg. S. Afr.*, 12.
- , L. F., and FOX, F. W. (1938) : The antiscorbutic value of some S. African foodstuffs. *S. Afr. med. J.*, 9, 181.

- LEVY, L. F., WEINTROUB, D. and FOX, F. W. (1936) : The food value of some common edible leaves. *S. Afr. med. J.*, **10**, 699.
- LOVERN, J. A. (1943) : The nation's food. VI—Fish as food. 3. The vitamin and mineral content of fish. *Chem. & Ind.*, **62**, 328.
- LUND, H. (1942) : Undersøgelser over karotin i frisk og torret lucerne. *Tidsskr. Planteavl.*, **46**, 275.
- LUNDE, G., KRINGSTAD, H., and OLSEN, A. (1938) : Untersuchungen über den gehalt an Vitamine B₁ und B₂. *Skr. norske Videnskakad.* **7**.
- MARSTON, H. R., and DAWBARN, M. C. (1941) : Food composition tables. *Austral. Coun. sci. indust. Res. Pamphlet No. 107*.
- MATAGRIN, A. (1939) : Le Soja.
- MATHIESON, E. (1938) : Vitamin C innholdet i hermetske baer, frukt og grønnsaker. *Tidsskr. Hermetikhindustr.*, **24**, 410.
- MCCANCE, R. A., and SHACKLETON, L. R. B. (1936) : The ionisable iron in foods. *Biochem. J.*, **30**, 582.
- , R. A., and SHIPP, H. L. (1933) : The chemistry of flesh foods and their losses on cooking. *Spec. Rep. Ser. Med. Res. Coun., Lond.*, No. 187.
- , R. A., and WIDDOWSON, E. M. (1935) : Phytin in human nutrition. *Biochem. J.*, **29**, 2694.
- , R. A., WIDDOWSON, E. M., and SHACKLETON, L. R. B. (1936) : The nutritive value of fruits, vegetables and nuts. *Spec. Rep. Ser. Med. Res. Coun., Lond.*, No. 213.
- , R. A., and WIDDOWSON, E. M. (1942) : The chemical composition of foods. *Spec. Rep. Ser. Med. Res. Coun., Lond.*, No. 235.
- MCCRISON, R., and MOORE, D. F. (1943) : Food. London.
- MCINTIRE, J. M., et al. (1941) : Nicotinic acid in meat products. *J. Nutrit.*, **22**, 535.
- MCVICAR, R. W., and BERRYMAN, G. H. (1942) : Nicotinic acid in foods. *J. Nutrit.*, **24**, 235.
- MICKELSON, O., et al. (1939) : The distribution of riboflavin in meat and meat foods. *J. Nutrit.*, **18**, 518.
- MILLER, C. D. (1927) : Food values of poi, taro and limu. *Bull. Bishop Mus., Honolulu*, No. 37.
- , C. D. (1933) : Japanese foods commonly used in Hawaii. *Hawaii agric. exp. Sta. Bull.* No. 68.
- , C. D., et al. (1936) : Some fruits of Hawaii. *Hawaii agric. exp. Sta. Bull.* No. 77.
- , C. D., et al. (1944) : Vitamin values of Hawaiian grown fruits and vegetables. *Hawaii agric. exp. Sta. Progress Note No. 36*.
- MITRA, K., and MITRA, H. C. (1943) : The determination by chemical methods of the food values of yet another batch of edibles. *Indian J. med. Res.*, **31**, 41.
- MORRIS, J. P., and OLIVEIRO, C. J. (1933) : Calcium in tropical foods. *Malay. med. J.*, **8**, 236.
- , J. P., and ROSEDALE, J. L. (1935) : The mineral content of some tropical foods. *Malay. med. J.*, **10**, 1.
- MOTTRAM, V. H., and RADLOFF, E. M. (1937) : Food tables. London.
- MUNSELL, H. E. (1940) : Vitamins and their occurrence in foods. *Milbank mem. Fd. quart. Bull.*, No. 18, 4.
- , H. E. (1942) : The riboflavin content of some common foods. *Food Res.*, **7**, 2.
- , H. E. (1943) : Vitamins and their occurrence in foods. *Puerto Rico J. publ. Hlth. trop. Med.*
- MYBURGH, S. J. (1935) : Carotene content of some S. African foods. *Onderstepoort J. vet. Sci.*, **5**, 475.
- NATIONAL RESEARCH COUNCIL, FOOD AND NUTRITION BOARD, U.S.A. (1945) : Tables of food composition (*unpublished*).
- NEWTON, C. L. (1931) : The vitamin content of turnip greens, collards, cantaloupes and peaches. *Georgia agric. exp. Sta. Bull.* No. 167.
- NICHOLLS, L. (1938) : Tropical nutrition. London.
- NIYOGI, S. P., et al. (1941) : Balanced diets. II—Studies on the nutritive value of fish. *Indian J. med. Res.*, **29**, 279.
- , S. P., et al. (1943) : Studies in fish liver oils. I—The biological assay of vitamins A and D in ghol- (*Sciaena miles*) and mushi- (*Scoliodon sororowah*) liver oils. II—The seasonal variation in the yield and vitamin A content of some fish liver oils. *Indian J. med. Res.*, **31**, 15, 21.
- OLLIVER, M. (1938) : The ascorbic acid content of fruits and vegetables. *Analyst*, **63**, 742.
- , M. (1943) : Ascorbic acid values of fruits and vegetables for dietary surveys. *Chem. & Ind.*, **62**, 146.
- PANNEKOEK-WESTENBURG, S. J. E., NIJHOLT, J. A., and VAN VEEN, A. G. (1940) : Voedingstabellen. *Geneesk. Tijdschr. Ned.-Ind.*, **33**, 80.
- PEIPING UNION MEDICAL COLLEGE HOSPITAL. (1934) : Outlines of diets. Peiping.
- PHILLIPS, J. B. (1934) : The utilisation of the soya bean. *J. Soc. chem. Ind., Lond.*, **53**, 29.
- PIPER, C. V., and MOKSK, W. J. (1923) : The soybean. New York.

- PLIMMER, R. H. A. (1921) : Analyses and energy values of foods. London.
- PYKE, M. (1940) : Observations on the distribution of vitamin B₁ in some plant families. *Biochem. J.*, 34, 330.
- RANGANATHAN, S. (1942) : The vitamin C content of some dehydrated vegetables and fruits and of some fruit juice preparations. *Indian J. med. Res.*, 30, 517.
- RAO, A. M., et al. (1942) : Vitamin B₁ studies. 2—B₁ content of some Indian foodstuffs by the thiochrome method using sensitive photo-electric fluorimeter. *Ann. Biochem. exp. Med.*, 2, 191.
- RAU, Y. V. S., and MURTY, V. V. S. (1942) : The utilisation of calcium from green leafy vegetables. *Ann. Biochem. exp. Med.*, 2, 87.
- RAYMOND, W. D. (1940 and 1941) : The nutritive value of some Tanganyika foods. I—Bananas. II—Cassava. *E. Afr. agr. J.*, 8, 105, 154.
- READ, B. E., LEE, W. Y., and CHENG, J. K. (1937) : Shanghai foods. *Chin. med. Ass. Spec. Rept.* No. 8, 2nd Ed.
- RHODESIAN DIVISION OF CHEMISTRY. (1942) : Analyses of Rhodesian foodstuffs. *Rhod. agric. J.*, 39, 5.
- ROBINSON, K. W., and COURTICE, R. (1940) : Factors affecting the vitamin C (ascorbic acid) content of fruit and vegetables. *Med. J. Aust.*, 1, 786.
- ROSE, M. S. (1937) : A laboratory handbook of dietetics. New York.
- ROSEDALE, J. L. (1935) : Chemical analyses of Malayan foods. Singapore.
- , J. L. (1939) : The nutritive value of rice. *J. Malaya Br. Brit. med. Ass.*, 8, 3.
- RUSSELL, W. C., et al. (1943) : The nicotinic acid content of common fruits and vegetables as prepared for human consumption. *J. Nutrit.*, 25, 275.
- SAFFRY, O. B., et al. (1940) : A biological assay of riboflavin in the liver of cow, calf, sheep, lamb and hog. *J. Nutrit.*, 20, 170.
- SAMPSON, H. C. (1936) : Cultivated crops of the British Empire and the Anglo-Egyptian Sudan (tropical and sub-tropical). *Kew Bull. Additional Series* No. 12.
- SARANGDHAR, P. N. (1942) : Vitamin A content of liver oils of some Indian elasmobranchs. *Indian J. med. Res.*, 30, 553.
- SCHWEIGERT, B. S., et al. (1943) : The retention of vitamins in meats during storage, curing and cooking. *J. Nutrit.*, 28, 75.
- SCOTT, M. A., RANDALL, F. E., and HESSEL, F. H. (1941) : A modification of the Snell-Strong microbiological method for determining riboflavin. *J. biol. Chem.*, 325, 141.
- SCOTT, R. R. (1934) : The preparation of maize flour in Tanganyika Territory. *S. Afr. med. J.*, 8, 399.
- SEN, J. N. (1917) : The composition of some Indian feeding stuffs. *Bull. agric. Res. Inst. Pusa*, No. 70.
- SHERMAN, W. C. (1924) : Food products. New York.
- , W. C. (1933) : Chemistry of food and nutrition. 4th Ed. New York.
- , W. C., and SALMON, W. D. (1939) : The carotene content of soybeans and cowpeas. *Food Res.*, 4, 371.
- SHOURIE, K. L., and SUNDARARAJAN, A. R. (1942) : The effect of milling on the vitamin B₁ (thiamine) and nicotinic acid content of Indian wheat products. *Indian J. med. Res.*, 30, 61.
- SNELL, E. E., and WRIGHT, L. D. (1941) : Microbiological method for the determination of nicotinic acid. *J. biol. Chem.*, 139, 675.
- SULLIVAN, R. A., et al. (1943) : The value of dairy products in nutrition. III—Vitamins in several varieties of cheese. *J. Nutrit.*, 25, 5.
- SWAMINATHAN, M. (1938) : A chemical method for the estimation of nicotinic acid in biological materials. *Indian J. med. Res.*, 28, 427.
- , M. (1941) : Further studies on the cyanogen-bromide method of estimating nicotinic acid in biological materials. *Indian J. med. Res.*, 29, 330.
- , M. (1941) : The effect of washing on the nicotinic acid content of raw and parboiled rice. *Indian J. med. Res.*, 29, 83.
- , M. (1942) : A fluorimetric method for the estimation of riboflavin in foodstuffs. *Indian J. med. Res.*, 30, 30.
- , M. (1942) : A simple procedure for estimating nicotinic acid in biological materials using the cyanogen-bromide aniline reagent. *Indian J. med. Res.*, 30, 401.
- , M. (1942) : Note on the vitamin B₁, riboflavin and nicotinic acid content of dried yeast. *Indian J. med. Res.*, 30, 405.
- , M. (1942) : The effect of washing and cooking on the vitamin B₁ content of raw and parboiled milled rice. *Indian J. med. Res.*, 30, 3.
- , M. (1944) : Nicotinic acid content of Indian foodstuffs. *Indian J. med. Res.*, 32, 39.
- TEPLY, L. J., et al. (1942) : The distribution of nicotinic acid in foods. *J. Nutrit.*, 23, 417.
- THOMAS, J. M., et al. (1942) : The nicotinic acid content of cereal products. *Cereal Chem.*, 19, 2.
- THORPE, E. (1927–1930) : A dictionary of applied chemistry. Revised ed. London.

- THORPE, J. F., and WHITELEY, M. A. (1934-1935) : Thorpe's dictionary of applied chemistry. Supplement. London.
- TRINIDAD, GOVT. OF. (1937) : Food in relation to health. London (Crown Agents).
- UGANDA, GOVT. OF. (1936) : Annual Report of Department of Agriculture, Part II.
- UNITED FRUIT Co. (1928) : Food value of the banana. Boston (U.S.A.).
- (1936 and 1937) : Nutritive and therapeutic values of the banana. Boston (U.S.A.).
- UNITED STATES DEPT. OF AGRICULTURE. (1933) : Report of Hawaii Agricultural Experiment Station.
- (1939) : Food and life. Yearbook.
- (BUREAU OF HOME ECONOMICS). (1943) : Tables of food values (*unpublished*).
- VAJIC, B. (1941) : Der Vitamin C-Gehalt im Weiss- und Rotkraut zu verschiedenen Jahreszeiten. *Biochem. Z.*, 309, 343.
- VAN VEEN, A. G. (1938) : Over cassave-bladeren, een hoogwaardige bladgroen, *Geneesk. Tijdschr. Ned.-Ind.*, 41, 2548.
- , A. G. (1936) : Visch als voedingsmiddel. *Geneesk. Tijdschr. Ned.-Ind.*, 76, 20.
- WAISMAN, H. A., et al. (1940) : The nicotinic acid potency of food materials and certain chemical compounds. *J. Nutrit.*, 19, 483.
- , H. A., and ELVEHJEM, C. A. (1941) : Chemical estimation of nicotinic acid and vitamin B₆. *Indust. & Eng. Chem.*, 33, 221.
- WEGNER, M. I., et al. (1942) : Influence of the method of preparation of sample on microbiological assay for riboflavin. *J. biol. Chem.*, 148, 547.
- WEHMER, C. (1929 and 1931) : Die Pflanzenstoffe. 2nd ed. Vols. I and II. Jena.
- WILSON, H. E. C., and ROY, G. K. (1938) : The flavine and vitamin B₆ content of Indian foodstuffs. *Indian J. med. Res.*, 26, 879.
- WINTON, A. L., and WINTON, K. B. (1935) : The structure and composition of foods. New York.
- WOLFF, E. (1880) : Aschen-analysen. Berlin.
- WOOD, T. B., and WOODMAN, H. E. (1939) : Rations for livestock. *Bull. Minist. Agric., Lond.*, No. 48.
- YOUNG, R. A. (1924) : The dasheen. *U.S. Fmrs' Bull.*, No. 1396.